EXTENSION OF LIE'S ALGORITHM; A POTENTIAL SYMMETRIES CLASSIFICATION OF PDES

By

Patrick Robert Doran-Wu

B. Eng. (Electrical/Electronic) University of Western Australia

M. Sc. (Mathematics) University of Oxford, U.K.

A THESIS SUBMITTED IN PARTIAL FULFILLMENT OF

THE REQUIREMENTS FOR THE DEGREE OF

DOCTOR OF PHILOSOPHY

in

THE FACULTY OF GRADUATE STUDIES

INSTITUTE OF APPLIED MATHEMATICS

DEPARTMENT OF MATHEMATICS

We accept this thesis as conforming

to the required standard

....

....

....

THE UNIVERSITY OF BRITISH COLUMBIA

October 1996

In presenting this thesis in partial fulfillment of the requirements for an advanced degree at the University of British Columbia, I agree that the Library shall make it freely available for reference and study. I further agree that permission for extensive copying of this thesis for scholarly purposes may be granted by the head of my department or by his or her representatives. It is understood that copying or publication of this thesis for financial gain shall not be allowed without my written permission.

Institute of Applied Mathematics
Department of Mathematics
The University of British Columbia
2075 Wesbrook Place
Vancouver, Canada
V6T 1Z1

Date:

Abstract

Symmetries of a system of differential equations are transformations which leave invariant the family of solutions of the system. Infinitesimal Lie symmetries of locally solvable analytic differential equations can be found by using Lie's algorithm. We extend Lie's algorithm to one which can calculate infinitesimal Lie symmetries of analytic systems of differential equations which are not locally solvable.

Local infinitesimal symmetries of differential equations are flows of vector fields which depend on local properties of solutions and have been extensively calculated and applied. In contrast infinitesimal nonlocal symmetries, which are flows of vector fields depending on nonlocal properties of solutions, have only recently been introduced. Using our extension of Lie's symmetry algorithm, we study the infinitesimal nonlocal symmetries of potential type introduced by Bluman, Kumei and Reid. We give verifiable criteria for useful potential systems and give a complete potential symmetries analysis for a class of nonlinear diffusion equations. We also find large classes of higher order scalar and systems of partial differential equations admitting potential symmetries.

Table of Contents

List of Figures

Acknowledgement

I wish to thank my supervisor Dr G.W. Bluman for suggesting that I study potential symmetries of PDEs, for his expert guidance, and his support during my PhD degree. I especially thank him for his speedy reading of my thesis drafts and for his insightful criticisms.

My thanks also go to Dr G.J. Reid for introducing me to Formal Theory of Integrability and to the world of symbolic programming. When I first arrived at U.B.C. he spent many tireless hours with me in the computer lab explaining the various features of his wonderful standard form package. I also thank him for reading my thesis drafts and for his helpful suggestions.

This thesis would not be possible without the constant support and encouragement of my wife Helen. If it were not for her, I would still be a frustrated computer programmer. Instead, I am now not only a Mathematician, but the father of two wonderful girls. To my daughters Rebecca and Connie, thank you for many hours of fun and delight.

I am forever indebted to my parents, Elizabeth and Philip. Their decision to migrate to Australia from Myanmar (formerly Burma), has given me and my siblings opportunities that would never have been possible otherwise. They have instilled in me the respect for good education and the drive for personal excellence.

I would also like to thank my fellow graduate students for their companionship over the past four years. The many jokes shared in the IAM lab and the lunch time card games have helped me keep sane. Special thanks go to Jennifer Enns, Pete Newbury and Jeff Orchard who helped me print out my thesis remotely while I was starting my post-doctoral position in Australia.

Funding for much of my stay in Canada was provided by the University Graduate Fellowship (U.B.C.), the Teaching and Research Assistantships and by my parents.

This thesis was typeset using LaTeX.

Chapter 1

Introduction

Examples of differential equations abound in all areas of science. They arise whenever one attempts to model any process (abstract or physical) that depends on continuously varying quantities and the rates at which they change. Once a differential equation has been obtained that adequately models the problem at hand, the question of existence and uniqueness of solutions for a given set of data (initial or boundary) must then be addressed. One may then be interested in obtaining numerical, asymptotic or exact analytical solutions.

There are many branches of mathematics which deal with the above mentioned problems. Broadly speaking, they come under the general heading of analysis. However, the most general techniques for finding exact solutions of differential equations come from the branch of mathematics called *Lie group analysis of differential equations* and its generalizations. This subject lies at the crossroads of analysis and differential geometry, and was originated by its founder, Sophus Lie. The research presented here lies in this field of mathematics. We first give a general overview of the historical and conceptual background to this thesis.

1.1 Brief Historical Background

Marius Sophus Lie was born in Nordfjordeid, a town in the western part of Norway, on December 17, 1842. His school teacher [6] was Ludvig Sylow, who inspired him on the works of N.H. Abel and E. Galois. Abel showed that it was impossible to solve polynomial equations of order greater than four by radical expressions. Galois subsequently gave an elegant theory for this by considering symmetries of these equations. Lie and Sylow collaborated on a careful editing of Abel's completed works. Lie had a close friendship with F. Klein that led to a long fruitful collaboration. Klein, who published his Erlangen program in 1872 (see [33, 27]), developed the thesis that geometry was the study of invariants of group actions on geometric objects. Lie's starting point had been in modern geometry, after coming across the works of J.V. Poncelet and J. Plücker, and soon turned his attention to the geometry of differential equations with all his genius.

Lie formally defined and initiated the mathematical study of continuous groups of transformations, now called Lie groups. He showed how these led to a symmetry theory of differential equations, in the same spirit as the Galois theory for polynomial equations. Of the many achievements of Lie, of particular relevance to us is his algorithm on the explicit determination of the continuous group of transformations (the admitted group) that leave invariant the solution space of a given differential equation. With this algorithm, he was able to classify all ordinary differential equations with respect to the admitted group, and to develop an integration theory for such equations:

> In a short communication to the Scientific Society of Göttingen (3 December 1874), I gave, among other things, a listing of all continuous transformation groups in two variables x, y, and especially emphasized that this might be made the basis of a classification and rational integration theory of all differential equations $f(x, y, y', \cdots, y^{(m)}) = 0$, admitting a continuous group of transformations. The great program sketched there I have subsequently carried out in detail. (S. Lie [41], p.187)

> I noticed that the majority of ordinary differential equations which were integrable by the old methods were left invariant under certain transformations, and that these integration methods consisted in using that property. Once I had thus represented many old integration methods from a common view-point, I set myself the natural problem: To develop a general theory of integration for all ordinary differential equations admitting finite or infinitesimal transformations. (S. Lie [42], p. iv)

Sophus Lie died on Feb 18, 1899 and left a profound legacy. His work has greatly influenced the development of mathematics to the present day. His Lie groups and Lie algebras pervade much of mathematics and physics. For example, the modern theory of elementary particles in physics would be significantly different without his work.

The modern theory of Lie groups is radically different from that originally formulated by Lie. Lie was interested in groups of transformations that act on the solution space of a differential equation. In general, these are *local* Lie groups, with transformations being defined if they are sufficiently close to the identity (otherwise graphs of functions may not be mapped to other graphs of functions). Besides E. Noether, who proved a remarkable theorem [46] showing the one-to-one correspondence between symmetries of a variational integral and the conservation laws of the associated Euler-Lagrange equations, the application of Lie groups to differential equations was scarcely further developed for almost fifty years. During that time, Lie group theory underwent a radical abstract reformulation at the hands of E. Cartan and those who followed him. Interest switched from local Lie groups to global Lie groups. Present day interests focus on special classes of global Lie groups (semi-simple, solvable, etc.) and on representation theory, where the group action on the underlying space is linear. This is in direct contrast to the local Lie groups studied by Lie which were never so elegant.

A major revival of interest in Lie's original application of Lie groups to differential equations was sparked off by the works of Ovsiannikov in the former USSR in the late 1950's and 1960's [49] and of Bluman in the West in the late 1960's and 1970's [11, 12]. In the 1980's, the advent of symbolic manipulation computer programs (see the review paper [28]), which take away the often tedious computations associated with Lie's algorithm, has also helped in drawing researchers into this area. The basic theory can be found in Ovsiannikov [50], Olver [47], Bluman and Kumei [15], and in many other texts that have followed since. Besides applications of the theory to particular differential equations of interest, research in this area focusses on extending the scope of the theory. For example, in this dissertation, we develop a new symmetry algorithm which extends Lie's algorithm for the calculation of symmetries of general systems of PDEs.

Using this new algorithm, we discover large classes of higher order scalar and systems of partial differential equations admitting potential symmetries, which are new classes of symmetries.

For a more detailed account of Lie the mathematician, see Baas [6], Ibragimov [30], and the references therein.

1.2 Symmetries of Differential Equations

A symmetry of a differential equation (DE) is a transformation mapping any solution to another solution of the DE. Given a solution of the DE, knowledge of an admitted symmetry then leads to the generation of another solution. This is one of the most basic applications of symmetry analysis of DEs and relies on the symmetry being explicitly known. Even if one starts with a simple solution, which can usually be found by inspection, one can obtain a highly nontrivial solution in this way. Continuous symmetries such as translations, scalings and rotations can often be found by inspection. Even some discrete symmetries may be stumbled upon. However, most symmetries are not so easily found and the fundamental problem is one of algorithmic determination of symmetries. We shall restrict ourselves to finding continuous symmetries. The task of finding discrete symmetries is still very much an open subject (see Reid et al [55]).

Symmetry is defined in terms of transformations on the solution space of a DE. However, explicit knowledge of a symmetry in terms of local coordinates also defines how derivatives of solutions are transformed. Consequently, one can concentrate on how the DE itself is transformed instead of how its solution space is transformed. This is very useful, since the solution space is not usually known a priori. If one naïvely applies an arbitrary transformation on a DE, the symmetry conditions turn out to be an overdetermined system of *nonlinear* DEs for the transformation, which cannot be solved in general. To proceed, Lie considered a special class of transformations called a one-parameter group of point transformations or, more generally, contact transformations. These transformations are essentially the flows associated with vector fields defined in the space of independent and dependent variables. These vector fields and their components are called infinitesimal generators and infinitesimals of the point transformations, respectively. The fundamental insight due to Lie was that the symmetry conditions for these one-parameter groups of point transformations turn out to be *linear* in the infinitesimals of the transformations. Being linear, these infinitesimal determining equations are much easier to analyse and, in a large number of cases, have been solved explicitly to yield the admitted group of point (or contact) symmetries. The derivation of the infinitesimal determining equations of any given DE is called *Lie's algorithm.* Henceforth, we shall refer to 'one-parameter groups of point symmetries' as simply 'point symmetries'.

The infinitesimal generators of point symmetries depend only on independent and dependent variables. Lie also considered contact symmetries, where the infinitesimal generators also depend on first order derivatives of dependent variables. Since the time of Lie, the symmetry theory of DEs has been extended and applied in many ways. Applications now include:

- Construction of new solutions from old ones.
- Integration of ordinary differential equations (ODEs).
- Invariant (similarity) reductions of partial differential equations (PDEs).
- Solving boundary value problems (BVPs).
- Linearization of PDEs.
- Generalized symmetries.
- Construction of conservation laws (Noether's theorem).
- Equivalence transformations.
- Nonclassical solutions.

The theory behind Lie's algorithm is given in §2. For now, let us consider two illustrative examples.

Example 1.2.1 *Elementary Example of Symmetry*

Consider the simple second order ODE

$$\frac{d^2y}{dx^2} = 0, \tag{1.1}$$

which has solutions consisting of all non-vertical lines in the plane $\mathbb{R}^2$, given by

$$y = mx + c, \qquad m, c \in \mathbb{R}.$$

Examples of symmetries of the ODE are translations in y,

$$T_\varepsilon(x, y) = (x, y + \varepsilon),$$

and rotations about the origin,

$$R_\varepsilon \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \cos\varepsilon & -\sin\varepsilon \\ \sin\varepsilon & \cos\varepsilon \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix},$$

where $\varepsilon \in \mathbb{R}$. Clearly such translations and rotations map any straight line in the plane to another straight line and hence are symmetries of the ODE.

Notice that different values of ε define different transformations T_ε. Moreover, this family of transformations has a group structure: The identity element is T_0, the binary operation is composition ($T_\delta \circ T_\varepsilon = T_{\delta+\varepsilon}$) which is closed and associative, and to each transformation T_ε there corresponds its inverse $T_{-\varepsilon}$. The same can be said of R_ε, except that in this case, not all values of ε are permitted if R_ε is to be a symmetry of the ODE. For instance $R_{\pi/2}$ rotates a horizontal line to a vertical line which is no longer a solution of the ODE. However, if ε is sufficiently close to zero, then R_ε is a symmetry of the ODE. In general, symmetry transformations may only be defined *locally* about the identity transformation (corresponding to $\varepsilon = 0$). Both T_ε and R_ε are examples of local Lie groups of point transformations (T_ε is also a *global* Lie group of point transformations). They correspond to the flows of the infinitesimal generators (vector fields) $\mathbf{X}$ and $\mathbf{Y}$ respectively, given by

$$\mathbf{X} = \partial_x, \qquad \mathbf{Y} = -y\,\partial_x + x\,\partial_y.$$

So far, we have concentrated on how transformations act explicitly on solutions of the ODE.

This is made possible since the ODE was simple enough so that its solution space is explicitly known. In general, this is not true and one has to deal with the equations themselves. For any ε, the translations $(\widetilde{x},\widetilde{y}) = T_\varepsilon(x,y)$ transform the ODE to the new ODE, $\partial^2\widetilde{y}/\partial\widetilde{x}^2 = 0$, which clearly has the same solutions as the original ODE. As for the rotations $(\widetilde{x},\widetilde{y}) = R_\varepsilon(x,y)$, the original ODE is transformed to the new ODE

$$\Delta^3 \frac{d^2\widetilde{y}}{d\widetilde{x}^2} = 0, \qquad \Delta = \left(\cos\varepsilon - \sin\varepsilon \frac{dy}{dx}\right),$$

where dy/dx can be expressed in terms of $\widetilde{x},\widetilde{y}$ and $d\widetilde{y}/d\widetilde{x}$. So long as $\Delta \neq 0$, this ODE has the same solution set as the original ODE. The nonvanishing of Δ essentially restricts ε to be sufficiently close to the identity to ensure that straight lines do not become vertical. For example, rotation of the line $y = 0$, through an angle of $\frac{\pi}{2}$ results in a vertical line and this is reflected by the fact that Δ vanishes for the line $y = 0$ when $\varepsilon = \frac{\pi}{2}$.

Lastly in this example, the general solution of the ODE can be obtained from rotations and translations of any one solution: Any non-vertical line can be reached by rotating any given non-vertical line line to achieve the right slope, followed by vertical translations to achieve the right y-intercept. In general, for the case of partial differential equations (PDEs) where the number of independent variables is greater than one, one cannot hope to generate the general solution from symmetries. However, the principle of mapping solutions to solutions still holds and one can hope to generate nontrivial solutions by applying symmetries to simple solutions. □

Example 1.2.2 *Mapping Solutions to Solutions*

Consider the partial differential equation

$$u_t = u_{xx},$$

known as the heat equation. By inspection, the heat equation has the simple solution $u = 1$. Applying Lie's algorithm leads to the infinitesimal generator

$$\mathbf{X} = xt\,\partial_x + t^2\,\partial_t - (\tfrac{1}{4}x^2 + \tfrac{1}{2}tu)\,\partial_u, \tag{1.2}$$

whose flow corresponds to the one-parameter Lie group of transformations

$$\widetilde{x} = \frac{x}{1-\varepsilon t}, \qquad \widetilde{t} = \frac{t}{1-\varepsilon t}, \qquad \widetilde{u} = u\sqrt{1-\varepsilon t}\exp\left[\frac{-\varepsilon x^2}{4(1-\varepsilon t)}\right]. \tag{1.3}$$

This transformation can be used to generate a new solution from the simple solution $u = 1$ as follows. The graph of this simple solution is given by

$$\{(x, t, 1) : x, t \in \mathbb{R}\}.$$

Now under the given symmetry transformation, any point $(x, t, 1)$ on this graph is transformed to a new point $(\widetilde{x}, \widetilde{t}, \widetilde{u})$, where $\widetilde{u}$ is the value of the new function $\widetilde{f}$ over the point $(\widetilde{x}, \widetilde{t})$, given by (1.3). This new function, obtained by expressing (x, t) in terms of $(\widetilde{x}, \widetilde{t})$ in (1.3c), is given by

$$\widetilde{u} = \widetilde{f}(\widetilde{x}, \widetilde{t}) = \frac{1}{\sqrt{1+\varepsilon\widetilde{t}}}\exp\left[\frac{\varepsilon\widetilde{x}^2}{4(1+\varepsilon\widetilde{t})}\right].$$

Hence, we have generated a highly nontrivial solution from a very simple solution, through the application of a symmetry.[1] □

Further examples of the applications of symmetry analysis can be found in the standard texts mentioned previously. However, we note in particular that there has been a lot of recent interest in the nonclassical method of Bluman and Cole [11] where the definition of invariant solutions of a DE is further extended. In the classical approach, one looks for solutions that are invariant under some point symmetry admitted by the DE. However, in the nonclassical approach, a special class of solutions of the DE are sought which admits a point symmetry that is *not admitted* by the DE as a whole. See Clarkson and Mansfield [24] for an algorithmic implementation of the nonclassical method and Bluman and Shtelen [19] for further extensions of the method. Other exact solution methods include the work of Ovsiannikov [50] on partially invariant solutions and Olver [48] on differential constraints. For work on the equivalence transformations, see Lisle [43].

[1] As an undergraduate, I did not like differential equations very much. However, as a Master's student searching for something to work on [25], I came across this very simple, but elegant application of symmetry analysis and was instantly won over to the study of Lie theory of DEs.

1.3 Potential Systems

As already mentioned, the symmetries considered by Lie were point symmetries where the infinitesimals depend on independent variables x and dependent variables u, and contact symmetries where the infinitesimals can also depend on first order derivatives of u. Lie theory of DEs is further extended by the consideration of generalized symmetries, where the infinitesimal generators can also depend on the derivatives of dependent variables up to some finite order. Point, contact and generalized symmetries are called *local symmetries*, since their infinitesimal generators depend only on local properties of solutions. In particular, at any point x, the infinitesimal generators depend only on x and the values of u and its derivatives at that point.

One can further enlarge the class of known symmetries of DEs by considering *nonlocal symmetries*, which are characterized by infinitesimal generators that are not of local type. For example, if the infinitesimal generator depends on integrals of dependent variables, then the corresponding symmetry would be nonlocal. In principle, a DE can admit many nonlocal symmetries, but the fundamental issue is in their algorithmic determination. Special and/or heuristic techniques have been employed by Akhatov et. al. [1], Konopelchenko and Mokhnachev [34], Kumei [37], Kapcov [32], Pukhnachev [52], and Krasil'shchik and Vinogradov [35, 36] to obtain nonlocal symmetries of DEs. In particular, the approach of Krasil'shchik and Vinogradov is restricted to PDEs with two independent variables. In this thesis, we will use the potential systems approach of Bluman, Kumei and Reid [16] which is a systematic method for finding nonlocal symmetries of PDEs with two or more independent variables. Nonlocal symmetries of ODEs require a different approach (see Bluman and Reid [17]) and are not considered in this thesis.

A simple minded way to find nonlocal symmetries of a given PDE is to apply Lie's algorithm with infinitesimal generators of the form

$$\mathbf{X} = \xi(x, u, u_{-1})\partial_x + \eta(x, u, u_{-1})\partial_u, \qquad u_{-1} = \int u\, dx.$$

If unsuccessful, then what other forms of infinitesimal generators should one consider next?

Clearly such an approach is *ad hoc* and a systematic procedure is needed.

In the potential systems approach, one uses a conservation law of the PDE $R\{u\}$ to form an associated potential system $S\{u,v\}$ involving the introduction of potentials v. Here, the potential variable v, which is algorithmically determined, replaces the formal variable u_{-1} in the above ad-hoc approach. By their very construction, potentials are nonlocal with respect to the dependent variable u of the original PDE $R\{u\}$. Since the solutions of $R\{u\}$ are nonlocally embedded in the solutions of $S\{u,v\}$, studying $S\{u,v\}$ may lead to new nonlocal information for $R\{u\}$ and vice versa. Also, this embedding of solutions is not one-to-one so that invertible mappings in (x,u,v)-space can lead to noninvertible mappings in (x,u)-space and vice versa. In this dissertation we only use the potential system $S\{u,v\}$ for finding nonlocal symmetries of $R\{u\}$. The use of potential systems to deduce other types of nonlocal information for $R\{u\}$ can be found in [3, 4, 18, 20].

Nonlocal symmetries of $R\{u\}$ may be found as point symmetries of $S\{u,v\}$ which do not project onto any point symmetry of $R\{u\}$. Nonlocal symmetries that arise in this way are called *potential symmetries*. Being realized as point symmetries of $S\{u,v\}$, Lie's algorithm[2] can be used to calculate these potential symmetries and, once found, all the applications of point symmetries outlined in the previous section are available. In particular, potential symmetries can lead to new information for R that is not obtainable via local symmetries of R. Applications of potential symmetries include:

- Noninvertible mappings of known solutions to new solutions.
- New invariant solutions of R.
- Exact solutions of new boundary value problems for R.
- Linearizations of R through *noninvertible* transformations.
- Nonlocal mappings between PDEs.
- Nonlocal conservation laws of PDEs.

[2]In §1.4, we will have more to say on the use of Lie's algorithm to find point symmetries of potential systems.

Let us now discuss, through illustrative examples, the construction of potential systems $S\{u,v\}$, how their point symmetries can lead to potential symmetries of the original PDE $R\{u\}$, and how potential symmetries can lead to new nonlocal information for $R\{u\}$.

Construction of Potential Systems

The idea of using a conservation law of a PDE to introduce potentials is not new. Applications have concentrated mainly on reducing the original system of PDEs to an equivalent system involving only the potentials. The main advantage of this is that the resulting system contains fewer equations and hence is easier to analyse.

Example 1.3.1 *Potentials in Physics*

In physics, the equations defining electromagnetic radiation are governed by Maxwell's equations

$$\frac{\partial \mathbf{E}}{\partial t} = \nabla \times \mathbf{B}, \qquad \frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E}, \qquad \nabla \cdot \mathbf{E} = 0, \qquad \nabla \cdot \mathbf{B} = 0,$$

where $\mathbf{E} \in \mathbb{R}^3$ is the electric field and $\mathbf{B} \in \mathbb{R}^3$ is the magnetic field. One can introduce a vector potential $\mathbf{A}$ such that $\mathbf{B} = \nabla \times \mathbf{A}$ is automatically divergence-free, i.e., $\nabla \cdot \mathbf{B} = 0$. Then obviously, $\nabla \times (\partial \mathbf{A}/\partial t + \mathbf{E}) = 0$. Consequently, one can introduce a scalar potential ϕ such that $\partial \mathbf{A}/\partial t + \mathbf{E} = \nabla \phi$. Maxwell's equations are then reduced to the following equivalent system:

$$\frac{\partial^2 \mathbf{A}}{\partial t^2} + \nabla \times (\nabla \times \mathbf{A}) = \nabla \frac{\partial \phi}{\partial t}, \qquad \nabla \cdot \frac{\partial \mathbf{A}}{\partial t} = \Delta \phi,$$

which involves only the potentials $\mathbf{A}$ and ϕ. $\square$

In the potential systems approach one does not seek to eliminate the original dependent variables to form a reduced system involving only the potentials, as was the case in the above example of Maxwell's equations. Rather, one replaces the given conservation law (divergence-free expression) with the equations defining the potentials to form a *potential system*, as shall be illustrated in the following two examples. The reduced equations can yield useful information for the original PDE [10]. However, from the perspective of finding nonlocal symmetries, the potential system allows these symmetries to be *explicitly* realized. We shall point this out when we come to discuss potential symmetries in §4.1.1.

Example 1.3.2 *Case of two independent variables*

In the case of two independent variables, the prototypical example is the nonlinear diffusion equation

$$\frac{\partial}{\partial t}(-u) + \frac{\partial}{\partial x}(K(u)u_x) = 0, \tag{1.4}$$

with corresponding potential system

$$\begin{aligned} v_x &= u, \\ v_t &= K(u)u_x, \end{aligned} \tag{1.5}$$

where u is the concentration, $K(u)$ is the concentration-dependent diffusivity and v is a scalar potential. Notice that taking the sum of the t-derivative of (1.5a) and the x-derivative of (1.5b) leads to the original PDE (1.4). We say that the original PDE is a *differential consequence* or an *integrability condition* of the associated potential system. □

Example 1.3.3 *Case of three independent variables*

In the case of three independent variables, the prototypical example is the nonlinear wave equation

$$\frac{\partial}{\partial t}(-u_t) + \frac{\partial}{\partial x}(C_1(u)u_x) + \frac{\partial}{\partial y}(C_2(u)u_y) = 0, \tag{1.6}$$

with corresponding potential system

$$\begin{aligned} -u_t - v_{3,x} + v_{2,y} &= 0, \\ C_1(u)u_x - v_{1,y} + v_{3,t} &= 0, \\ C_2(u)u_y - v_{2,t} + v_{1,x} &= 0, \end{aligned} \tag{1.7}$$

where $C_1(u)$ and $C_2(u)$ are the wave speeds, $v = (v_1, v_2, v_3)$ is a vector potential, and $v_{i,x_j} = \partial v_i/\partial x_j$ ($x_j = t$, x or y). As in the previous case, the original wave equation is a differential consequence of the associated potential system, as can be seen by taking the sum of the t-derivative of (1.7a), the x-derivative of (1.7b) and the y-derivative of (1.7c). □

The precise construction of potential systems is described in §4.1. For now, it suffices to say that the procedure is algorithmic: In [26] Haager et al. have developed a software package which automates the construction process.

Potential Symmetries and Applications

Point symmetries of $R\{u\}$ are of the form

$$\mathbf{X}^R = \xi^R(x,u)\,\partial_x + \eta^R(x,u)\,\partial_u,$$

whereas point symmetries of $S\{u,v\}$ are of the form

$$\mathbf{X}^S = \xi^S(x,u,v)\,\partial_x + \eta^S(x,u,v)\,\partial_u + \rho^S(x,u,v)\,\partial_v.$$

It turns out that if $S\{u,v\}$ admits $\mathbf{X}^S$, then $R\{u\}$ must admit

$$\widehat{\mathbf{X}}^S = \xi^S(x,u,v)\,\partial_x + \eta^S(x,u,v)\,\partial_u.$$

By comparing $\mathbf{X}^R$ and $\widehat{\mathbf{X}}^S$, one can see that a new symmetry has been found for $R\{u\}$ if (ξ^S, η^S) depends on v. These new symmetries, called potential symmetries, are nonlocal symmetries of $R\{u\}$ since v is nonlocal with respect to u.

Example 1.3.4 *Potential Symmetry*

Let $R\{u\}$ be the nonlinear diffusion equation (1.4) with associated potential system $S\{u,v\}$, given by (1.5). If $K(u) = \exp(a \arctan u)/(1+u^2)$, where a is a constant, then $S\{u,v\}$ admits the infinitesimal generator [16]

$$\mathbf{X}^S = v\partial_x + at\partial_t - (1+u^2)\partial_u - x\partial_v.$$

Since the infinitesimal corresponding to x depends on v, we have found a potential symmetry of $R\{u\}$. In particular, the flow of $\mathbf{X}^S$ corresponds to the symmetry transformation of $S\{u,v\}$ given by

$$\begin{aligned}
\tilde{x} &= x\cos\varepsilon + v\sin\varepsilon, \\
\tilde{t} &= e^{a\varepsilon}t, \\
\tilde{u} &= \tfrac{u\cos\varepsilon - \sin\varepsilon}{\cos\varepsilon + u\sin\varepsilon}, \\
\tilde{v} &= -x\sin\varepsilon + v\cos\varepsilon,
\end{aligned}$$

which is defined for all $\varepsilon \in \mathbb{R}$ sufficiently close to zero. This transformation induces a symmetry transformation of $R\{u\}$ obtained by projection onto (x, u)-space:

$$\begin{aligned}
\tilde{x} &= x\cos\varepsilon + (\textstyle\int u dx)\sin\varepsilon, \\
\tilde{t} &= e^{a\varepsilon}t, \\
\tilde{u} &= \tfrac{u\cos\varepsilon - \sin\varepsilon}{\cos\varepsilon + u\sin\varepsilon}.
\end{aligned}$$

This induced symmetry transformation is clearly nonlocal. By projecting $\mathbf{X}^S$ to (x, u)-space, one obtains the nonlocal infinitesimal generator

$$\hat{\mathbf{X}}^S = (\int u dx)\partial_x + at\partial_t - (1 + u^2)\partial_u,$$

admitted by $R\{u\}$. □

The following example further illustrates why potential systems and potential symmetries are worthy of study.

Example 1.3.5 *Noninvertible Linearizations*

Consider the nonlinear system of PDEs $R\{u^1, u^2\}$, given by

$$u^1_x - u^2_t = 0,$$

$$u^2_t - u^1 u^2 - u^1 - u^2 = 0,$$

which describes fluid flow through a reacting medium. These equations, also known as the Thomas equations, were studied by Thomas [64] (see also Whitham [71]). When applying the linearization algorithms [38, 15], one can show that there exists no invertible transformation which linearizes R. Evidently, the point symmetries of R are not sufficiently rich enough.

However, a noninvertible linearization of R can be found through a potential system of R:

Using the first equation in R which is a conservation law, one can form the associated system $S\{u^1, u^2, v\}$, given by

$$v_t = u^1,$$

$$v_x = u^2,$$

$$u_t^2 - u^1 u^2 - u^1 - u^2 = 0.$$

One can show that S admits an infinite-parameter family of point transformations with infinitesimal generator

$$\mathbf{X} = e^v \left\{ (\psi_t + u^1\psi)\partial_{u^1} + (\psi_x + u^2\psi)\partial_{u^2} + \psi\partial_v \right\},$$

where $\psi(x,t)$ is an arbitrary function satisfying the linear PDE

$$\psi_{xt} - \psi_x - \psi_t = 0.$$

Consequently, the linearization algorithm leads to the following *invertible* mapping

$$\begin{aligned} z_1 &= x, \qquad & w^1 &= -e^{-v}, \qquad & w^3 &= e^{-v}u^1, \\ z_2 &= t, \qquad & w^2 &= e^{-v}u^2, \end{aligned} \tag{1.8}$$

which transforms S to the linear system

$$w^1_{z_1} = w^2, \qquad w^1_{z_2} = w^3, \qquad w^2_{z_2} = w^2 + w^3.$$

Moreover, projecting the transformations (1.8) to (x,u)-space, one obtains the *noninvertible* transformation

$$\begin{aligned} z_1 &= x, \qquad & w^1 &= -e^{-\int u^1 dt}, \\ z_2 &= t, \qquad & w^2 &= u^2 e^{-\int u^1 dt}, \end{aligned}$$

which maps R to the linear system

$$w^1_{z_1} = w^2, \qquad w^2_{z_2} = w^2 + w^1_{z_2}.$$

Hence, we have found a noninvertible linearization of R through potential symmetries of R. $\square$

1.4 A New Symmetry Algorithm For Systems of PDEs

Lie's algorithm successfully leads to the point symmetries admitted by scalar PDEs and systems of PDEs of Cauchy-Kovalevskaya type. However, we have encountered problems when applying this algorithm to more general systems of PDEs, such as potential systems. As such, we have had to derive a new symmetry algorithm to overcome these problems. To do this requires us to synthesize ideas from the Formal Theory of Integrability and the existing Lie Theory of DEs. We now sketch out the main points.

Recall that a symmetry of an n-th order system R of PDEs is a transformation mapping any solution to another solution of R, i.e., a transformation that leaves the solution space of R invariant. Since the solution space of R is not known explicitly in general, Lie's algorithm deals solely with the equations of R. The equations of R define a locus of points (subvariety) in the space of independent variables, dependent variables and all derivatives of dependent variables up to order n (n-jet space or n-th extended space). Since solutions of R lie in the locus of R, one arrives at the symmetries of the solution space of R by finding the symmetries of the locus of R. There are two important caveats to this.

The first caveat is that symmetries of the locus of R are only guaranteed to map points in the locus to other points in the locus: They may not map functions to functions. As such, while they are guaranteed to map solutions of R to other points in the locus of R, they may not map solutions to other solutions of R. One ensures that solutions of R are mapped to other solutions by seeking symmetries that belong to the class of extended point transformations.

The second caveat is that one may not obtain the full set of symmetries of R unless certain differential consequences of the equations of R are used. Appending differential consequences to the equations of R leads to a new system $\widetilde{R}$ which has the same solution space as R, but a different locus. As such, the symmetries of R and $\widetilde{R}$ are the same, but the symmetries of their respective loci may be different. It is true that any such system $\widetilde{R}$ can be used to find symmetries of R. However, in general the full set of point symmetries of R may not be uncovered. In short, to obtain all the point symmetries of R, one must apply Lie's algorithm to a *locally solvable* system $\widetilde{R}$, i.e., $\widetilde{R}$ must be such that given any point in its locus there must pass a solution of R. If $\widetilde{R}$ is not locally solvable, symmetries of its locus are required to leave invariant, not only the solutions of $\widetilde{R}$, but also regions of the locus through which there are no solutions. This leads to stronger conditions than necessary on the symmetry transformations.

For analytic systems[3] of PDEs, an n-th order system of PDEs is locally solvable if the system contains all its differential consequences up to order n. It turns out that scalar PDEs and systems of PDEs of Cauchy-Kovalevskaya type are locally solvable and, hence, Lie's algorithm successfully leads to all the point symmetries admitted by such PDEs. The situation for more general systems of PDEs is not so simple, as we now illustrate in the following two examples.

Example 1.4.1 Let R be the system of second order PDEs given by

$$\begin{aligned} u_{xx} &= v, \\ u_t &= v. \end{aligned} \tag{1.9}$$

By differentiating (1.9b), we obtain the second order differential consequences

$$\begin{aligned} u_{xt} &= v_t, \\ u_{tt} &= v_x. \end{aligned} \tag{1.10}$$

However, we have not yet achieved local solvability. There is one further second order differential consequence obtained by equating mixed partial derivatives $u_{txx} = u_{xxt}$ to obtain

$$v_{xx} = v_t. \tag{1.11}$$

The system $\widetilde{R}$ given by (1.9), (1.10) and (1.11) is locally solvable since there are no further differential consequences up to order two. Hence, to find all the point symmetries of R, one must apply Lie's algorithm to the system $\widetilde{R}$. $\square$

Here is an example of a system of PDEs where local solvability cannot be achieved for any finite order.

[3] By staying in the analytic regime, we avoid certain smooth DEs for which there exist no solutions (see Lewy's counter example [39]).

Example 1.4.2 Consider the third order scalar PDE

$$-D_tF(x,t,u^{(2)})+D_xG(x,t,u^{(2)})=0, \tag{1.12}$$

which has the associated potential system S, given by

$$\begin{aligned} v_x &= F, \\ v_t &= G, \end{aligned} \tag{1.13}$$

where F and G are fixed, but not explicitly given. When looking for all the point symmetries of S, which is of order two, one must first make the system locally solvable by uncovering all its differential consequences up to order two. Such differential consequences can occur: For example, when $(F,G)=(u_{xx}+u,u_{xx})$, S has the second order differential consequence $v_{xx}=v_{tx}+u_x$. Unfortunately, since F and G are not explicitly given (such a situation will arise in §5.1 when we study symmetries of potential systems), one cannot deduce all differential consequences of S up to order two. The best one can do is to differentiate the equations of S to obtain

$$\begin{aligned} v_{xx} &= D_xF, \\ v_{xt} &= D_tF, \\ v_{tt} &= D_tG, \end{aligned} \tag{1.14}$$

and the original scalar PDE (1.12) which arises through the compatibility condition $v_{xt}=v_{tx}$. The system $\widehat{S}$, given by (1.12), (1.13) and (1.14), is of third order and to achieve local solvability one must determine all differential consequences of $\widehat{S}$ up to order three. But this is not possible for the same reason that S cannot be made locally solvable. This problem persists after any number of differentiations of the equations of S. Consequently S cannot be made locally solvable at any finite order. As will be shown in §3.2, even the infinite system $\widetilde{S}$ obtained by differentiating the equations of the system (1.12) and (1.13) to all orders is not locally solvable. However, it turns out that such an infinite system satisfies the weaker property of *analytic local solvability* and this will be sufficient for our purposes. Even so, the application of Lie's algorithm to such an infinite system $\widetilde{S}$ is clearly not feasible. □

The above example illustrates the problems that one may encounter when applying Lie's algorithm to find symmetries of systems of PDEs. In §3.3, we show how one can relax the local solvability requirement. Essentially, we show that the symmetry conditions for any locally solvable system of PDEs can be reduced to an equivalent set of conditions involving significantly fewer equations. In the new symmetry algorithm that results, only the equations of the given system of PDEs are required at the first step. At a later step in the algorithm (the substitution step), only a finite number of differential consequences of the system are required.

For scalar PDEs and systems of PDEs of Cauchy-Kovalevskaya type, our symmetry algorithm is essentially the same as Lie's algorithm. For more general systems of PDEs, our algorithm is much more efficient: When finding symmetries of the system (1.13), only a finite number of its differential consequences are required, whereas Lie's algorithm requires the infinite system $\widetilde{S}$ described in Example 1.4.2. Even when finding symmetries of the system (1.10), Lie's algorithm requires all the equations of the locally solvable system (1.9), (1.10) and (1.11) from the very beginning, whereas our symmetry algorithm does not need to use the differential consequences (1.10) and (1.11) until the less computationally intensive substitution step.

1.5 Outline of Thesis

The following is assumed throughout the thesis:

Blanket locality and analyticity assumption: *We always assume that all differential equations, their solutions and mappings are local analytic functions of their arguments. For simplicity of exposition we will discuss local properties using local coordinates.*

This blanket analyticity assumption may be relaxed in some cases. Where it is crucial to have analyticity, we shall explicitly state it.

We anticipate that this dissertation will be of interest to at least three different audiences. One such audience will be interested mainly in the potential systems approach for finding nonlocal symmetries of PDEs. Another audience will be interested in our extension of Lie's algorithm and its theoretical basis. The third audience will be researchers in the field of symbolic computation who will be interested in the algorithms of this dissertation as well as their computer implementations. To help each reader access the specific information they need, we feel it essential to provide a 'road map' designed for each of these three audiences. We will do this after giving a brief description of each chapter. A more detailed description of the contents of each chapter is provided at the beginning of each chapter.

The standard theory behind Lie's algorithm is presented in §2. The symmetry transformations we consider belong to the class of transformations associated with flows of vector fields. We seek such transformations which leave invariant the solution space of the given system R of PDEs. As discussed in §1.4, one arrives at the symmetries of R through the symmetries of the locus of R, by restricting to the class of extended point transformations and by requiring that R be locally solvable. As such, the theory of extended point transformations and their corresponding extended vector fields is provided. (The issue of local solvability will be tackled in §3.) Symmetries of the locus are given by vector fields which are tangent to the locus. By restricting to extended vector fields the associated tangency conditions, called the infinitesimal symmetry conditions for R, lead to the required symmetries of R. The resulting algorithm is called Lie's algorithm. A discussion of the various other symmetry formulations, other algorithms and their shortcomings is provided. In particular, we point out that a commonly used and accepted algorithm for the calculation of symmetries of systems of PDEs appears to lack theoretical justification.

In §3, a new symmetry algorithm which extends Lie's algorithm to systems of PDEs is presented. In order to do this, there are two questions to be answered. Firstly, how does one achieve local solvability in general? Secondly, how can one find the point symmetries admitted by systems of PDEs, such as (1.13), whose (analytic) locally solvable form consists of an infinite number of equations? The Formal Theory of Integrability of Riquier-Janet answers the first question. Here we follow the more efficient approach of Reid. By a finite number of operations involving differentiations, back substitutions and applications of the Implicit Function Theorem, a *prolonged standard form* is constructed. We show that for any system of PDEs, it is the associated prolonged standard form for which Lie's algorithm applies.

To address the second question, we show how the infinitesimal symmetry conditions for a prolonged standard form can be reduced to an equivalent set of symmetry conditions involving significantly fewer equations. This leads to a new symmetry algorithm which is more efficient than Lie's algorithm. If the starting system of PDEs consists of a finite number of equations, then only a finite number of equations from the prolonged standard form is required in this new algorithm. This is true even if the prolonged standard form comprises of an infinite number of equations.

In §4, we present the mathematical framework of the potential systems approach for finding nonlocal symmetries of systems of PDEs. We provide details of how to find conservation laws, how to construct potential systems, how to delineate which of these systems are useful for finding potential symmetries, and how one can iterate this process to obtain higher generation useful potential systems. A complete potential systems analysis of the nonlinear diffusion equation is provided which shows how potential symmetries can arise. The construction of conservation laws is through linear combinations, involving coefficients called factors, of the equations of the system. We show how these factors can be used to derive necessary conditions for the linearization of systems of PDEs.

To date, all examples of potential symmetries involve scalar PDEs of order two and systems of PDEs of order one. In §5, we tackle the problem of finding higher order scalar and systems of PDEs admitting potential symmetries. Preliminary work on this problem appears in [51], where necessary conditions for such PDEs are claimed, but where no examples were given. It turns out that there is an error in their claims, since they neglect to form a locally solvable system before applying Lie's algorithm. We correct their claims using the new symmetry algorithm of §3. We then proceed to construct large classes of higher order scalar and systems of PDEs which admit potential symmetries.

Our results are summarized in §6, where possible future directions for this research are also discussed.

1.5.1 Specific Guide to Thesis

The reader, who is interested only in the potential systems approach and who is already familiar with Lie's algorithm, can start immediately at §4 and read on. However, in §5 we make extensive use of our new symmetry algorithm. Unless the reader is willing to read §2 and §3 for the complete theory, the reader is advised to do the following: Read §1.4 to get the basic concept of local solvability and the need of our new symmetry algorithm. Look at Theorem 3.3.3 which is the basis of our new symmetry algorithm, given by Algorithm 3.3.4, as well as the example that follows this theorem.

The reader should be familiar with all aspects of Algorithm 3.3.4 except for the use of the prolonged standard form in the substitution step (step 2). Just think of the prolonged standard form as providing all necessary substitutions that are derived from the equations of the given system. The algorithms for achieving a prolonged standard form are summarized in Appendix A. However, the reader should be able to follow the calculations of §5 without reading Appendix A. If the reader is still unconvinced for the need of the new symmetry algorithm, consider this. Any algorithm may lead to infinitesimal generators that can be verified, by other means, as symmetries of the given system of PDEs. But the point is, where is the guarantee that the algorithm used leads to the full set of the admitted infinitesimal generators? Through Theorem 3.3.3, we provide a sound theoretical basis for our algorithm.

The reader who is interested in our new symmetry algorithm and the theory behind it, should read §2 and §3. The proof of Theorem 3.3.3, which our new symmetry algorithm relies on, is of a technical nature and is given in Appendix C. For applications of our symmetry algorithm, see §5, where the symmetries of potential systems are investigated. Much of our effort here is spent on deriving the prolonged standard forms, which is an essential ingredient in the new symmetry algorithm, for the given potential systems. Read §1.3 to find out how potential systems are constructed and why they are important to study. We also direct the reader to Definition 4.0.1 and Theorem E.0.1, where the prolonged standard form has been used to define local symmetries and to state the Fréchet formulation for finding them, respectively.

The reader interested in algorithms and their computer implementations is directed to Lie's algorithm given by Algorithm 2.3.8, to a commonly used variant of Lie's algorithm given by Algorithm 2.3.9, and to our new algorithm given by Algorithm 3.3.4. Notice that the new algorithm does not require the given system to be in locally solvable form. Also, notice that the substitution step (step 2), which is required in each algorithm, has been made more explicit (and more easy to implement on a computer) in the new algorithm, through the use of the prolonged standard form. The algorithms for achieving prolonged standard form are given in Appendix A. If unfamiliar with the Formal Theory of Integrability of Riquier-Janet and the improved algorithms of Reid, then the reader is referred to §3.1. See Definition 4.0.1, Theorem E.0.1 and Theorem 4.2.5, where the prolonged standard form has been used to define local symmetries, to state the Fréchet formulation for finding them, and to state the adjoint theorem for finding factors leading to conservation laws. These particular formulations can lead to improved algorithms for finding local symmetries and factors. Lastly, we point out that a computer package [26] now exists which automates various aspects of the potential systems approach.

Chapter 2

Lie's Algorithm

In this chapter, we develop the standard theory behind Lie's algorithm which is used to determine all the point symmetries of a given PDE. In §2.1, we study transformation groups. The transformations that Lie considered are those represented by the flows of (analytic) infinitesimal generators (vector fields). Given an infinitesimal generator, the corresponding set of transformations, which satisfy the group axioms, is called a one-parameter Lie group of point transformations. A set of infinitesimal generators generates a multi-parameter Lie group if and only if they form a Lie algebra. In §2.1.4, the infinitesimal generators admitted by a system of algebraic equations are derived. One requires that the infinitesimal generators are tangent to the locus of the system and this leads to the algebraic infinitesimal symmetry conditions.

The n-jet (extended) space, i.e., the space of independent variables, dependent variables and all derivatives of the dependent variables up to order n, is described in §2.2. The locus of a system of DEs is the set of points in n-jet space which satisfies the equations of the system. As mentioned in §1.4, the symmetries of the locus lead to the symmetries of the system of DEs, provided that the system is locally solvable and the symmetry transformations map functions to functions. To ensure the latter, one restricts oneself to Lie groups of extended point transformations which are the flows of vector fields, called extended infinitesimal generators, defined in n-jet space.

In §2.3, we derive the symmetry conditions for a system of PDEs. As in the case of algebraic systems, one requires the extended infinitesimal generators to be tangent to the locus of the system. This leads to Lie's algorithm for determining the point symmetries admitted by the system of PDEs. A discussion of the various other forms of the commonly used symmetry conditions and algorithms together with their theoretical shortcomings is provided. The local solvability requirement of Lie's algorithm can lead to problems and, hence, an extension of Lie's algorithm to overcome these problems is needed.

2.1 Transformation Groups

Definition 2.1.1 Let $M = \mathbb{R}^p$, with coordinates $x = (x_1, x_2, \cdots, x_p)$. A *point transformation* of a space M is an (analytic) (C^∞) mapping $\widetilde{x} = \tau(x)$ such that τ is one-to-one and onto. The transformation corresponding to the inverse of τ is denoted by τ^{-1}.

More generally, one can allow M to be any differentiable manifold[1] with local coordinates x, and τ defined only locally so that the rotations R_ε of Example 1.2.1 are also considered as transformations. In this thesis we shall not always explicitly emphasize the local nature of our statements, but it will always be assumed. Also, we will be dealing almost exclusively with *one-parameter Lie groups* of transformations and as such the unqualified term 'transformation' will aways be taken to be a 'point transformation'. We will see that these transformations are essentially the flows of vector fields on M. Given such a characterization, we will be able to write down the symmetry conditions for algebraic systems of equations in terms of these vector fields. This will then pave the way for determining the infinitesimal symmetry conditions of DEs, given in the next section. Let us first review the more familiar concept of flows of vector fields.

2.1.1 Flows of Vector Fields

Definition 2.1.2 Let M be a differentiable manifold and $TM|_x$ be the space of tangent vectors to M at x. A *vector field* $\mathbf{X}$ on M assigns a tangent vector $\mathbf{X}|_x \in TM|_x$ to each point $x \in M$.

[1]The precise definition of an r-dimensional differentiable manifold M can be found in [69, 47]. It shall be sufficient for us to treat M as a space made up of subsets of $\mathbb{R}^r$ that are patched together in a suitable manner. Since we will always work locally, the global topology of a manifold will not affect us.

An *integral curve* of $\mathbf{X}$ through $x = x_0$ is a parametrized curve $x = \phi_{x_0}(\varepsilon)$ passing through x_0 and whose velocity $\dot{\phi}_{x_0}(\varepsilon)$ coincides with the vector field $\mathbf{X}$, at any point along the path:

$$\phi_{x_0}(0) = x_0, \qquad \dot{\phi}_{x_0}(\varepsilon) = \mathbf{X}|_{\phi_{x_0}(\varepsilon)}, \; \forall\, \varepsilon \in I, \tag{2.1}$$

where I is some open interval of $\mathbb{R}$ containing the origin[2]. Different integral curves through a given point can have different domains of definitions I and the unique one corresponding to the maximum domain of definition is called the *maximal integral* curve through the point.

Example 2.1.3 An example of an integral curve is the path traced out by a particle that is being carried along by a river. Here, M is the surface of the river and the vector field $\mathbf{X}$ at any point $x \in M$ is given by the surface velocity of the river at that point. Suppose that the flow is steady. Then these velocities, and hence $\mathbf{X}$, do not change with time. Let ε be the time, x_0 be the location of a particle at $\varepsilon = 0$ and $\phi_{x_0}(\varepsilon)$ be the path traced out by the particle over time. Then $\phi_{x_0}(\varepsilon)$, $\varepsilon \in I$, is an integral curve of $\mathbf{X}$ through $x = x_0$ for some open interval I containing the origin. Assuming that the river is infinitely long with no sinks or sources, the maximum integral curve of $\mathbf{X}$ through $x = x_0$ is the unique integral curve where $I = \mathbb{R}$. □

We denote the maximal integral curve of $\mathbf{X}$ passing through $x \in M$ by $\Psi(\varepsilon, x)$ and call Ψ the *flow* generated by $\mathbf{X}$. The flow has the properties:

$$\Psi(\delta, \Psi(\varepsilon, x)) = \Psi(\delta + \varepsilon, x), \qquad x \in M, \tag{2.2}$$

for all $\delta, \varepsilon \in I$ such that both sides are defined (one cannot flow past a sink),

$$\Psi(0, x) = x, \qquad \text{and} \qquad \frac{\partial}{\partial \varepsilon}\Psi(\varepsilon, x) = \mathbf{X}\Big|_{\Psi(\varepsilon,x)}. \tag{2.3}$$

Here is how flows of vector fields give rise to transformations on the underlying space. Let $\tau_\varepsilon(x) = \Psi(\varepsilon, x)$. Then for each fixed ε, $\tau_\varepsilon : M \to M$ defines a transformation on M. To see this, recall the analogy of a vector field as representing a steady state surface velocity of a river. To determine how τ_ε transforms any given $x \in M$, just imagine dropping a particle on the

[2]We further emphasize that (2.1b) is not only satisfied at the point $x = x_0$ (so that $\dot{\phi}_{x_0}(0) = \mathbf{X}|_{x_0}$), but also at all other points along the curve $x = \phi_{x_0}(\varepsilon)$, $\forall\, \varepsilon \in I$.

surface of the river at x and letting it flow for a time ε. The resulting location of the particle is $\tau_\varepsilon(x)$. Different values of ε lead to different transformations τ_ε and the set of all these is called a *one-parameter family* of transformations.

In terms of the local coordinates x, a vector field has the form

$$\mathbf{X}|_x = \xi^1(x)\,\frac{\partial}{\partial x_1}\bigg|_x + \xi^2(x)\,\frac{\partial}{\partial x_2}\bigg|_x + \cdots + \xi^p(x)\,\frac{\partial}{\partial x_p}\bigg|_x, \tag{2.4}$$

where $(\partial/\partial_{x_i})|_x$, $i = 1, \cdots, p$, forms a basis for TM at x. Henceforth, we shall drop the symbol $|_x$ and it should be clear, from the context, which point x or tangent space $TM|_x$ one is referring to. We shall also use ∂_{x_i} to denote ∂/∂_{x_i} and assume ξ are analytic functions of x so that we deal exclusively with analytic vector fields. Using (2.4), it is convenient to also express (2.1) in terms of local coordinates:

Theorem 2.1.4 *The one-parameter family of transformations $\tilde{x} = \tau_\varepsilon(x)$ associated with an analytic vector field $\mathbf{X}$, given by (2.4), is the unique solution to the autonomous system of ordinary differential equations (ODEs)*

$$\frac{\partial \tilde{x}_i}{\partial \varepsilon} = \xi^i(\tilde{x}), \quad \tilde{x}(0) = x. \tag{2.5}$$

Solving (2.5) to obtain the transformations τ_ε is often referred to as *exponentiating* $\mathbf{X}$. In particular, using (2.5) one can show that the power series expansion of τ_ε is given by

$$\tau_\varepsilon(x) = \exp(\varepsilon\mathbf{X})\,x = \Big(1 + \varepsilon\mathbf{X} + \frac{\varepsilon^2}{2}\mathbf{X}^2 + \cdots\Big)\,x.$$

Example 2.1.5 Let $M = \mathbb{R}$ with coordinates x and consider the vector field $\mathbf{X} = x^2\partial_x$. Then the corresponding one parameter family of transformations is given by

$$\tau_\varepsilon(x) = \exp(\varepsilon\mathbf{X})x = \frac{x}{1 - \varepsilon x}. \tag{2.6}$$

Notice that when $x > 0$, then τ_ε is defined only for $\varepsilon \in I_x$ where $I_x = (-\infty, 1/x)$. $\square$

Using the properties (2.2) and (2.3) of flows, one can show that the transformations τ_ε satisfy the axioms of a group. In particular, the identity element is τ_0, the binary operation between transformations is given by composition, $\tau_\varepsilon \circ \tau_\delta = \tau_{\delta+\varepsilon}$, which is clearly associative. Lastly the inverse of τ_ε is just $\tau_{-\varepsilon}$. Let us now explore the group structure of such transformations.

2.1.2 Lie Groups

Definition 2.1.6 A *group* is a set G together with a binary operation $* : G \times G \to G$ satisfying the following axioms:

Associativity:	For all $g, h, k \in G$, $g * (h * k) = (g * h) * k$.
Identity Element:	There exists an element $e \in G$, called the identity element, such that for all $g \in G$, $e * g = g = g * e$.
Inverses:	For each $g \in G$, there exists an inverse, denoted by g^{-1}, such that $g * g^{-1} = e = g^{-1} * g$.

Example 2.1.7 An example of a group is the set of integers $\mathbf{Z}$ together with addition as the associative binary operation. The identity element is 0 and the inverse of each integer $j \in \mathbf{Z}$ is $-j$. One often denotes this group by $(\mathbf{Z}, +)$. Other examples of groups are $(\mathbb{R}, +)$, the set of real numbers with addition, and $(\mathbb{R}^+, \cdot)$, the set of positive real numbers with multiplication. $\square$

Unlike, $(\mathbf{Z}, +)$, the group $(\mathbb{R}, +)$ has a binary operation which is an analytic map $+ : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$. Such groups are called Lie groups.

Definition 2.1.8 An r-parameter *Lie group* is a group $(G, *)$ which is also an r-dimensional differentiable manifold M such that both the binary operation $*$ and the inversion map $\varepsilon \mapsto \varepsilon^{-1}$ are analytic.

Henceforth, M will denote an p-dimensional differentiable manifold.

Definition 2.1.9 Let $(G, *)$ be an r-parameter Lie group. An r-parameter *Lie Group of transformations* is a collection of transformations $\mathcal{G}$ on M, together with a map $\tau : G \to \mathcal{G}$, which maps ε to τ_ε, such that

- τ_e is the identity map;
- $\tau_\delta \circ \tau_\varepsilon = \tau_{\delta * \varepsilon}$;
- $(\tau_\varepsilon)^{-1} = \tau_{\varepsilon^{-1}}$;
- $\tau_\varepsilon(x)$ is analytic in x and ε;
- The binary operation $*$ is analytic in both of its arguments.

In general, we are interested in *local* Lie groups of transformations where the transformations τ_ε may only be defined when ε is sufficiently close to the identity e, which is taken to be zero. Without loss of generality, we assume that any transformations considered are sufficiently close to the identity transformation. The need for the analyticity requirements for Lie groups of transformations is so that we can relate these to flows of vector fields. This will become clear shortly.

We have mentioned that flows of vector fields lead to families of transformations which form a local Lie group. In this regard, one can make a more precise statement:

Theorem 2.1.10 *Given any vector field* $\mathbf{X}$ *on* M*, the transformations* $\tau_\varepsilon = \exp(\varepsilon\mathbf{X})$ *form a one-parameter local Lie group of transformations acting on* M*.*

As already mentioned, one-parameter Lie groups of transformations arising as flows of a vector fields have a very simple composition law (cf. (2.2)). However, Lie groups of transformations can have more complicated composition laws in general.

Example 2.1.11 An example of a one-parameter Lie group of transformations on $M = \mathbb{R}^2$ is given by

$$\tau_\varepsilon \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} (1+\varepsilon)x \\ (1+\varepsilon)^2 y \end{pmatrix}, \qquad \varepsilon \in I = (-1, \infty). \tag{2.7}$$

The underlying Lie group is $(I, *)$, where the group binary operation is given by

$$\varepsilon_1 * \varepsilon_2 = \varepsilon_1 + \varepsilon_2 + \varepsilon_1\varepsilon_2, \qquad \varepsilon_1, \varepsilon_2 \in I, \tag{2.8}$$

the identity is zero, and the inverse $\varepsilon^{-1} = -\varepsilon/(1+\varepsilon)$ for all $\varepsilon \in I$. Clearly the axioms of Definition 2.1.9 are satisfied for τ_ε. $\square$

Given any one-parameter local Lie group $\mathcal{G}$ of transformations, Lie [40] proved that there always exists a re-parametrization (a particular choice of local coordinates for the underlying Lie group G) such that $\mathcal{G}$ becomes equivalent to the flow of some vector field:

Theorem 2.1.12 *Consider a one-parameter local Lie group of transformations $\mathcal{G}$ on M, given by $\widetilde{x} = \tau_\varepsilon(x)$ where $\tau_\varepsilon = (\tau_\varepsilon^1, \cdots, \tau_\varepsilon^p)$. Form the associated vector field $\mathbf{X} = \xi^i(x)\,\partial_{x_i}$, where*

$$\xi^i(x) = \left.\frac{\partial}{\partial\varepsilon}\tau_\varepsilon^i(x)\right|_{\varepsilon=0}. \tag{2.9}$$

Then there exists a reparametrization $\varepsilon = \varepsilon(\delta)$ such that $\tau_{\varepsilon(\delta)}(x) \equiv \exp(\delta\mathbf{X})x$.

The proof of this theorem, which can be found in [15], relies on the analyticity assumptions in Definition 2.1.9 in order to construct certain power series expansions. The re-parametrization $\varepsilon = \varepsilon(\delta)$ is a different choice of local coordinates for the Lie group G, such that the group binary operation in the original coordinates ε becomes addition in the new coordinates δ.

Definition 2.1.13 Consider a one-parameter Lie group of transformations τ_ε on M with *infinitesimals* $(\xi^1, \cdots, \xi^p)$, given by (2.9). Each ξ^i is called the *infinitesimal* of x_i and the corresponding vector field $\mathbf{X} = \xi^i\partial_{x_i}$ is called the *infinitesimal generator* of τ_ε.

Example 2.1.14 The infinitesimal generator associated with the one-parameter Lie group of transformations (2.7) is given by $\mathbf{X} = x\partial_x + 2y\partial_y$. The flow of $\mathbf{X}$ yields the one-parameter Lie group of transformations

$$\widetilde{\tau}_\delta\begin{pmatrix} x \\ y \end{pmatrix} = \exp(\delta\mathbf{X})\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} e^\delta x \\ e^{2\delta}y \end{pmatrix}, \qquad \delta \in (-\infty, \infty).$$

The re-parametrization that relates τ_ε to $\widetilde{\tau}_\delta$ is given by $\varepsilon = e^\delta - 1$. $\square$

Theorem 2.1.12 can be generalized to an r-parameter group of transformations $\mathcal{G}$ as follows. Let the Lie group $(G, *)$ have local coordinates $\varepsilon = (\varepsilon_1, \cdots, \varepsilon_r) \in G$ and $\tau_\varepsilon \in \mathcal{G}$. Form the set of r associated infinitesimal generators

$$\mathbf{X}_\alpha = \xi^{i\alpha}(x)\,\partial_{x_i}, \qquad \alpha = 1, \cdots, r,$$

where the infinitesimals $\xi^{i\alpha}$ are given by

$$\xi^{i\alpha} = \left.\tfrac{\partial}{\partial \varepsilon_\alpha}\tau^i_\varepsilon(x)\right|_{\varepsilon=0}.$$

Each transformation $\tau_\varepsilon \in \mathcal{G}$ can be realized as either

$$\exp(\lambda_1 \mathbf{X}_1)\exp(\lambda_2 \mathbf{X}_2)\cdots\exp(\lambda_r \mathbf{X}_r),$$

or

$$\exp(\delta \mathbf{X}), \tag{2.10}$$

where

$$\mathbf{X} = \mu_1 \mathbf{X}_1 + \mu_2 \mathbf{X}_2 + \cdots + \mu_r \mathbf{X}_r, \tag{2.11}$$

for all real numbers $\lambda_i, \delta, \mu_\alpha$, with λ_i, δ sufficiently small. Hence, the r infinitesimal generators $\mathbf{X}_\alpha$ contain all the information needed to reconstruct the r-parameter Lie group of transformations $\mathcal{G}$. (See [15] for more details.) Henceforth we will refer to any Lie group of transformations as simply a Lie group.

It is natural to consider the r-dimensional vector space $\mathcal{L}$ with basis given by the infinitesimal generators $\{\mathbf{X}_1, \cdots, \mathbf{X}_r\}$ over the real numbers since for any $\mathbf{X} \in \mathcal{L}$, given by (2.11), the corresponding transformation (2.10) lies in $\mathcal{G}$ for sufficiently small δ. It turns out that $\mathcal{L}$ is endowed with an additional algebraic structure which we shall now investigate.

2.1.3 Lie Algebras

Definition 2.1.15 A *Lie Algebra* $\mathcal{L}$ is a vector space over some field $\mathcal{F}$ with an additional binary operation (*the commutator*), such that for all $a, b \in \mathcal{F}$ and $\mathbf{X}, \mathbf{Y}, \mathbf{Z} \in \mathcal{L}$, the following properties are satisfied:

$$\begin{array}{ll} \textit{Closure.} & [\mathbf{X},\mathbf{Y}] \in \mathcal{L}; \\ \textit{Bilinearity.} & [\mathbf{X}, a\mathbf{Y} + b\,\mathbf{Z}] = a[\mathbf{X},\mathbf{Y}] + b[\mathbf{X},\mathbf{Z}]; \\ \textit{Anticommutativity.} & [\mathbf{X},\mathbf{Y}] = -[\mathbf{Y},\mathbf{X}]; \\ \textit{Jacobi Identity.} & [\mathbf{X},[\mathbf{Y},\mathbf{Z}]] + [\mathbf{Y},[\mathbf{Z},\mathbf{X}]] + [\mathbf{Z},[\mathbf{X},\mathbf{Y}]] = 0. \end{array} \tag{2.12}$$

Given any finite dimensional Lie algebra $\mathcal{L}$ with basis $\{\mathbf{X}_1, \cdots, \mathbf{X}_r\}$, Lie showed that the closure property leads to

$$[\mathbf{X}_\alpha, \mathbf{X}_\beta] = C^\gamma_{\alpha\beta}\mathbf{X}_\gamma \qquad \text{where} \qquad \alpha, \beta, \gamma = 1, \cdots, r,$$

for some constants $C^\gamma_{\alpha\beta}$, called the structure constants of $\mathcal{L}$. The anticommutativity property leads to

$$C^\gamma_{\alpha\beta} = -C^\gamma_{\beta\alpha};$$

and the Jacobi identity leads to

$$C^\rho_{\alpha\beta}C^\delta_{\rho\gamma} + C^\rho_{\beta\gamma}C^\delta_{\rho\alpha} + C^\rho_{\gamma\alpha}C^\delta_{\rho\beta} = 0.$$

We will be confining ourselves to the case where $\mathcal{L}$ is the vector space given by a basis of vector fields $\{\mathbf{X}_1, \cdots, \mathbf{X}_r\}$, $\mathcal{F}$ is the field of real numbers and the commutator is defined as follows:

Definition 2.1.16 Let $\mathbf{X} = \xi^i \frac{\partial}{\partial x^i}$ and $\mathbf{Y} = \gamma^i \frac{\partial}{\partial x^i}$ be two vector fields. Their *commutator* is the vector field

$$[\mathbf{X},\mathbf{Y}] = \mathbf{XY} - \mathbf{YX} = \left(\xi^j \frac{\partial \gamma^i}{\partial x^j} - \gamma^j \frac{\partial \xi^i}{\partial x^j}\right) \frac{\partial}{\partial x^i}\,. \tag{2.13}$$

Can any set of vector fields $\{\mathbf{X}_1, \cdots, \mathbf{X}_r\}$ define a Lie algebra? One can easily verify that the commutator bracket (2.13), by its very definition, satisfies the bilinearity and anticommutativity properties as well as the Jacobi identity. However, it is not true in general that the closure property holds for any set of vector fields. For example, $\{\mathbf{X}_1 = \partial_x, \mathbf{X}_2 = x\partial_y\}$ is not closed under the commutator bracket. However the closure property does hold for infinitesimal generators of an r-parameter Lie group.

Theorem 2.1.17 *The infinitesimal generators $\{\mathbf{X}_\alpha\}$, $\alpha = 1, \cdots, r$, of an r-parameter Lie group $\mathcal{G}$ form an r-dimensional Lie algebra $\mathcal{L}^r$.*

This follows from the fact that $\mathcal{G}$ is closed under composition of its members. In particular, given any two infinitesimal generators $\mathbf{X}_\alpha$ and $\mathbf{X}_\beta$ of $\mathcal{G}$, their corresponding transformations $\exp(\varepsilon\mathbf{X}_\alpha)$ and $\exp(\varepsilon\mathbf{X}_\beta)$ both belong to $\mathcal{G}$ and hence so does

$$\tau_\varepsilon = \exp(-\sqrt{\varepsilon}\mathbf{X}_\beta)\exp(-\sqrt{\varepsilon}\mathbf{X}_\alpha)\exp(\sqrt{\varepsilon}\mathbf{X}_\beta)\exp(\sqrt{\varepsilon}\mathbf{X}_\alpha),$$

for ε sufficiently small. By Theorem 2.1.12 the infinitesimal generator corresponding to τ_ε is given by

$$\left.\frac{\partial\tau_\varepsilon}{\partial\varepsilon}\right|_{\varepsilon=0+} = [\,\mathbf{X}_\alpha, \mathbf{X}_\beta\,].$$

Consequently, the infinitesimal generators of $\mathcal{G}$ are closed under commutation.

2.1.4 Symmetries of Algebraic Equations

Consider the system of *algebraic equations* defined on M:

$$F_\mu(x) = 0, \qquad \mu = 1, \cdots, l, \tag{2.14}$$

where $l \leq p$ and $F \in C^\infty(M)$. The *solutions* of (2.14) are given by the locus of points

$$\varrho_F = \{x \in M \colon F(x) = 0\} \subset M.$$

Definition 2.1.18 A *symmetry* of a system of equations (2.14) is a transformation τ mapping any solution to another solution of the system:

$$\tau(\varrho_F) \subset \varrho_F,$$

where ϱ_F is the locus of solutions of the system. A Lie group $\mathcal{G}$ acting on M is a *symmetry group* of the system, if and only if for all $\tau_\varepsilon \in \mathcal{G}$, τ_ε is a symmetry of the system, for ε sufficiently small.

In other words, $\mathcal{G}$ is a symmetry group of the system if and only if for all $\tau_\varepsilon \in \mathcal{G}$,

$$F_\mu(\tau_\varepsilon(x)) = 0 \qquad \text{whenever} \qquad x \in \varrho_F.$$

These conditions are, in general, nonlinear in τ_ε and hence very difficult to analyse. Using the infinitesimal generators of $\mathcal{G}$, an equivalent statement for these symmetry conditions can be made which turns out to be more useful in practice. As we shall see, the infinitesimal formulation of the symmetry conditions will be indispensable in the case of DEs. This was the fundamental insight due to Lie [40]. Before stating the infinitesimal symmetry conditions, one must first define what is meant by a system of equations having maximal rank.

Definition 2.1.19 The system of equations (2.14) on M is of *maximal rank* if and only if the Jacobian matrix $(\partial F_\mu/\partial x_i)$ is of rank l at every solution $x \in \varrho_F$. If a system is of maximal rank, then it is said to satisfy the *maximal rank condition.*

The following theorem and the subsequent example are from [47, §2.1].

Theorem 2.1.20 (Infinitesimal Symmetry Conditions) *Consider the system (2.14) with locus of solutions* ϱ_F *and which is of maximal rank. Then a Lie group* $\mathcal{G}$ *acting on* M *is a symmetry group of the system if and only if every infinitesimal generator* $\mathbf{X}$ *of* $\mathcal{G}$ *satisfies*

$$\mathbf{X}[F_\mu(x)] = 0,\ \mu = 1, \cdots, l, \qquad \forall\, x \in \varrho_F. \tag{2.15}$$

Example 2.1.21 Let $\mathcal{G} = SO(2)$ be the one parameter rotation group in the plane $M = \mathbb{R}^2$, with infinitesimal generator $\mathbf{X} = -y\partial_x + x\partial_y$ and consider the algebraic equation

$$F(x, y) = x^4 + x^2y^2 + y^2 - 1 = 0.$$

Clearly (2.15) holds since

$$\mathbf{X}(F) = -2xy(x^2 + 1)^{-1}F(x, y).$$

In addition, the maximal rank condition is satisfied since the Jacobian

$$\left(\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y}\right) = (4x^3 + 2xy^2, 2x^2y + 2y),$$

vanishes only at $(x, y) = (0, 0)$ which is not a solution of $F(x, y) = 0$. By Theorem 2.1.20, $SO(2)$ is a symmetry group of $F(x, y) = 0$. Indeed,

$$F(x, y) = (x^2 + 1)(x^2 + y^2 - 1) = 0,$$

has S^1, the unit circle with origin for centre, as the solution set; rotations in the plane map points in S^1 to other points in S^1. $\square$

Example 2.1.22 To illustrate the importance of the maximal rank condition, let $M = \mathbb{R}$, $\mathcal{G}$ be the one-parameter group of translations with infinitesimal generator $\mathbf{X} = \partial_x$ and consider the equation $F(x) = (x-1)^2 = 0$ which has locus of solutions $\varrho_F = \{x : x = 1\}$. We have $\mathbf{X}(F) = 2(x-1)$ so that (2.15) is satisfied. However, we know for certain that $\mathcal{G}$ is not a symmetry group of ϱ_F. The problem is that the Jacobian $\partial F/\partial x = 2(x-1)$ vanishes on ϱ_F, so that the maximal rank condition does not hold and Theorem 2.1.20 does not apply. Of course, in this example one could have simplified the original equation to $x - 1 = 0$ since both have the same locus of solutions ϱ_F. The new equation does satisfy the maximal rank condition, but now (2.15) does not hold and one concludes that $\mathcal{G}$ is not a symmetry group of $F(x) = 0$. However, the maximal rank condition is invaluable when considering more complicated systems where it may be unclear whether such simplifications are needed in order to apply Theorem 2.1.20. □

The following theorem will prove useful later:

Theorem 2.1.23 *Consider the system (2.14) with solutions $\varrho_F \subset M$ and symmetry group $\mathcal{G}$. Suppose H is a function on M such that $H(x) = 0$, $\forall x \in \varrho_F$. Then*

$$\mathbf{X}[H(x)] = 0, \qquad \forall x \in \varrho_F.$$

Proof. Since $\mathcal{G}$ is a symmetry group of the system, given any $x \in \varrho_F$ and $\tau_\varepsilon \in \mathcal{G}$, we have $\tau_\varepsilon(x) \in \varrho_F$ for ε sufficiently small. In other words, $H(\tau_\varepsilon(x)) = 0$. Now τ_ε is given by exponentiating some infinitesimal generator $\mathbf{X}$ in the Lie algebra of $\mathcal{G}$. Hence, we have

$$0 = H(\tau_\varepsilon(x)) = H(\exp(\varepsilon \mathbf{X})\, x), \qquad \forall x \in \varrho_F.$$

Differentiating this with respect to ε and setting $\varepsilon = 0$ leads to the desired result. □

2.2 Differential Equations

It is natural to consider a system R of n-th order DEs as a system of algebraic equations in *n-jet space*, i.e., the space of independent variables, dependent variables and derivatives of dependent variables up to order n. A principle motivation for doing this is that the symmetry conditions for algebraic equations have already been determined in the last section. How does the group thus calculated relate to the symmetries of R? By passing to the algebraic regime, we lose the various relationships between a dependent variable and its derivatives. Without any a priori restrictions on the transformations considered, the algebraic symmetry conditions may result in transformations that do not map analytic functions to analytic functions, let alone mapping solutions to other solutions of R. Consequently, we are led to consider only certain types of transformations (Lie groups of extended transformations) when applying the algebraic symmetry conditions. It turns out that we will also require the system to be locally solvable. Let us first review some elements of n-jet space.

2.2.1 Jet Space

Let $X = \mathbb{R}^p$, with coordinates $x = (x_1, x_2, \cdots, x_p)$, be the space representing the independent variables, and let $U = \mathbb{R}^q$, with coordinates $u = (u^1, u^2, \cdots, u^q)$, be the space representing the dependent variables. The space $M = X \times U$ is called the *base space*. Of particular interest are functions $u = f(x)$, which are identified with their graphs

$$\Gamma_f = \{(x, f(x)) : x \,\epsilon\, \Omega\} \subset X \times U,$$

where $\Omega \subset X$ is the domain of f. Consider the k-th order partial derivatives of $f(x)$, given by

$$\partial_J f^\alpha(x) = \frac{\partial^k f^\alpha(x)}{\partial x_{j_1} \partial x_{j_2} \cdots \partial x_{j_k}},$$

where the multi-index $J = (j_1, \cdots, j_k)$ is an unordered k-tuple of integers $1 \le j_1, \cdots, j_k \le p$ and the order of J, denoted by $|J|$, is the number of elements in J (k in this case). There are $q \cdot p_k$ such partial derivatives, where

$$p_k = \binom{p+k-1}{k}.$$

In order to build a space to represent these derivatives of f, one *extends* or *prolongs* the base space as follows. Let $U_k = \mathbb{R}^{q \cdot p_k}$ have coordinates u_J^α, for all $\alpha = 1, \cdots, q$ and for all $J = (j_1, \cdots, j_k)$, so that $U^{(n)} = U \times U_1 \times \cdots \times U_n$ represents the space of all partial derivatives of $u = f(x)$ up to order n. Also, let $u^{(n)}$ denote a typical point in $U^{(n)}$. By convention, $u_0^\alpha = u^\alpha$ is called the zeroth order derivative of u^α, $U^{(0)} = U_0 = U$, and J is given by the single multi-index 0 when $k = 0$. The dimension of $U^{(n)}$ is $q \cdot p^{(n)}$ where

$$p^{(n)} = p + p_1 + p_2 + \cdots + p_n = \binom{p+n}{n}.$$

Example 2.2.1 Consider the case $p = 2$, $q = 1$ so that $X = \mathbb{R}^2$ has coordinates $(x_1, x_2) = (t, x)$, and $U = \mathbb{R}$ has the single coordinate u. Now U_1 is isomorphic to $\mathbb{R}^2$ with coordinates (u_t, u_x) representing all first order partial derivatives of u with respect to t and x. Likewise, $U_2 \simeq \mathbb{R}^3$ with coordinates (u_{tt}, u_{tx}, u_{xx}) represents all second order partial derivatives of u. Lastly, $U^{(2)} \simeq \mathbb{R}^6$ with coordinates $u^{(2)} = (u, u_t, u_x, u_{tt}, u_{tx}, u_{xx})$, is the space of all partial derivatives of u up to order 2. $\square$

The space $X \times U^{(n)}$ is called the n-th *extension space* or the n-th order *jet space* of the underlying base space $X \times U$. It represents the space of all independent variables, dependent variables and all derivatives of dependent variables up to order n. To graph a function $u = f(x)$ in this extension space, one must calculate the values of all the partial derivatives of f up to order n, in the domain of f. One then forms the induced function $u^{(n)} = \mathrm{pr}^{(n)} f(x)$, called the *$n$-th prolongation* of f, defined by the equations

$$u_J^\alpha = \partial_J f^\alpha(x).$$

The extended or prolonged graph is then given by

$$\mathrm{pr}^{(n)}\Gamma_f = \Gamma_f^{(n)} = \{(x, \mathrm{pr}^{(n)} f(x))\} \subset X \times U^{(n)}.$$

Example 2.2.2 Consider the case $p = 1$, $q = 1$ so that $X \times U^{(1)} \simeq \mathbb{R}^3$ with coordinates (x, u, u_x). The graph of the function $f(x) = \sin x$ is $\Gamma_f = \{(x, \sin x) : x \in \mathbb{R}\}$, and the first prolonged graph is given by

$$\Gamma_f^{(1)} = \{(x, \sin x, \cos x) \colon x \in \mathbb{R}\} \subset X \times U^{(1)}.$$

Also, consider the following locus of points

$$\varrho = \{(x, \sin x, 1) \colon x \in \mathbb{R}\} \subset X \times U^{(1)}.$$

ϱ does not correspond to the extension of any graph $u = f(x)$. $\square$

The above example illustrates that any arbitrary locus of points $\varrho \subset X \times U^{(n)}$ will not in general be the prolonged graph of some function $u = f(x)$. However, given any point in n-jet space, one can always find an n-th prolonged graph of some function that passes through the point.

Theorem 2.2.3 *Given any point $P(x, u^{(n)}) \in X \times U^{(n)}$, there is an n-th prolonged graph $\Gamma_f^{(n)}$ of some function $u = f(x)$ that passes through P.*

The proof of this theorem relies on constructing a Taylor polynomial of order n. Here, we will just illustrate the case $n = 1$, with $X = \mathbb{R}^2$ and $U = \mathbb{R}$. Let X and U have coordinates (x, t) and u respectively. Then given any point $P(x_0, t_0, u(x_0, t_0), u_x(x_0, t_0), u_t(x_0, t_0)) \in X \times U^{(1)}$, the Taylor polynomial

$$f(x, t) = u(x_0, t_0) + u_x(x_0, t_0)(x - x_0) + u_t(x_0, t_0)(t - t_0), \tag{2.16}$$

has the first prolonged graph $\Gamma_f^{(1)}$ passing through P.

There are two types of differentiations that can be performed on $X \times U^{(n)}$, depending upon how one views this space. *Partial differentiation* involves the partial differential operators ∂_{x_i} and $\partial_{u_J^\alpha}$, which treat $X \times U^{(n)}$ as just Euclidean space $\mathbb{R}^N$ of dimension $N = p + q \cdot p^{(n)}$. Since no relationships between any of the coordinates $(x, u^{(n)})$ are assumed, $\partial_{x_i} g(x, u^{(n)})$ is calculated while keeping fixed all the coordinates except for x_i. A similar statement can be made for $\partial_{u_J^\alpha}$. *Total differentiation* on the other hand, involves the total derivative operators D_{x_i}, which respect the various relationships between the coordinates of $X \times U^{(n)}$. In particular, the total derivative operators are defined through the following identity:

$$\left(D_{x_i} g(x, u^{(n)})\right)\Big|_{\Gamma_f^{(n+1)}} \equiv \partial_{x_i}\left(g(x, u^{(n)})|_{\Gamma_f^{(n)}}\right). \tag{2.17}$$

Example 2.2.4 Let $X = \mathbb{R}^2$ and $U = \mathbb{R}$ with coordinates (x, t) and u respectively. The function $g(x, u^{(1)}) = x u_x^2$, has the following partial derivatives:

$$\partial_t g = 0, \qquad \partial_x g = u_x^2, \qquad \partial_u g = 0, \qquad \partial_{u_t} g = 0, \qquad \partial_{u_x} g = 2x u_x.$$

The total derivative of g with respect to t is calculated as follows:

$$\Big(D_t g\Big)\Big|_{\Gamma_f^{(2)}} = \partial_t\Big(x f_x^2(x,t)\Big) = 2x f_x f_{xt}.$$

This together with a similar calculation for $D_x g$ gives

$$D_t g = 2x u_x u_{xt}, \qquad D_x g = u_x^2 + 2x u_x u_{xx}.$$

$\square$

One can easily show that the total derivative operator D_{x_i} is given by

$$D_{x_i} = \partial_{x_i} + u_i^\alpha \partial_{u^\alpha} + \cdots + u_{Ji}^\alpha \partial_{u_J^\alpha} + \cdots, \tag{2.18}$$

where summation over the repeated indices $\alpha = 1, \cdots, q$, $J = (j_1, \cdots, j_k)$ and $Ji = (j_1, \cdots, j_k, i)$ is assumed. This is an infinite sum, but when applied to functions $g(x, u^{(n)})$, only a finite number of terms are needed. It is clear that this identity yields the same expressions for $D_t g$ and $D_x g$ as given in Example 2.2.4. The advantage of using this identity for D_{x_i} is that one avoids having to make the replacements $u^{(n)} = \mathrm{pr}^{(n)} f(x)$ in (2.17). It is often convenient to use D_i to mean D_{x_i} and D_J to denote the k-th order total derivative operator $D_{j_1} D_{j_2} \cdots D_{j_k}$.

The total derivative operators D_i encode the relationships between the various coordinates of $X \times U^{(n)}$, namely

$$u_J^\alpha = D_J(u^\alpha).$$

The prolongation structure of n-jet space is discussed in detail in [60]. For our purposes, it will be sufficient to view $X \times U^{(n)}$ as $(\mathbb{R}^N, D_i)$, i.e., the Euclidean space of dimension $N = p + q \cdot p^{(n)}$ together with the prolongation structure encoded in the total derivative operators D_i. We are now ready to discuss what we mean by a system of differential equations.

Definition 2.2.5 A system R of n-th order DEs is given by the system of algebraic equations on n-jet space:

$$\Delta_\mu(x, u^{(n)}) = 0, \qquad \mu = 1, \cdots, l, \tag{2.19}$$

where Δ_μ are real valued functions on $X \times U^{(n)}$. A solution of the DE (2.19) is a function $u = f(x)$ such that

$$\Delta(x, u^{(n)})\Big|_{\Gamma_f^{(n)}} = 0. \tag{2.20}$$

We must emphasize that by a solution of R, one really means a function $u = f(x)$ such that, after replacing u and its derivatives by $f(x)$ and its corresponding derivatives, Δ evaluates to zero for all x in the domain of $f(x)$. Having said this, it is often convenient to consider the individual points in $X \times U^{(n)}$ that satisfy (2.19):

Definition 2.2.6 [47, p.96*ff*] Let ϱ_Δ be the locus of points (subvariety) in n-jet space given by

$$\varrho_\Delta = \{(x, u^{(n)})\colon \Delta(x, u^{(n)}) = 0\} \subset X \times U^{(n)}.$$

In other words, ϱ_Δ consists of the roots of the algebraic equations $\Delta : X \times U^{(n)} \to \mathbb{R}^l$.

It is natural to ask what the relationship is between the solutions of R and the locus of points ϱ_Δ. From (2.20), the extended graphs of solutions of R must lie in ϱ_Δ:

Theorem 2.2.7 *Let R be an n-th order system of DEs. Then $u = f(x)$ is a solution of R if and only if its n-th extended graph lies in ϱ_Δ:*

$$\Gamma_f^{(N)} \subset \varrho_\Delta. \tag{2.21}$$

Example 2.2.8 Let $X = U = \mathbb{R}$ and R be the ODE, $\Delta = u_x + u^2 = 0$. The general solution of R is $f(x) = \frac{1}{x-c}$, for any $c \in \mathbb{R}$, since $\Delta(x, f(x), f'(x))$ evaluates to zero for all $x \neq c$. One also has

$$\varrho_\Delta = \{(x, u, -u^2)\colon x, u \in \mathbb{R}\} \subset X \times U^{(1)};$$

$\Gamma_f^{(1)}$ given by the functions

$$u = (x - c)^{-1}, \qquad u_x = -(x - c)^{-2};$$

and, as seen in Figures 2.1, (2.21) holds. $\square$

In general, given any point P in ϱ_Δ there is no guarantee that there exists an extended graph of some solution $u = f(x)$ of R passing through P. Theorem 2.2.3 still applies in that one can always find an extended graph of some function $u = g(x)$ passing through P. However, $u = g(x)$ will not, in general, be a solution of R.