

Is it possible matching higher-twist contributions with chiral perturbation theory?

D.A. Volkova¹, N.A Gramotkov¹, O.V. Teryaev¹ and O.P. Solovtsova²

Joint Institute for Nuclear Research, Dubna, Russia

Sukhoi State Technical University of Gomel, Gomel, Belarus

Using the example of the QCD analysis of the precise low-energy data on the polarized Bjorken sum rule, we show how it is possible to make a transition between two regimes of the QCD expansions down to low Q^2 scales. As it is well known, at low momentum transfer, $Q < 1 \sim \text{GeV}$, the description of the perturbative part of the Bjorken integral in the framework of standard perturbation theory encounters serious difficulties due to unphysical features of the usual perturbative running coupling. To avoid this difficulty, we use an analytic running coupling which, without introducing additional parameters, eliminates the unphysical features of the perturbative coupling. Since the theoretical description of the Bjorken sum rule involves not only a series in powers of α_s but also a series in powers of $1/Q^2$ (higher twist contributions summing into an unknown function), we will use the technique of matching the function at large Q^2 and behavior at small Q^2 near zero by involving the Gerasimov--Drell--Hearn sum rule. The essence of the "matching" method is that the sum rule is applied to the region of large values of Q^2 , where it works well, and then continues to the region of small Q^2 . This allows us to obtain information about the behavior of structural functions in the region where experimental data are not available. The region near $Q^2 = 0$ is of particular interest because it corresponds to small distances between interacting particles. In this region, the structural functions can experience significant changes associated with the manifestation of the inner degrees of freedom of hadrons. A qualitative description of the region near $Q^2 = 0$ can be obtained by analyzing the behavior of structural functions at small values of Q^2 . This approach in different loop level of the perturbative part gives a stable good agreement with the experimental data in the whole region up to zero momentum transfer.