

Observation of New Modes of Multi-Body Decays of $^{252}\text{Cf(sf)}$

Yu.V. Pyatkov^{1,2}, D.V. Kamanin², Z.I. Goryainova², E.A. Kuznetsova²,
A.N. Solodov², O.V. Strelakovsky², V.E. Zhuchko³, A.O. Zhukova², S.M. Wyngaardt³

¹*National Research Nuclear University “MEPhI”, 115409 Moscow, Russia*

²*Joint Institute for Nuclear Research, 141980 Dubna, Russia*

³*University of Stellenbosch, Stellenbosch, Western Cape, South Africa*

In our previous publications [1-4], we discussed various manifestations of the decay channel of low excited heavy nuclei, called collinear cluster tri-partition (CCT). New modes of ternary and likely quaternary decays of $^{252}\text{Cf(sf)}$ were observed using the “double-hit” approach. The experiments were performed at the COMETA, a double-armed, mosaic, time-of-flight spectrometer of fission fragments [5]. Digital images of all the detector signals were obtained using multichannel fast flash-digitizer. Off-line processing of the recorded data allowed us to select the decay events where two fragments were detected in the same PIN diode during the time-selection gate of 200 ns. For the selected events, the precission configuration of the mother nucleus seems to be a channel consisting of different magic nuclei.

References

1. Yu.V. Pyatkov et al., Eur. Phys. J. A **45**, 29 (2010).
2. Yu.V. Pyatkov et al., Eur. Phys. J. A **48**, 94 (2012).
3. Yu.V. Pyatkov et al., Phys. Rev. C **96** (2017) 064606.
4. Yu.V. Pyatkov et al., Eurasian Journal of Physics and Functional Materials, v.4, №1 (2020) 13–18.
5. Yu.V. Pyatkov et al., Eur. Phys. J. A **48**, 94 (2012).