

Объединенный
институт
ядерных
исследований
Дубна

E1-85-547


V.M.Bystritsky, J.Wozniak

USE OF A LARGE-VOLUME
NE-213 SCINTILLATOR
FOR INVESTIGATIONS
OF THE $\pi\mu^- \rightarrow {}^4\text{He} + 2n + \mu^-$ REACTION

Submitted to "Acta Phys. Polonica"

1985

By now the reaction



is not studied well. Only two complete kinematic experiments have been carried out at the energies 1.39 MeV ^{1/1} and 40 keV ^{1/2} with ambiguous results as to interpretation of the role of (n-n) and (α -n) interactions between reaction products in the final state. Noteworthy is that the initial orbital moment of this system $L=0$. In this case three reaction channels are possible:


1) production of the ${}^6\text{He}$ compound nucleus followed by its break-up into an α -particle and two neutrons;

2) production of ${}^5\text{He}$ in the ground state $P_{3/2}$ and a neutron;

3) production of ${}^5\text{He}$ in the excited state $P_{1/2}$ and a neutron.

For the triton energy 0.5 MeV and $\theta = 90^\circ$ the branching ratio for those channels is 0.7:0.2:0.1 ^{1/3}.

The investigations of the $tt\mu^-$ -molecule formation followed by the synthesis in it



showed that, unlike reaction (1), reaction (2) starts from the state $L=1$ and, consequently, the relation between the reaction channels may change. However, it is possible to obtain information on contributions of (n-n) and (α -n) interactions between the reaction products in the final state. We believe it is possible by comparison of amplitude spectra of registered neutrons with the simulated distributions

obtained with the help of assumed matrix element which describes reaction (2).

In Ref. ^{1/4} the neutron registration efficiency has been calculated for reaction (2) using a 1-litre detector with the NE-213 scintillator, and amplitude distributions have been plotted with allowance for "pure" (n-n) and (α -n) interactions in the final state (chosen as boundary cases). They are given in




Fig. 1. Amplitude distributions of neutrons for reaction (2), calculated for the detector with 1 l of the NE-213 scintillator.

Fig. 1. As is seen, no definite conclusions on the relation between (n-n) and (α -n) interactions are possible. To answer the question, whether the amplitude distributions of registered neutrons allow in general information on the mechanism of reaction (2), we have performed the necessary calculations using a neutron detector of practically full absorption with 200 l of the NE-213 scintillator.

Fig. 2. shows the distributions of the total energy of two neutrons produced in reaction (2) for (n-n) and (α -n) interactions. In Fig. 3 one can see the calculated amplitude distributions obtained with the help of a 200 l detector for those two variants of reaction (2) product interactions in the final state. The calculations were performed using the Monte-Carlo method. The algorithm for the calculations and the geometrical dimensions of the neutron detector are given in Ref. ^{15/}.

The differential cross section for reaction (2) where two particles i,j interact strongly in the final state can be written down in a simplified form:

$$d\sigma = \text{const } F_{ij}^2(k_{ij}) dR, \quad (3)$$

where dR is the reaction phase space element; $\hbar k_{ij}$ is the rela-

tive momentum of the particles i,j; F_{ij}^2 is the enhancement factor, describing the particle interaction in the final state.

The enhancement factor has been calculated in the Watson-Migdal approximation

$$F_{ij}^2 = k_{ij}^{-2} \cdot \sin^2 \delta_{ij}, \quad (4)$$

where δ_{ij} is the scattering phase shift.

For the (n-n) interaction the scattering phase shift has been parametrized using the effective radius approximation with the scattering length $a_{nn} = -17$ fm and the effective radius $r_0 = 2.65$ fm. For the (α -n) interaction the phase shift parametrization was taken from Ref. ^{16/}. The channel ratio $P_{3/2}:P_{1/2}$ was 1:1 in the calculations.

Comparison of Figs. 1 and 3 makes it obvious that the difference of hypotheses of (n-n) and (α -n) interactions becomes significant when a large-volume neutron detector is used. Besides, attention should be paid to the following fact: when the experimental spectrum coincides with the simulated one, the energy distribution of α -particles becomes known, which is very important for determination of the probability of the muon shaking-off from ${}^4\text{He}\mu$ produced in reaction (2). This is significant for the efficiency of the μ -catalysed fusion.

In conclusion we note that a large-volume scintillator detector can be employed for checking the theoretical description of reaction (2) mechanism.

The authors are grateful to V.S. Melezhhik for useful discussions.

References

1. Kühn B. et al. Nucl.Phys., A183 (1972) 640.
2. Iarose-Poutissou R. and Jeremie H. Nucl.Phys., A218 (1974) 559.
3. Ajzenberg-Selove F. and Lauritsen T. Nucl.Phys., A227 (1974) 1.
4. Bystritsky V.M. et al. Acta Phys.Polonica, B15 (1984) 689.
5. Bystritsky V.M. et al. JINR, E1-84-735, Dubna, 1984; to be published in Nuclear Instruments and Methods.
6. Arndt R.A. et al. Nucl.Phys., A209 (1973) 447.




Fig. 2. Event distribution for the reaction $tt\mu \rightarrow {}^4\text{He}+2n+\mu$ over the total energy of two neutrons.




Fig. 3. Amplitude distributions of neutrons for reaction (2), calculated for the detector with 200 l of the NE-213 scintillator.

В Объединенном институте ядерных исследований начал выходить сборник "Краткие сообщения ОИЯИ". В нем будут помещаться статьи, содержащие оригинальные научные, научно-технические, методические и прикладные результаты, требующие срочной публикации. Будучи частью "Сообщений ОИЯИ", статьи, вошедшие в сборник, имеют, как и другие издания ОИЯИ, статус официальных публикаций.

Сборник "Краткие сообщения ОИЯИ" будет выходить регулярно.

The Joint Institute for Nuclear Research begins publishing a collection of papers entitled *JINR Rapid Communications* which is a section of the JINR Communications and is intended for the accelerated publication of important results on the following subjects:

Physics of elementary particles and atomic nuclei.
Theoretical physics.
Experimental techniques and methods.
Accelerators.
Cryogenics.
Computing mathematics and methods.
Solid state physics. Liquids.
Theory of condensed matter.
Applied researches.

Being a part of the JINR Communications, the articles of new collection like all other publications of the Joint Institute for Nuclear Research have the status of official publications.

JINR Rapid Communications will be issued regularly.



Быстрицкий В.М., Возняк Я.

E1-85-547

Использование большого сцинтиллятора для исследования реакции $t\mu^- \rightarrow {}^4\text{He} + 2n + \mu^-$

Обсуждается вопрос использования детектора с большим объемом сцинтиллятора для исследования механизма реакции ядерного синтеза $t\mu^- \rightarrow {}^4\text{He} + 2n + \mu^-$ путем сравнения измеренных амплитудных распределений зарегистрированных нейтронов с расчетными распределениями нейтронов, полученных в предположении определенного вида матричного элемента реакции.

Работа выполнена в Лаборатории ядерных проблем ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1985

Bystritsky V.M., Wozniak J.

E1-85-547

Use of a Large-Volume NE-213 Scintillator for Investigations of the $t\mu^- \rightarrow {}^4\text{He} + 2n + \mu^-$ Reaction

Use of a large-volume NE-213 scintillator for investigations of the mechanism of the nuclear synthesis reaction $t\mu^- \rightarrow {}^4\text{He} + 2n + \mu^-$ is discussed. The investigations are supposed to be performed by comparison of the measured amplitude distributions of registered neutrons with the calculated neutron distributions obtained for an assumed form of the matrix element of the reaction.

The investigation has been performed at the Laboratory of Nuclear Problems, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1985