

**ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА**

E1-85-55

M.N.Khachaturian

**THIRTY-YEAR WORK
WITH LEAD-GLASS CHERENKOV
 γ -SPECTROMETERS AT DUBNA**

Submitted to All-Union Seminar
on Cherenkov Detectors and Their
Application in Science
and Technology (Moscow, 1984)

1985

One of the first Cherenkov γ -spectrometers was proposed and constructed at JINR in 1954-55 and intended for the detection of antiprotons and antineutrons at the 10 GeV proton synchrotron^{1/} (Fig.1).

To increase the energy resolution of lead-glass γ -spectrometers which is to a great extent dependent on the optical transparency of a radiator, in 1955-56 new lead-glasses were made from extra pure materials. This allowed the energy resolution of lead-glass γ -spectrometers to be considerably increased. In 1957-59 one of these counters was used in measurements of the Panofsky ratio^{2/} (Fig.2).

At the end of the fifties a great deal of information on the interaction cross sections of charged hadrons was obtained by a number of laboratories. The situation was less favourable for the production cross sections of neutrons with nucleons and nuclei for which there were practically no experimental data above 1.4 GeV. This circumstance was naturally limited by checking a variety of theoretical models which predict the behaviour of nucleon cross sections at high energies. The lack of experimental data for neutron cross sections was related to the fact that existing neutron detectors based on measuring the energy of recoil protons by means of absorbers could be used only over an energy range of less than 1 GeV. As energy increases above 1 GeV, the proton paths become many times larger than the mean nuclear free path in an absorber, and this results in a dramatic decrease of detector efficiency. In order to measure the neutron cross sections for GeV energies, a device based on other principles of energy measurement should be used. For high energy neutron cross section measurements in 1958 we proposed a lead-glass Cherenkov counter several nuclear free paths in thickness, i.e., a hadronic total absorption detector (Fig.3). This detector was used for neutron cross section measurements with protons, neutrons and nuclei over an energy range of 2-10 GeV^{3/}.

The construction of Cherenkov counters having a high energy resolution made it possible to measure for the first time the charge exchange scattering cross section of π^- -mesons on hydrogen over an energy range of 2 GeV^{4/}. A study of the charge exchange reaction $\pi^-p \rightarrow \pi^0n$ (1) is of great interest for checking the principles of field theory and also theoretical models in which particle interactions for high energies are interpreted as an exchange of virtual particles.

Using kinematical features of the reaction (1) at high energies, in 1962 it was pointed to the possibility of measuring the charge exchange cross section using a characteristic maximum in the γ -quanta spectrum arising from the absorption of practically monoenergetic π^0 -mesons in a lead-glass counter. A $18 \times 18 \times 30$ cm³ lead-glass counter was constructed for these measurements^{5/}. A study of the γ -spectrometer on a monoener-

Fig.2. A schematic diagram of the experimental set-up for measuring the Panofsky ratio. The radiator is $50 \times 50 \times 20$ cm³ in size. The radiation length is 2.5 cm.





Fig.3. The Cherenkov lead-glass calorimeter (1957) used to measure the interaction cross sections between 2-10 GeV neutrons and protons, neutrons and nuclei. The Pb glass size is $50 \times 50 \times 50$ cm³. The radiation length is 2.5 cm. The energy resolution for 1 GeV electrons is 14% (HWHM).






Fig.4. A schematic diagram of the Cherenkov lead-glass spectrometer. The radiator dimensions are $18 \times 18 \times 30 \text{ cm}^3$. The radiation length is 2.5 cm. The energy resolution, R for 4 GeV electrons is 5% (HWHM).




Fig.5. The amplitude spectra for 2 and 4 GeV electrons. The energy resolution is shown on the curves.

getic electron beam demonstrated a surprising result: over an energy range of 2-4 GeV the counter resolution was practically constant and very high (about 5%) (Figs. 4 and 5). The obtained data suggested that relatively small ($7.2 \times 7.2 \times 12 \text{ rad. lengths}$) counters could be used for the spectrometry of γ -quanta

with an energy of several GeV at a high energy resolution. These results opened up possibilities of constructing multichannel lead-glass hodoscopes.

A qualitatively new method^{/6/} proposed in 1964 permitted one to measure the effective mass of electromagnetic particles by means of spark chambers and lead-glass counters.

The operation principle of a γ -mass-spectrometer is very simple and based on the possibility of measuring simultaneously the direction and energy of electrons and γ -quanta. A high accuracy in measuring the energy and the possibility of identifying the nature of the electromagnetic particles in lead-glass are important when very rare processes are under study. The development of this method and the construction of a high-efficiency detector, i.e., a two-arm Cherenkov mass-spectrometer^{/7/}, made it possible to carry out a series of complicated experiments to prove the existence of rare electromagnetic decays of ρ^0 and ϕ -mesons into electron-positron pairs and to measure their partial widths^{/8/}.

The partial widths "vector meson-lepton pair" is directly related to the transition constants "photon-vector meson" that are of fundamental importance in the vector dominance model and its numerous applications. This mass-spectrometer allowed one to observe the structure in the differential cross section of the reaction $\pi^- p \rightarrow \eta^- n$ at $-t < 0.1 \text{ (GeV/c)}^2$ ^{/9/}, to establish the




Fig.6. The two-arm lead-glass γ -mass-spectrometer. The radiator is $50 \times 50 \times 30 \text{ cm}^3$ in size. The radiation length is 2.5 cm. The energy resolution for 4 GeV electrons is 5% (HWHM).

existence of η' (958)-meson decays into two γ -quanta and to measure their relative probability^{/10/}.

To search for heavy vector mesons, in 1967 we proposed a multichannel Cherenkov lead-glass mass-spectrometer which can be considered as a second generation of mass spectrometers. The new set-up (Fig.7a,b) consists of 90 lead-glass modules^{/11/}. The use of new optical high-transparency glasses, photomultipliers with high sensitivity, new technical aids and materials made it possible to increase significantly the energy resolution R of lead-glass counters. For example, for electrons with an energy of 4 GeV $R = 2.3\%$ (HWHM).

The set-up includes: 1. beam detectors composed of scintillation counters and proportional chambers; 2. γ -quantum detectors composed of 32 wire spark chambers with magnetostrictive information readout. The total number of wires is 64000. The wire chambers are separated into two identical arms of 16 chambers. The chambers are grouped in four: in order to remove ambiguities in the correlation of sparks between different planes, two of them turn through an angle of 17° . Copper converters 1.2 rad lengths in thickness are placed between the groups of chambers, and their





Fig.7. The hodoscopic Cherenkov γ -mass-spectrometer. The hexagonal module is 35 cm deep. The radiation length is 2.5 cm. The energy resolution for 4 GeV electrons is 2.3% (HWHM).




Fig.8. The effective mass spectrum for 2γ events measured by means of the hodoscopic Cherenkov γ -mass-spectrometer.

thickness is a compromise between detection efficiency and angular resolution. The direction of γ -quanta is measured by a group of 4 chambers positioned directly after the converter in which γ -conversion occurred. The first group of 4 chambers from a target is used for γ -quantum identification.

Four identical scintillation fodoscopes, each consisting of 10 elements 100 cm long and 10 cm wide, oriented in the horizontal and vertical planes, are placed between the wire chambers and Cherenkov γ -spectrometers.

The lead-glass counters for γ -ray energy measurements consist of 90 hexagonal prisms 14 radiation lengths deep, each viewed by a 17 cm phototube. The counters are grouped in 45 in each of the two arms. The set-up operates on-line with a computer. The effective mass distribution of 2γ events measured by the γ -mass-spectrometer is presented in Fig.8.

A series of experiments has been performed using the multi-channel γ -mass-spectrometer. Among them the following experiments should be noted:

1. The measurement of the differential cross sections of the reaction $\pi^-p \rightarrow \eta^0n$ from 3 to 5 GeV in the region of momentum transfers from 0 to $0.3 (\text{GeV}/c)^2$. In the differential cross sections an appreciable minimum was observed in the forward direction (having confirmed the data obtained previously^{/9/}) indicating a significant role of the amplitude with changing helicity^{/13/}.

2. The measurement of the differential cross sections of η -meson production by π^- 's on different nuclei which allowed one to measure the interaction cross section between η -mesons and nucleons for a momentum of $3.3 \text{ GeV}/c$ ^{/14/}.

3. The study of cumulative π^0 -meson production at an angle of 180° in the reaction $\pi^-C \rightarrow \pi^0(180^\circ)X$ for $3.8 \text{ GeV}/c$. Experimentally the value of $\langle Q \rangle$ was found to be 0.16 ± 0.01 ($\langle Q \rangle$ is the average number of nucleons in the cumulative volume). This fact was an independent support of a striking universality of the parameter $\langle Q \rangle$ describing the quark-parton structure functions of nuclei^{/15/}.

Increasing the number of spectrometric channels, using filmless readout chambers instead of optical spark chambers and computers, the possibilities of the method of Cherenkov gamma-mass-spectrometers were considerably extended. However, these advantages could not get rid of a variety of grave difficulties. In fact, in order to detect γ -quanta by means of spark chambers, it is necessary to use converters the total thickness of which is limited by the requirement for high energy resolution. For 10 GeV γ -quanta this thickness is not larger than one radiation length^{16/}. In this case the conversion efficiency of the mass-spectrometer is proportional to $\epsilon_k \sim (0.5)^n$, where n is the number of γ -quanta. Large errors in measuring decay angles of γ -quanta which are due to bremsstrahlung of conversion pairs and multiple scattering are another consequence of using converters. It should be emphasized that spark chambers in similar devices have large longitudinal dimensions and a dead time of the order of milliseconds which has an essential influence on the geometric efficiency and fast operation of the mass-spectrometer.

Due to these disadvantages, the problem of construction of γ -mass-spectrometers without spark chambers is of great interest. This problem is complicated because the coordinate of particles should be measured with an accuracy comparable to that achieved by means of modern chamber technique.

In 1972 we proposed one of the solutions of this problem - a cell structure γ -spectrometer consisting of a large number of independent spectrometer-modules of small transverse dimensions^{17/}.

The first question which should be answered can be formulated as follows: what transverse dimensions must the module of the γ -spectrometer have in order to measure the γ -quantum coordinates with an accuracy of ~ 1 mm? The character of the transverse electromagnetic shower development for electrons with an energy of 1 GeV shows that the radius of the shower is weakly dependent on energy and is equal approximately to $1X_m$, where X_m is a Moliere unit. If the transverse dimensions of the spectrometer module satisfy the condition $D_m/D_s < 1$, where D_m and D_s are respectively the module and electromagnetic shower diameters, the shower energy is distributed in the module group. The analysis of the energy distribution between the modules in the group permits the γ -quantum coordinate to be localized by measuring the shower center of gravity. However, a detailed consideration of the problem indicates that in addition to (1) it is necessary to fulfil the second condition: the energy resolution of the spectrometer and hence of each module must be sufficiently high so that the energy difference in a module may exceed measurement errors due to small shifts of the shower axis. The Monte-Carlo method was used for qualitative estimates. This method

made it possible to determine errors in measuring the γ -quantum coordinates ($\Delta X, \Delta Y$) in the spectrometer versus

- 1) the module size " D_m ";
- 2) the coordinates " X, Y " of the γ -quantum entry point to the spectrometer;
- 3) the γ -quantum incident angle " θ " (angle between the γ -quantum direction and the module axis);
- 4) the energy resolution " $\Delta E/E$ " of the γ -spectrometer.

The Monte-Carlo results have shown that the γ -quantum coordinates and direction can be measured with an accuracy of $\Delta X = \Delta Y = +0.5$ mm and $\Delta\theta = +0.5^\circ$, respectively, for the module 1.2 rad. units in diameter if $\Delta E/E = 2.5\%$.

These ideas forming the basis of γ -mass-spectrometers of the third generation have been realized in a series of experimental developments both in USSR and abroad^{18/}.

REFERENCES

1. Pantujev V.S., Khachaturian M.N., Chuvilo I.V. JINR, P-291, Dubna, 1959; Pantujev V.S., Khachaturian M.N., Chuvilo I.V. Pribory i Tekhnika Experimenta, 1960, 1, p.19.
2. Dunaitsev A.F. et al. Proc. of the Ann.Int.Conf. on High Energy Physics at Rochester, 1960, p.181.
3. Khachaturian M.N., Pantujev V.S. Nucl.Instr. and Meth., 1963, 25, p.173; Ozhdyan L. et al. Journ. of Exp. and Theor.Phys., 1962, 42, p.392; Khachaturian M.N. et al. Journ.of Nucl.Phys., 1965, 1, p.134; Khachaturian M.N., Pantujev V.S. Phys.Lett., 1963, 7, p.80.
4. Azimov M.A. et al. JINR, P-1782, Dubna, 1964; Azimov M.A. et al. JINR, P-2436, Dubna, 1965; Azimov M.A. et al. JINR, 2592, Dubna, 1966.
5. Azimov M.A., Pantujev V.S., Khachaturian M.N. Nucl.Instr. and Meth., 1966, 39, p.325.
6. Хачатурян М.Н. Авторское свид. СССР №182249 от 25.05.66 г. Бюл. ОИПОТЗ, 1966, №11, с.47. Azimov M.A. et al. Pribory i Tekhnika Experimenta, 1965, 4, p.223; Khachaturian M.N. et al. JINR, P-2436, Dubna, 1965; Azimov M.A., Khachaturian M.N. JINR, B7-2070, Dubna, 1964; Azimov M.A., Khachaturian M.N. JINR, 173, Dubna, 1964.
7. Khachaturian M.N. et al. Nucl.Instr. and Meth., 1967, 51, p.309; Astvatsaturov R.G. et al. Pribory i Tekhnika Experimenta, 1972, 5, p.21. Astvatsaturov R.G. et al. JINR, 13-6712, Dubna, 1972.
8. Azimov M.A. et al. JINR, E1-3148, Dubna, 1967; Khachaturian M.N. et al. Phys.Lett., 1967, 7, p.349; Azimov M.A.

- et al. Journ.of Nucl.Phys., 1967, 6, p.515; Astvatsaturov R.G. et al. Phys.Lett., 1968, 27B, p.45. Диплом на открытие №94. от 28.04.71 г. Бюл. ОИПОТЗ, 1971, №15, с.3.
9. Hladky J. et al. Phys.Lett., 1970, 31B, p.475.
 10. Khachaturian M.N. et al. 14 Int.Conf. on High Energy Phys., Vienna, 1968, p.466.
 11. Astvatsaturov R.G. et al. Nucl.Instr. and Meth., 1973, 107, p.105; Astvatsaturov R.G. et al. Nucleonika, 1974, 19, p.575. Khachaturian M.N. et al. Nucl.Instr. and Meth., 1979, 16, p.343.
 12. Khachaturian M.N. et al. Pribory i Tekhnika Experimenta, 1979, 4, p.57.
 13. Astvatsaturov R.G. et al. Journ.of Nucl.Phys., 1978, 27, p.401; Arkhipov V.V. et al. JINR, E1-11596, Dubna, 1978,
 14. Arkhipov V.V. et al. Journ.of Nucl.Phys., 1983, 10, p.986; Arkhipov V.V. et al. Journ. of Nucl.Phys., 1984, 39, p.1206.
 15. Astvatsaturov R.G. et al. Journ.of Nucl.Phys., 1981, 34, p.1504; Arkhipov V.V. et al. Journ.of Nucl.Phys., 1982, 35, p.1206; Arkhipov V.V. et al. Journ.of Nucl.Phys., 1983, 38, p.674.
 16. Azimov M.A. et al. Pribory i Tekhnika Experimenta, 1965, 4, p.223.
 17. Хачатурян М.Н. Авт.свид. СССР №508764 от 30.03.76 г. Бюл. ОИПОТЗ, 1976, №12, с.119. Khachaturian M.N., Melkumov G.L. Nucl.Instr. and Meth., 1976, 135, p.505.
 18. Binon F. et al. Nucl.Instr. and Meth., 1981, 188, p.507; Engelmann R. et al. Nucl.Instr. and Meth., 1983, 216, p.45.

WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?
 You can receive by post the books listed below. Prices - in US \$,
 including the packing and registered postage

	Proceedings of the VII All-Union Conference on Charged Particle Accelerators. Dubna, 1980. 2 volumes.	25.00
	Proceedings of the VIII All-Union Conference on Charged Particle Accelerators. Prqtvino, 1982. 2 volumes.	25.00
D2-81-543	Proceedings of the VI International Conference on the Problems of Quantum Field Theory. Alushta, 1981.	9.50
D1,2-81-728	Proceedings of the VI International Seminar on High Energy Physics Problems. Dubna, 1981.	9.50
D17-81-758	Proceedings of the II International Symposium on Selected Problems in Statistical Mechanics. Dubna, 1981.	15.50
D1,2-82-27	Proceedings of the International Symposium on Polarization Phenomena in High Energy Physics. Dubna, 1981.	9.00
D2-82-568	Proceedings of the Meeting on Investigations in the Field of Relativistic Nuclear Physics. Dubna, 1982	7.50
D3,4-82-704	Proceedings of the IV International School on Neutron Physics. Dubna, 1982	12.00
D11-83-511	Proceedings of the Conference on Systems and Techniques of Analytical Computing and Their Applications in Theoretical Physics. Dubna, 1982.	9.50
D7-83-644	Proceedings of the International School-Seminar on Heavy Ion Physics. Alushta, 1983.	11.30
D2,13-83-689	Proceedings of the Workshop on Radiation Problems and Gravitational Wave Detection. Dubna, 1983.	6.00
D13-84-63	Proceedings of the XI International Symposium on Nuclear Electronics. Bratislava, Czechoslovakia, 1983.	12.00
E1,2-84-160	Proceedings of the 1983 JINR-CERN School of Physics. Tabor, Czechoslovakia, 1983.	6.50
D2-84-366	Proceedings of the VII International Conference on the Problems of Quantum Field Theory. Alushta, 1984.	11.00
D1,2-84-599	Proceedings of the VII International Seminar on High Energy Physics Problems. Dubna, 1984.	12.00

Orders for the above-mentioned books can be sent at the address:
 Publishing Department, JINR
 Head Post Office, P.O.Box 79 101000 Moscow, USSR

Received by Publishing Department
 on January 25, 1985.

COMMUNICATIONS, JINR RAPID COMMUNICATIONS, PREPRINTS, AND PROCEEDINGS OF THE CONFERENCES PUBLISHED BY THE JOINT INSTITUTE FOR NUCLEAR RESEARCH HAVE THE STATUS OF OFFICIAL PUBLICATIONS.

JINR Communication and Preprint references should contain:

- names and initials of authors,
- abbreviated name of the Institute (JINR) and publication index,
- location of publisher (Dubna),
- year of publication
- page number (if necessary).

For example:

1. *Pervushin V.N. et al. JINR, P2-84-649, Dubna, 1984.*

References to concrete articles, included into the Proceedings, should contain

- names and initials of authors,
- title of Proceedings, introduced by word "In;"
- abbreviated name of the Institute (JINR) and publication index,
- location of publisher (Dubna),
- year of publication,
- page number.

For example:

Kolpakov I.F. In: XI Intern. Symposium on Nuclear Electronics, JINR, D13-84-53, Dubna, 1984, p.26.

Savin I.A., Smirnov G.I. In: JINR Rapid Communications, N2-84, Dubna, 1984, p.3.

Хачатурян М.Н.

E1-85-55

30 лет работы с черенковскими γ -спектрометрами из свинцового стекла в Дубне

Работа посвящена истории развития методики черенковских γ -спектрометров из свинцового стекла в Лаборатории высоких энергий Объединенного института ядерных исследований.

Работа выполнена в Лаборатории высоких энергий ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1985

Khachaturian M.N.

E1-85-55

Thirty-Year Work with Lead-Glass Cherenkov γ -Spectrometers at Dubna

This paper is devoted to the history of construction and development of lead-glass Cherenkov γ -spectrometers at the Laboratory of High Energies, JINR, Dubna.

The investigation has been performed at the Laboratory of High Energies, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1985