

ОБЪЕДИНЕННЫЙ  
ИНСТИТУТ  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ  
ДУБНА

1155  
83  
1

E11-82-853

3 75

**A.Kondor**

**METHOD OF CONVERGENT WEIGHTS –  
AN ITERATIVE PROCEDURE  
FOR SOLVING FREDHOLM'S  
INTEGRAL EQUATIONS  
OF THE FIRST KIND**

Submitted to "Nuclear Instruments and Methods"  
(Letters)

**1982**

## 1. UNFOLDING PROBLEMS AND INTEGRAL EQUATIONS

Unfolding problems arise in measuring the density  $u(x)$  of some quantity  $x$  when only the distorted (smeared) distribution of  $x$  can be observed,  $x$  may be one dimensional or higher dimensional and  $x$  may be a random variable or a certain physical quantity which can be described by a density function  $u(x)$ . Since the principles of the problems which are considered here are always the same independently of the dimension of  $x$ , in the following, for convenience,  $x$  is always considered one dimensional and taking values between 0 and 1.

A common unfolding problem is the following. Let  $x$  be a random variable with probability density function  $u(x)$ . In a great number of cases a real measurement of  $x$  does not yield an ideal sample of the  $x$ -values from the density  $u(x)$  but one has distorted values  $y$  with a density function  $v(y)$  where the distortion is caused by some measurement noise  $w$  which is independent of  $x$  and which has the probability density  $p(w)$ . This situation leads to the well-known problem of the solution of the convolution integral equation of the first kind.

Very often, however, the independent noise model is not general enough such that more complicated equations than the convolution one have to be solved. First of all the measurement distributions can depend on  $x$  such that  $p$  becomes the function of two variables and the equation relating the distributions  $u, v$  and  $p$  generalizes to the Fredholm's integral equation of the first kind:

$$v(y) = \int_0^1 p(x, y) u(x) dx, \quad 0 \leq y \leq 1, \quad (1)$$

where  $p(x, y) \geq 0$  for  $0 \leq x, y \leq 1$ . Secondly, it can happen that no  $y$ -value is measured even if an  $x$ -value occurred. This fact may be expressed by the function

$$A(x) = \int_0^1 p(x, y) dy, \quad 0 \leq x \leq 1 \quad (2)$$

for this, in this case, the inequality

$$0 \leq A(x) \leq 1, \quad 0 \leq x \leq 1 \quad (3)$$

holds. The function  $A$  is called the acceptance function of the experimental device represented in our case by  $p(x, y)$ . Suppose

that

$$A(x) < 1 \quad (4)$$

for all the  $x$  in some interval in  $[0,1]$ . It then follows from (1) to (4) that

$$\int_0^1 v(y) dy = \int_0^1 \int_0^1 p(x,y) u(x) dx dy = \int_0^1 A(x) u(x) dx < 1 \quad (5)$$

if it is assumed that the order of integration in the double integral in (5) can be changed. The occurrence of this inequality also causes problems in practice.

Two more problems can be considered. The density  $v$  can only be measured with some error and often the function  $p$  is not exactly known. For instance, in high energy physics, either  $p$  can only be estimated by Monte-Carlo simulation or due to computer cost, for an assumed  $u_0$  the corresponding  $v_0$  can be estimated only. Another well-known problem comes from the fact that the problem of solving (1) for given  $p$  and  $v$  is an ill-posed mathematical problem. That means, for small changes in  $v$  large changes in  $u$  may be caused. Many different proposals to solve equation (1) have been made ranging from series solutions over least square solutions to regularization. Since there is no unique method which solves the problem in general, sometimes different methods may have to be tried. This implies that it is important to have different methods available.

In this letter an iterative method is proposed whose convergence properties have not yet been proved mathematically. However, numerous cases of a limited class of equations (1) with measurement noise on  $v$  and  $p$  have shown that good results may be obtained in practice.

In section 2 the class of equations (1) is given to which the iterative method is applied together with the iteration scheme. In section 3 numerical examples are presented.

## 2. THE METHOD OF CONVERGENT WEIGHTS

In order to construct an iteration for solving Eq.(1) the following assumptions are made to define the class of equations to which the iteration is applied.

- (a) The kernel function  $p(x,y)$  is non-negative and continuous on  $[0,1] \times [0,1]$  except possibly isolated integrable singularities.
- (b) The acceptance function  $A(x)$  is positive on  $[0,1]$

$$0 < A(x) = \int_0^1 p(x,y) dy, \quad 0 \leq x \leq 1. \quad (6)$$

(c)

$$g(y) = \int_0^1 p(x,y) f(x) dx > 0, \quad 0 \leq y \leq 1 \quad (7)$$

for every integrable positive function  $f(x)$ .

- (d) For a given  $v=v(y)$  which is assumed to be positive and continuous on  $[0,1]$  the integral equation

$$v(y) = \int_0^1 p(x,y) u(x) dx, \quad 0 \leq y \leq 1 \quad (8)$$

has a unique continuous solution  $u=u(x)$  for  $0 \leq x \leq 1$ .

(e)

$$\int_0^1 \int_0^1 p(x,y) u(x) dx dy = \int_0^1 \int_0^1 p(x,y) u(x) dy dx. \quad (9)$$

Before constructing the iterative scheme a consequence of the above assumptions will be discussed. Suppose one has a series of iterative approximations  $\{u^{(j)}(x)\}_0^\infty$  of the exact solution  $u(x)$ . Using Eq.(7) one can calculate  $v^{(j)}(y)$  the iterative right-hand side

$$v^{(j)}(y) = \int_0^1 p(x,y) u^{(j)}(x) dx, \quad 0 \leq y \leq 1. \quad (10)$$

Introducing  $\delta^{(j)}(x)$ , the difference of  $u(x)$  and  $u^{(j)}(x)$ , the following equation can be found

$$\left[1 - \frac{1}{r^{(j)}(y)}\right] v(y) = \int_0^1 p(x,y) \delta^{(j)}(x) dx, \quad 0 \leq y \leq 1, \quad (11)$$

where

$$\delta^{(j)}(x) = u(x) - u^{(j)}(x) \quad \text{and} \quad r^{(j)}(y) = \frac{v(y)}{v^{(j)}(y)}$$

Eq.(11) proves that constructing a convergent to the unity series  $\{r^{(j)}\}_0^\infty$  under the assumptions (c) and (d) one will have a convergent series of the corresponding  $\{u^{(j)}\}_0^\infty$  with the limit  $u^\infty = u(x)$  being the solution of Eq.(8).

Forming such an iteration at first we transform the original equation to one having a normalized kernel. It is easy to do via introducing the so-called importance function

$$z(x) = A(x) u(x). \quad (12)$$

It then follows from (1), (2) and (6) that

$$v(y) = \int_0^1 q(x,y) z(x) dx, \quad 0 \leq y \leq 1, \quad (13)$$

where

$$q(x,y) = \frac{p(x,y)}{A(x)}$$

and  $q(x,y)$  is normalized to the unity. Eq.(13) has the property of normalization

$$\int_0^1 v(y) dy = \int_0^1 \int_0^1 q(x,y) z(x) dx dy = \int_0^1 z(x) dx. \quad (14)$$

This normalization may have a regularization effect on the iterative solution, so Eq. (14) is used as a motivation for requesting to satisfy Eq.(14) by all the iterants. This requirement implies the following form of the iterational equation. Choosing a starting function  $u^{(0)}(x)$  one can calculate  $r^{(0)}(y)$  as

$$r^{(0)}(y) = \frac{v(y)}{v^{(0)}(y)}, \quad 0 \leq y \leq 1$$

and  $z^{(0)}(x)$  as

$$z^{(0)}(x) = A(x) u^{(0)}(x), \quad 0 \leq x \leq 1$$

and then the further iterants  $z^{(j)}(x)$  will be given by

$$z^{(j+1)}(x) = \int_0^1 r^{(j)}(y) q(x,y) dy \cdot z^{(j)}(x), \quad 0 \leq x \leq 1, \quad (15)$$

where equations (7) (13) were used and  $u^{(0)}(x)$  was supposed to be arbitrary but satisfying assumptions (a)-(d).

Eq. (15) has two characteristic features. All the  $z^{(j)}(x)$  satisfy the normalization requested for the exact solution  $z(x)$  by Eq.(14) and the definition of  $q(x,y)$  in (13) guarantees the convergence of  $z^{(j)}(x)$  in the cases when  $r^{(j)}(y) \rightarrow 1$ . The reason of choosing the name the Method of Convergent Weights (MCW) for this iteration was the central role of the convergence of the series  $r^{(j)}$  to the unity. The ratio  $r^{(j)}$  is called a weight because it guarantees as a weighting function the satisfying of the normalization (Eq.(14)) until the series of  $z^{(j)}$  has not been converged.

The basic equation of the iteration can be transformed to the following one

$$u^{(j+1)}(x) = \int_0^1 r^{(j)}(y) q(x,y) dy u^{(j)}(x), \quad 0 \leq x \leq 1 \quad (16)$$

which is equivalent to

$$u^{(j+1)}(x) = (1 + \mathcal{F}^{(j)}) u^{(j)}(x), \quad (17)$$

where

$$\mathcal{F}^{(j)} = \int_0^1 \frac{v(y) - v^{(j)}(y)}{v^{(j)}(y)} q(x,y) dy.$$

One can see that the multiplicative operator  $\mathcal{F}^{(j)}$  depends on its arguments via an integral of the relative error of the iterative right hand side

$$\frac{Bu - Bu^{(j)}}{Bu}; \quad Bf = \int_0^1 p(x,y) f(x) dx.$$

Eqs. (16) and (17) are equivalent in the case of exact values but in their numerical realization Eq.(17) has advantages giving higher stability against the rounding errors and those of the discretization of the integrals.

### 3. TEST CASES

Two test cases will be presented to demonstrate the convergence and the stability of the MCW. Both the cases are created on the analogy of real physical experiments.

#### Case 1

In this test case we demonstrate the solution of the one-dimensional smearing problem by the example of a purified numerical model of a real experimental problem<sup>1/</sup>. The problems arising from the finite experimental resolution and the imperfect acceptance of a real experiment we modelize by a Fredholm's integral equation of the first kind:

$$v_t(y) = \int_a^b p_t(x,y) u_t(x) dx, \quad (T1)$$

where

$$a = 4, \quad b = 24$$

$$u_t(x) = 1600/x^2$$

$$p_t(x,y) = [A - B(5-x)^2 - \sum_{k=1}^3 C_k \exp\{-D_k(x-x_k)^2\}] x$$

$$\times \exp\{-E(x-y)^2\} F$$

$$A = 1.2$$

$$B = \begin{cases} 1 & \text{for } x < 5 \\ 0.0023 & \text{for others} \end{cases}$$

$$C_k = 0.4; 0.3; 0.2$$

$$D_k = 5; 5; 1$$

$$x_k = 7; 9; 13$$

$$E = 10$$

$$F = \begin{cases} 1 & \text{for } |x-y| < 0.08x + 0.88 \\ 0 & \text{for others} \end{cases}$$

and the right hand side  $v_i(y)$  is given by straightforward integration (see Fig.1).




Fig.1. Functions  $u_t(x)$  and  $v_t(y)$  for the Case 1.

We have chosen 50 equidistant coarse mesh points for discretizing of functions of one variable. The pseudo-experimental errors of the RHS we had from the following expression:

$$\tilde{v}_i = v_i(x_i)(1 + \theta_i p), \quad (T2)$$

$$i = 1, 2, \dots, 50,$$

where:  $\theta_i$  are equally distributed random numbers from the interval  $[-1, 1]$  and  $p = 0.05$ .

The possible errors of the kernel function we have modeled in the following manner:

$$p_{i,k} = p_t(x_i, y_k)(1 + \theta_{i,k} \tilde{p}), \quad (T3)$$

$$i, k = 1, 2, \dots, 50,$$

where

$$\tilde{p} = \max\{0.5, (1 + p\sqrt{|i-k|})\}.$$




Fig.2. Fluctuations in the iterated solution in the case of +5% error in the RHS and +(5±50)% error in the kernel function.




Fig.3. Changes in the stability of the iterated solution applying smoothing transformations to the kernel function.

The discretization of Eq. (T1) has been provided using the Simpson's quadrature with 20 fine mesh intervals in each coarse one.

The resulting quadratic system of linear equations has been solved by using the MCW and the standard Seidel's method, the later used for comparison.

For the purpose of illustrating the application of the MCW we tested the convergence in the ideal case, i.e., with errorless  $p_t(x,y)$  and  $v_t(y)$  but having the errors of the discretization (Fig.1). We started with a trial function  $u^{(0)} = \text{constant}$ . The relative error of the iterant was found to converge to within +0.5% in 7 iterations. The stability of the iteration has been proved in 1000 iteration steps in which case the iterant remained in the +0.5% error corridor.

Modelizing the experimental errors as formulated above we have got the following results. In the case of  $\tilde{v}$  and  $p_t$  the result from 1 iteration is shown in Fig.2 by full line, while the case of  $\tilde{v}$  and  $\tilde{p}$  is illustrated by triangles. The parallel computation by the Seidel's iteration leads to a solution with fluctuations in a range of +100% for the case of  $\tilde{v}$  and  $\tilde{p}$ .

The stability of the MCW can be improved by two transformations of the basic equations both having smoothing character. At first one can transform the discretized system of linear equa-

tions to the normalized one multiplying by the transposed matrix of the system. As a second step one can apply a proper power function (with an exponent  $0 < \kappa < 1$ ) for substituting  $r^{(j)}(y)$  by  $[r^{(j)}(y)]^\kappa$  in Eq. (16). Slowed down the iteration one has to choose a more realistic trial function than  $u^{(0)} = \text{const.}$  The change in the stability of the MCW has been investigated in 7 iterations choosing

$$u^{(0)}(x) = 1100/x^2(1+2\sin(0.2x-0.8))$$

which differs from  $u_t(x)$  in a range of  $\pm(20 \div 100)\%$ . The smoothing properties of the above transformations for  $\kappa=1/2$  are shown in Fig.3.

### Case 2

For the second test case we have chosen a typical ill-posed problem. Our test case is based on that of the monography of Tikhonov and Arsenin<sup>1/2/</sup>.

It can be constructed the following mathematical experiment on the analogy of the unfolding procedure of irradiation's spectra. Suppose that the physical process under discussion led to solving the following integral equation:

$$v_t(y) = \int_a^b p_t(x,y) u_t(x) dx, \quad (T4)$$

where

$$a=0 \quad b=10$$

$$u_t(x) = (1 - \exp\{-A(b-x)^2\}) + B \sin\left(\frac{\pi \cdot x}{b}\right) + \sum_{k=1}^2 C_k \exp\{-D_k(x-x_k)^2\}$$

with

$$A = 2$$

$$B = 0.5$$

$$C_k = 6; 6$$

$$D_k = 3; 4$$

$$x_k = 3; 7$$

$$p_t(x,y) = (1 - \frac{x}{y}) \gamma(y-x);$$

$$\gamma(z) = \begin{cases} 1 & \text{for } z > 0 \\ 0 & \text{for others} \end{cases}$$

and  $v_t(y)$  is given by straightforward integration (see Fig.4).




Fig.4. Functions  $u_t(x)$  and  $v_t(y)$  for the Case 2.




Fig.5. Comparison of fluctuations excited by  $\pm 5\%$  of the RHS for large numbers of iterations.

The discretization has been provided as in Case 1. The errors of the RHS have been modeled by the formula given in ref.<sup>1/2/</sup>:

$$v_i = v(y_i) (1 + \theta_i \sqrt{\frac{3(b-a)}{b^3-a^3}} \cdot p); \quad i = 1, 2, \dots, 50,$$

where  $\theta_i$  are random numbers from the interval  $[-1, 1]$  and  $p=0.05$ .

Due to the integral in Eq. (16) the convergence of the MCW is extremely slow for this test case which gives good possibilities of investigating the numerical stability of the iteration. For the ideal case ( $p=0$ ) the result of 4000 steps is presented in Fig.5 in comparison with the exact solution. Choosing  $p=0.05$  one can see (Fig.5) that the characteristic fluctuations of  $u^{(k)}$  appear enough late for distinguishing real peaks from ones arising due to errors of the RHS.

### 4. CONCLUSIONS

Introducing the acceptance and the importance functions related to the kernel function of Fredholm's integral equations of the first kind has led to the construction of an iterative procedure which seems to be satisfactorily stable against the errors of the RHS and the kernel function. The convergence is controlled by the integral of the relative error of the iterated RHS that restricts the iteration to be recommended for solving integral equations with nonsmooth RHS.

The effect of the transformation to the normal equation as well as that of the introduction of a power function of the RHS' ratio (in Eq. (16)) is to reduce the instability of the MCW due to the errors in the RHS and the kernel function. Application of an exponent in the power function far from the unity makes the convergence slow down, so in such a case one has to choose a more realistic trial function than  $u^{(0)} = \text{const.}$

It is hoped that improvements in the direction of regularization of the basic equation will permit the method to be applied in a wide range of the ill-posed problems.

#### 5. ACKNOWLEDGEMENTS

The author would like to thank Dr. B.Schorr (CERN) and Dr. G.Emelyanenko (JINR) for helpful discussions.

#### REFERENCES

1. Bollini D. et al. Phys.Rev.Lett., 1981, 104B, p.403.
2. Tikhonov A.N., Arsenin V.Ya. Metody resheniya nekorrektnykh zadach (Methods for Solution of Ill-Posed Problems). "Nauka", M., 1979.

Received by Publishing Department  
on December 14, 1982.

#### WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices - in US \$,  
including the packing and registered postage

D-12965 .	The Proceedings of the International School on the Problems of Charged Particle Accelerators for Young Scientists. Minsk, 1979.	8.00
D11-80-13	The Proceedings of the International Conference on Systems and Techniques of Analytical Computing and Their Applications in Theoretical Physics. Dubna, 1979.	8.00
D4-80-271	The Proceedings of the International Symposium on Few Particle Problems in Nuclear Physics. Dubna, 1979.	8.50
D4-80-385	The Proceedings of the International School on Nuclear Structure. Alushta, 1980.	10.00
	Proceedings of the VII All-Union Conference on Charged Particle Accelerators. Dubna, 1980. 2 volumes.	25.00
D4-80-572	N.N.Kolesnikov et al. "The Energies and Half-Lives for the $\alpha$ - and $\beta$ -Decays of Transfermium Elements"	10.00
D2-81-543	Proceedings of the VI International Conference on the Problems of Quantum Field Theory. Alushta, 1981	9.50
D10,11-81-622	Proceedings of the International Meeting on Problems of Mathematical Simulation in Nuclear Physics Researches. Dubna, 1980	9.00
D1,2-81-728	Proceedings of the VI International Seminar on High Energy Physics Problems. Dubna, 1981.	9.50
D17-81-758	Proceedings of the II International Symposium on Selected Problems in Statistical Mechanics. Dubna, 1981.	15.50
D1,2-82-27	Proceedings of the International Symposium on Polarization Phenomena in High Energy Physics. Dubna, 1981.	9.00
D2-82-568	Proceedings of the Meeting on Investigations in the Field of Relativistic Nuclear Physics. Dubna, 1982	7.50
D9-82-664	Proceedings of the Symposium on the Problems of Collective Methods of Acceleration. Dubna, 1982	9.20
D3,4-82-704	Proceedings of the IV International School on Neutron Physics. Dubna, 1982	12.00

Orders for the above-mentioned books can be sent at the address:  
Publishing Department, JINR  
Head Post Office, P.O.Box 79 101000 Moscow, USSR

## SUBJECT CATEGORIES OF THE JINR PUBLICATIONS

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches
19.	Biophysics

Кондор А. Метод сходящихся весов - E11-82-853  
итерационная процедура для решения интегральных уравнений  
Фредгольма первого рода

Представлен метод решения интегральных уравнений Фредгольма первого рода для случая положительных ядра и свободного члена. Итерационное решение является удовлетворительно стабильным при наличии возмущений статистического характера. Этот факт и сам основной принцип итерации вытекают из включения в алгоритм функции акцептанса, соответствующего ядру интегрального оператора. Основная черта метода - итерация при помощи гладких функций - станет преимущественной по сравнению со стандартными методами для таких интегральных уравнений с гладкими решениями, у которых как ядро, так и свободный член выразительно негладкие, что часто встречается при решении проблем смещения данных в экспериментах высокой энергии.

Работа выполнена в Лаборатории ядерных проблем ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1982

Kondor A. Methods of Convergent Weights - E11-82-853  
an Iterative Procedure for Solving Fredholm's Integral  
Equations of the First Kind

A method is presented for solving Fredholm's integral equations of the first kind for the case of positive kernel function and right-hand side one. The iterative solution seems to be satisfactorily stable against perturbations of statistical nature. This feature and the basic principle of the iteration itself come from including into the algorithm the so-called acceptance function related to the kernel of the integral operator. The main characteristic of the method - to iterate via smooth functions - becomes to be advantageous as compared to standard methods in such cases of integral equations with smooth solutions when both the kernel function and the right-hand side function are strongly nonsmooth, which is often the case at the solution of the smearing problems of the high energy physics' experiments.

The investigation has been performed at the Laboratory of Nuclear Problems, JINR.  
Preprint of the Joint Institute for Nuclear Research. Dubna 1982