



ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

3795/2-80

11/8-80

E15-80-351

V.B.Belyaev, M.Daum*, V.S.Evseev, R.Frosch,*
V.S.Kurbatov, J.McCulloch*,*V.M.Sidorov,
E.Steiner,* J.Wrzecionko

EXPERIMENTAL STUDY OF
 $\bar{\pi}$ DOUBLE CHARGE EXCHANGE
WITH ${}^7\text{Li}$

Submitted to "Nuclear Physics"

* SIN, Swiss Institute for Nuclear Research,
Villigen, Switzerland.

** Present address: Marconi Space and Defence
System, The Groove, Warren Lane, Stanmore,
Middlesex HA7 4LY, England.

1980

1. INTRODUCTION

The double charge exchange reaction of pions, i.e., $\pi^{\pm} + (Z, A) \rightarrow \pi^{\mp} + (Z \pm 2, A)$, is a good tool for studying nuclear states with proton excess or neutron excess. In particular, it may be helpful in searching for new nuclei. The feasibility of such experiments has been demonstrated by many groups (see, e.g., refs. ^{1-10/}). As the cross sections for double charge exchange are small (typically 10^{-30} to 10^{-29} cm² for a given final nuclear state), experiments of this kind are of special interest if they are carried out at high-current accelerators ("meson factories").

We have performed a study of the reaction



in the high intensity pion channel $\pi\text{E}1$ ^{'11'} at SIN. We used stacks of nuclear emulsion plates to detect the π^{+} mesons and measure their energy. The preparation and scanning of the emulsion plates was done at the Laboratory of Nuclear Problems at JINR.

One purpose of our study was the search for the possible particle-stable nucleus ${}^7\text{H}$, i.e., the search for the reaction



The question of the existence of a particle-stable heavy hydrogen isotope ${}^7\text{H}$ arose after the discovery of the isotope ${}^8\text{He}$ ^{'12,13'} which has the same neutron number. As the π^{-} energy and the π^{+} angle are fixed, reaction (2) above would produce a peak in the π^{+} energy spectrum, situated near the high energy end of the continuous π^{+} spectrum due to double charge exchange with prompt disintegration of the residual nucleus. This disintegration spectrum was recorded in our experiment down to low π^{+} energies.

The stopped π^{+} can be detected in the developed emulsion, among the many π^{-} tracks, because of their characteristic decay $\pi^{+} \rightarrow \mu^{+} + \nu_{\mu}$ (whereas the stopped π^{-} are captured in nuclei and produce stars). The energy of the π^{+} and π^{-} is determined by measuring their range in the emulsion. The nuclear emulsion technique is suitable for a first experiment because one can record the whole scattered π^{-} and π^{+} energy spectrum simultaneously with a very simple experimental set-up.




Fig. 1. Experimental arrangement.
 1. Incident π^- beam envelope.
 2. Multiwire proportional chamber used as beam profile monitor.
 3. ${}^7\text{Li}$ target.
 4. Stack of nuclear emulsion plates.
 5. Lead bricks.
 6. Supporting table.

2. DESCRIPTION OF THE EXPERIMENT

Our experimental arrangement is shown

in Fig. 1. Negative pions from the πE1 channel¹¹ pass through a thin multiwire proportional chamber¹⁴ used to monitor the beam profile. The magnets and collimators of the πE1 channel were set for a central π^- momentum of 200 MeV/c and a momentum bite of 2% FWHM. The beam was focused upon a 0.8 g/cm² thick ${}^7\text{Li}$ target oriented as shown in Fig. 1. This target was irradiated for three hours with a total of 5×10^{11} negative pions. A stack of 30 nuclear emulsion plates NIKPHI BR-2 (plate dimensions 110x30x0.6 mm³) was placed at 30° to the beam at a distance of 425 mm from the target center. The emulsion stack was shielded with lead bricks as indicated in Fig. 1.

As mentioned in section 1, the emulsion plates were used to detect secondary pions of both signs. For test purpose we also irradiated further stacks of emulsion plates after replacing the ${}^7\text{Li}$ target by targets of beryllium and polyethylene.

3. DETERMINATION OF THE INCIDENT π^- ENERGY

The π^- elastically scattered from the three targets mentioned above allowed an independent determination of the beam energy and energy spread. The observed π^- range distributions are shown in Fig. 2 as solid lines. For all three targets the count of scattered π^- was concentrated on the domain of ranges expected for elastic scattering. The experimental π^- range




Fig. 2. Distribution of π^- ranges in the nuclear emulsion for the three scattering targets used, in the region of elastic π^- scattering. Solid lines: Observed events. Dashed lines: Gaussian fits; see text (section 3).

Table

Results on elastic π^- scattering; cf. section 3.

Target	\bar{R} (mm)	ΔR (mm)	\bar{E}_π (MeV)	ΔE_π (MeV)	χ^2
${}^7\text{Li}$	103.8	4.4	100.0	2.3	19.7
CH_2	103.4	3.2	99.4	1.2	6.7
${}^9\text{Be}$	103.0	3.5	99.3	1.5	8.8

distributions between 99 and 109 mm were approximated by Gaussian distributions (dashed lines in Fig. 2). In the table the mean projected range \bar{R} , the observed r.m.s. range spread ΔR , the corresponding mean incident π^- energy \bar{E}_π , and the incident pion r.m.s. energy spread ΔE_π are given. In the calculation of \bar{E}_π a correction to the projected range and straggling¹⁵ were taken into account. In the case of ${}^7\text{Li}$ the fit is considerably worse than for the other targets (see the table), possibly because of inelastic π^- scattering




Fig.3. Histogram with error bars: Distribution of π^+ kinetic energies derived from the observed distribution of π^+ ranges in the emulsion plates irradiated from the ${}^7\text{Li}$ target; the interval from 96 to 103 MeV was not scanned for π^+ because of large π^- rate; see text. The horizontal line at an ordinate of 0.95 represents an estimate of the π^+ background not

related to the ${}^7\text{Li}$ target. Vertical arrows indicate binding energies of hypothetical ${}^7\text{H}$ ground state; cf. reaction (2). Curves 1, 2 and 3 are phase space predictions based on reactions (3), (4) and (5), respectively.

leading to the first excited state of ${}^7\text{Li}$ (excitation energy 0.48 MeV). In the case of polyethylene the contribution from elastic π^- scattering by hydrogen was neglected since the corresponding cross section (≈ 13 mb) is much smaller than the cross section for elastic π^- scattering by carbon nuclei (≈ 280 mb)^{16/}. The resulting incident pion energy, averaged over the three targets, is $E_\pi = 99.6$ MeV. The experimental uncertainty of this E_π value is estimated to be about 2 MeV.

The πE1 channel magnet and collimator settings mentioned in section 2 lead to a mean beam energy $E_\pi = 104.3$ MeV and an r.m.s. energy spread $\Delta E_\pi = 1.4$ MeV. In this case, the estimated uncertainty of E_π is 1 MeV. The value $\Delta E_\pi = 1.4$ MeV agrees well with lines 2 and 3 of the table; in the case of the first line (${}^7\text{Li}$ target) the emulsion value may be too large because of inelastic π^- scattering, as discussed above.

Our final estimates of the mean π^- energy and r.m.s. energy spread upstream of the ${}^7\text{Li}$ target are (102 ± 2) MeV and (1.4 ± 0.2) MeV, respectively.

4. RESULTS FOR DOUBLE CHARGE EXCHANGE

In scanning the photoemulsion plates irradiated from the ${}^7\text{Li}$ target we found 257 π^+ mesons with ranges from 9 to 119 mm. The interval from 99 to 109 mm was not scanned for π^+ because of the presence of a large background due to stopping π^- mesons elastically scattered by lithium nuclei. In fig.3 the doubly differential distribution $d^2N / (dE_\pi / d\Omega)$ for the observed π^+ mesons is shown. Curve 1 in fig.3 corresponds to the π^+ energy distribution expected for the reaction



with all neutrons in the final state unbound. In the calculation a constant matrix element was assumed, i.e., the usual phase space distribution was calculated for a final state consisting of eight particles. Curve 2 in fig.3 is the expected π^+ energy distribution for the reaction



where the four final neutrons are, as before, assumed to be unbound (the final state consists of six particles). Curve 3 in fig.3 is for the hypothetical reaction



where now the four neutrons in the final state are assumed to be bound with zero binding energy. All curves in fig.3 are normalized to the experimental spectrum in the energy interval from 20 to 65 MeV. This normalization was performed after subtraction of a flat (energy-independent) background from the experimental data. This flat background was determined from the events with $E_\pi > 65$ MeV; it is attributed to a π^+ background in the experimental area not related to the ${}^7\text{Li}$ target. The total background is estimated as 110 events, so we are left with about 150 double charge exchange events. Vertical arrows in Fig.3 correspond to the π^+ energies at which peaks would be expected in the spectrum if bound states of one proton and six neutrons with binding energy ΔE existed; negative values of ΔE correspond to the production of a "virtual" (particle-unstable) excited nuclear state. In the experimental spectrum of fig.3 there are no significant peaks for $E_\pi > 65$ MeV. If a peak due to ${}^7\text{H}$ production is assumed to be of a Gaussian shape with an r.m.s. width of 2 MeV (π^+ range straggling and the energy distribution of the incident π^- beam are being taken into account), then it is possible to determine,




Fig. 4. Basic diagram for double charge exchange of pions discussed in ref. (17) and used for fig. 5.

from the data shown in fig. 3, the upper limit of the differential cross section ($d\sigma/d\Omega$) for the production of ($p + 6n$) in bound states with binding energy from 0 to 25 MeV or in narrow excited states with negative "binding energy" from - 5 to 0 MeV. Our result is

$$(d\sigma/d\Omega) < 1.0 \times 10^{-31} \text{ cm}^2/\text{ster} \text{ (90\% C.L.)} \quad (6)$$


This upper limit may be compared to the cross sections for other double charge exchange reactions leading to nuclear ground states, e.g.,



which has been studied in refs. ^{7,8,10}. The authors ⁸ find a cross section of $(3 \pm 1) \times 10^{-31} \text{ cm}^2/\text{ster}$ at a π^+ energy of 148 MeV and $(2.1 \pm 0.8) \times 10^{-31} \text{ cm}^2/\text{ster}$ at 187 MeV; the scattering angle was 18° in both cases. At 0° the cross section is much larger ⁷; in a recent more complete study of reaction (7) a diffraction minimum of the differential cross section has been observed ¹⁰. It is not impossible that for our reaction (2) at $E_{\pi^-} = 102 \text{ MeV}$ there is a similar diffraction minimum near our angle of 30°. We therefore conclude from our data that the existence of a narrow ⁷H state with a binding energy between - 5 MeV and + 25 MeV is rather unlikely but not safely excluded. Our upper limit (eq. (6)) is four times smaller than that of Gilly et al. ⁴ for the same reaction (2), at 0°, obtained at a fixed secondary π^+ energy of 180 MeV and a variable incident π^- energy ($180 \text{ MeV} \leq E_{\pi^-} \leq 260 \text{ MeV}$).

In the breakup region ($E_{\pi} < 65 \text{ MeV}$) the experimental energy spectrum of fig. 3 cannot be explained on the basis of processes with six to eight known non-interacting particles in the final state (curves 1 and 2 in fig. 3). The hypothetical reaction (5), represented by curve 3, gives a fairly good fit to the data, and the same is true for reactions similar to (5) but with five or six neutrons, bound with zero binding energy, plus a deuteron or a proton, in the final state.

Fig. 5. Same as fig. 3; theoretical predictions based on ref. (17). Curves 1, 2, 3; see caption to fig. 3.



We could not determine the π^+ energy spectrum for $E_{\pi} < < 20 \text{ MeV}$, because the corresponding (upstream) part of the emulsion was too strongly blackened by charged particles. Assuming that the π^+ spectrum in that region ($E_{\pi} < < 20 \text{ MeV}$) is described by the phase space curve for reaction (5) we obtain

$$(d\sigma/d\Omega)_{\text{tot}} = (4.2 \pm 1.7) \times 10^{-30} \text{ cm}^2/\text{ster} \quad (8)$$

for the differential cross section, integrated over π^+ energies, of reaction (1). The uncertainty of this result (one standard deviation) is mainly due to the estimated uncertainty of the extrapolation to $E_{\pi} = 0$ discussed above, to the uncertainty of the total number of incident π^- mesons and to the statistical error of the number of counted π^+ mesons.

5. DISCUSSION AND OUTLOOK

Our experimental data imply an important contribution from the interaction of several nucleons and possibly the π^+ in the final state of reaction (1). This conclusion is not altered by the considerations of ref. ¹⁷ where it is suggested that double charge exchange is described by the diagram of fig. 4. Calculations that we have performed with this diagram for reactions (3), (4) and (5) have given π^+ spectra which practically coincide with those calculated over phase space. The results are shown in fig. 5; the curves are seen to be very similar to the result of the corresponding phase space calculations (fig. 3).

Curve 3 of fig. 3 and fig. 5 corresponds to reaction (5). As mentioned in the previous section, other reactions give a similarly good fit, if there are three particles or resonances in the final state. The corresponding general diagram is shown in fig. 6. Here X and Y are nucleon clusters. In order to test,




Fig.6. Diagram for double charge exchange with two nucleon clusters X and Y.

e.g., reaction (5), an experiment in which coincidences of π^+ and ${}^3\text{H}$ are recorded, ought

to be carried out. Besides, the systems $\pi^+ + p$ or $\pi^+ + {}^3\text{H}$ could be detected and used for missing mass spectra. The presence of lines in such spectra would indicate the existence of a group of correlated neutrons.

An alternative explanation of our experimental π^+ spectrum is based on the hypothesis that a three-particle resonance πNN with isospin component $T_z = 0$ is formed with large probability in nuclei. Then the process occurs according to the diagram of fig.7. During the lifetime of the resonance B the remaining nucleons scatter and the π^+ spectrum again corresponds to a three-particle final state ($\pi^+ \text{nn}$). References ¹⁸ and ¹⁹ contain indications of a strong attraction and even of a resonance arising in the system $\text{N} + \Lambda_{33}$ in the state with $J^\pi = 2^+$ and $T = 2$. If this resonance contributes to the double charge exchange on ${}^7\text{Li}$, then one may expect that the process on ${}^8\text{Li}$ will produce the same spectrum of π^+ mesons as on ${}^7\text{Li}$. Also, this πNN resonance model can be tested by investigating the dependence of the π^+ spectrum on the incident π^- energy.

Finally it should be noted that our data (fig.3) contain possible evidence for a broad excited state of ${}^7\text{H}$ at an excitation energy of about 25 MeV ($\Delta E = -25$ MeV). A further study of this energy region with improved statistics would help to clarify this.

ACKNOWLEDGEMENTS

We thank Dr. G.H.Eaton, P.Gheno, M.Moser and M.Plotrowski for their help with the data taking, the SIN accelerator crew for the provision of stable beam and the scanning group at JINR for their excellent work. We would also like to thank Professor J.P.Blaser for his support of our experiment and, concerning two of us (V.S.E. and V.M.S.), for the kind hospitality at SIN.




Fig.7. Diagram with a resonance B of two nucleons and a pion.

REFERENCES

1. Batusov Yu.A. et al. JETP, 1964, 46, p.817.
2. Batusov Yu.A. et al. JINR, P1-3248, Dubna, 1967.
3. Sperinde J. et al. Phys.Lett., 1970, 32B, p.185.
4. Gilly L. et al. Phys.Lett., 1965, 19, p.335.
5. Boynton P.E. et al. Phys.Rev., 1968, 174, p.1083.
6. Chultem D. et al. Nucl.Phys., 1979, A316, p.290.
7. Marks T. et al. Phys.Rev.Lett., 1977, 38, p.149.
8. Corfu R. et al. Phys.Lett., 1977, 69B, p.301.
9. Seth K.K. et al. Phys.Rev.Lett., 1978, 41, p.1589.
10. Seth K.K. et al. Phys.Rev.Lett., 1979, 43, p.1574.
11. Frosch R., Funk P. SIN report TM-30-15, 1974.
12. Batusov Yu.A. et al. Phys.Lett., 1966, 22, p.487.
13. Poskanzer A.M. et al. Phys.Rev.Lett., 1966, 17, p.1271.
14. Eaton G.H., Pozar F. SIN report TM-33-02, 1975.
15. Powell C.F., Fowler P.H., Perkins D.H. The Study of Elementary Particles by the Photographic Method, Pergamon Press, 1959.
16. Barashenkov V.S. et al. JINR, P2-4520, Dubna, 1969.
17. Germond J.F., Wilkin C. Lett.Nuovo Cim., 1975, 13, p.605.
18. Arenhövel H. Nucl.Phys., 1975, A247, p.475.
19. Belyaev V.B., Moller K., Simonov Yu.A. Journ.of Phys.G, Nucl.Phys., 1979, 5, No.8, p.1057.

Received by Publishing Department
on May 15 1980.