

Объединенный
институт
ядерных
исследований
Дубна

E17-84-659

E.I.Kornilov, V.B.Priezzhev

**PSEUDO-CRITICAL BEHAVIOUR
OF SMOOTHNESS
OF SELF-AVOIDING LOOPS**

Submitted to "Journal of Physics A"

1984

In a recent paper^{/1/} the statistical behaviour of self-avoiding loops on d-dimensional hypercubic lattices has been numerically studied. The partition function of the investigated model is defined by Rys and Helfrich

$$Z(t) = \sum_{n,s} g(n,s) m^s t^n, \quad (1)$$

where t is the monomer fugacity and $g(n,s)$ denotes the number of configurations of self-avoiding loops with total length n in units of the lattice spacing, containing s nonintersecting contours with multiplicity m . Different values of m describe different physical systems. So, the limit $m \rightarrow 0$ corresponds to the single polymer ring without self-intersection.


The aim of the present letter is to consider the case $m=1$ and to discuss an exactly solved model related to the self-avoiding ring polymers model in the simple square lattice. In this case we can rewrite the partition function (1) in the form:

$$Z(t) = \sum_n G(n) t^n, \quad (2)$$

where $G(n)$ is the number of all configurations of self-avoiding loops with total length n .

Consider a square $M \times N$ lattice with periodic boundary conditions. The problem of calculating the partition function (2) is equivalent to solving the 8-vertex model with the following parameters

$$\omega_1 = 0, \quad \omega_2 = 1, \quad \omega_i = t, \quad i = 3, \dots, 8, \quad (3)$$



where all types of configurations labelled by the index i are shown in fig.1. Unfortunately, the set of parameters (3) belongs neither to the Baxter condition^{/3/}

Fig.1. Typical loop configuration. Site of type 1 is forbidden.

$$\omega_1 = \omega_2, \quad \omega_3 = \omega_4, \quad \omega_5 = \omega_6, \quad \omega_7 = \omega_8 \quad (4)$$

nor to the free-fermion condition^{/4/}

$$\omega_1 \omega_2 + \omega_3 \omega_4 = \omega_5 \omega_6 + \omega_7 \omega_8 \quad (5)$$

under which the 8-vertex model has an exact solution.

Several years ago an auxiliary model has been proposed^{/5/} which is closely related to the self-avoiding loops problem. The idea is to put $\omega_3 = \omega_4 = \sqrt{2} t \sin \alpha$ (3) and to consider the sites of types 3 and 4 as defects on a polymer chain. Every two defects double the number of the loops configurations, so, the partition function of the auxiliary model can be represented in the form:

$$Z^*(t) = \sum_n G^*(n) t^n. \quad (6)$$

Here $G^*(n)$ is the number of arrangements of a set of closed chains with total length n in which the above property of defects is fulfilled. The partition function (6) can be easily calculated by the method of Pfaffians. In the large N and M limit the result is^{/5/}

$$Z^*(t) = \exp \left\{ \frac{MN}{8\pi^2} \int_0^{2\pi} \int_0^{2\pi} \ln [1 + 4t^2 + 2^{3/2} t \cos \alpha + 2^{3/2} t (1 + 2^{1/2} t \cos \alpha) \cos \beta] d\alpha d\beta \right\}. \quad (7)$$

The model has an Ising-like singularity at $t_c = 2^{-3/2}$, e.g., the specific heat tends to infinity by the law $c \sim \ln |t - t_c|$.


In this letter we shall use the properties of correlation functions of the auxiliary model to relate the partition functions (2) and (6) and to derive the critical behaviour of the original model.

Consider the correlation function corresponding to the defect density $n_D(t)$ defined as the probability of a site configuration weighted by ω_3 or ω_4 . By the standard procedure described by Montroll^{/6/} the correlation function is given by

$$n_D = 2|X + Q|^{1/2}, \quad (8)$$

where X is the 4×4 matrix

$$X = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \quad (9)$$



and the components of 4x4 matrix Q are

$$Q_{ii} = 0, \quad i = 1, 2, 3, 4$$

$$Q_{12} = Q_{24} = \frac{1}{(2\pi)^2} \int_0^{2\pi} \int_0^{2\pi} d\alpha d\beta \{1 + 2t^2 + 2^{3/2} t \cos \alpha + 2^{1/2} t(1 + 2^{1/2} t \cos \alpha) \cos \beta\} / D(t), \quad (10)$$

$$Q_{13} = Q_{24} = 0$$

$$Q_{14} = -\frac{1}{2\pi^2} \int_0^{2\pi} \int_0^{2\pi} d\alpha d\beta (1 + 2^{1/2} t \cos \alpha + 2^{1/2} t \cos \beta) / D(t),$$

$$Q_{23} = \frac{-2^{1/2} t}{(2\pi)^2} \int_0^{2\pi} \int_0^{2\pi} d\alpha d\beta \{(1 + 2t^2 + 2^{3/2} t \cos \alpha) \cos \beta + 2^{3/2} t(1 + 2^{1/2} t \cos \alpha)\} / D(t),$$


$$Q_{ij} = -Q_{ji},$$

where $D(t) = 1 + 4t^2 + 2\sqrt{2}t \cos \alpha + 2\sqrt{2}t(1 + \sqrt{2}t \cos \alpha) \cos \beta$.

The plot of the ratio of the density of defects $n_D(t)$ to the polymer density $\rho = t(\partial/\partial t) \ln Z^*(t)$:

$$s(t) = \frac{n_D(t)}{\rho(t)} \quad (11)$$

is shown in fig.2. It is natural to call the function $s(t)$ "smoothness" of polymer rings since the sites weighted by ω_3 and ω_4 correspond to rectilinear segments of the polymer. One can see



that the smoothness function behaves like an order parameter. The function $s(t)$ is almost constant above the critical point t_c and rapidly decreases below t_c . The plateau of the smoothness is not just constant, but weakly changes and reaches its limit value $s_\ell = 1/2$ when $t \rightarrow \infty$.

Fig.2. Smoothness s and reduced smoothness \bar{s} against fugacity.

Up to now we were dealing with the exactly solved auxiliary model. To consider the problem of calculation of the partition function (2), we should make some model assumptions. First suppose that the smoothness behaves as a true order parameter, i.e., $s(t)$ is constant above t_c . Second, assume s_ℓ to be the value of this constant. Then for large n we have the following equality

$$G(n) = G^*(n) 2^{-n s_\ell / 2} \quad (12)$$

since the presence of each two defects doubles the number of possible configurations in the original problem, and the number of defects is ns_ℓ .

From (12) a relation follows between the partition functions $Z(t)$ and $Z^*(t)$ above t_c :

$$Z(t) = \sum_n G(n) t^n = \sum_n G^*(n) 2^{-n s_\ell / 2} t^n = \sum_n G^*(n) \tilde{t}^n = Z^*(\tilde{t}), \quad (13)$$

where the new variable \tilde{t} is introduced:

$$\tilde{t} = t/2^{s_\ell / 2}, \quad t > t_c. \quad (14)$$

The singularity of $Z^*(t)$ is known, and therefore we can find the critical point of the model of self-avoiding loops writing:

$$t_c = \tilde{t}_c \cdot 2^{s_\ell / 2} = 2^{-5/4} = 0,4204... \quad (15)$$

This result can be compared with the numerical data of Karowski et al.^{1/} These authors have observed infinite fluctuations in the system of loops at temperature $\beta_0 = 0.86$. Taking into account their definition $t = e^{-\beta}$ we get from (15) $\beta_0 = 5/4 \ln 2 = 0.866...$ in a good agreement with numerical data. Relation (13) together with (14) means that the thermodynamic functions in the self-avoiding loops model have the same singularity as in the auxiliary model above t_c , i.e., the specific heat $c \sim \ln(t - t_c)$, which is also consistent with the results of the numerical analysis.

It should be stressed that we cannot draw any conclusions about the type of singularity below t_c since $s(t)$ is no longer constant at $t < t_c$.

To clarify the origin of the critical behaviour of the smoothness function, we have considered the contribution of the smallest loops (the elementary square in fig.1) to the value of $s(t)$. By definition the proper smoothness of such loops is zero. The density of squares ρ_{sq} can be calculated by the method of Pfaffians. The result of these calculations is presented in fig.2, where \bar{s} is given by

$$\bar{s} = \frac{n_D}{\rho - 4\rho_{sq}} \quad (16)$$

One can see that deleting of the elementary squares makes the smoothness function more even. We conclude that the pseudo-critical behaviour of $s(t)$ is due to the presence of small loops and the smoothness of the long polymer chains depends weakly on the fugacity.

REFERENCES

1. Karowski M. et al. J.Phys.A: Math.Gen., 1983, 16, p. 4073.
2. Rys F., Heilfrich W. J.Phys.A: Math.Gen., 1982, 15, p. 599.
3. Baxter R.I. Ann.Phys., 1972, 70, p. 193.
4. Fan Ch., Wu F.Y., Phys.Rev., 1970, B2, p. 723.
5. Priezzhev V.B. Zh.Eksp.Theor.Fiz., 1978, 74, p. 1177.
6. Montroll E.W. In: Applied Combinatorial Mathematics ed. E.E.Beckenbach (New York, London, Sydney: Jonh Wiley and Sons, Inc.), 1964.

Корнилов Е.И., Приезhev В.Б.

E17-84-659

Псевдокритическое поведение гладкости самоизбегающих полимеров

Обсуждается модель самоизбегающих полимеров на квадратной решетке. Вводится вспомогательная модель, одна из корреляционных функций которой /названная гладкостью/ обладает псевдо-критическим поведением. Этот факт позволяет вычислить критическую точку t_c основной модели и определить тип особенности выше t_c .

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1984

Kornilov E.I., Priezzhev V.B.

E17-84-659

Pseudo-Critical Behaviour of Smoothness of Self-Avoiding Loops

It is shown that the critical point and the logarithmic singularity of the specific heat for the self-avoiding-loop model on the square lattice can be found by introducing an exactly solvable auxiliary model. Evaluations of correlation functions of the latter model show the pseudo-critical behaviour of the density of rectilinear polymer segments.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1984

Received by Publishing Department
on October 29, 1984.