

СООБЩЕНИЯ
ОБЪЕДИНЕННОГО
ИНСТИТУТА
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

C. 54

E17-88-488 e

W.Chmielowski, K.Knast¹, S.Kielich¹

**HYDRODYNAMICAL FIELD FLUCTUATIONS
IN NON-EQUILIBRIUM QUASI-STATIONARY
STATE DUE TO TEMPERATURE GRADIENT
I. GENERAL THEORY²**

¹Nonlinear Optics Division, Institute of Physics,
A.Mickiewicz University, Grunwaldzka 6, 60-780, Poznan, Poland

²This work was carried out under Research Project CPBP 01.12


1988

1. Introduction

One of the essential problems of the theory of irreversible processes concerns the influence of various perturbations on the thermodynamical equilibrium of systems. Two kinds of perturbations — mechanical and thermal — are generally distinguished in the statistical thermodynamics of irreversible processes¹⁾. According to Kubo's terminology^{2,3)}, mechanical perturbations originate in the influence of external (e.g., electric) fields, an action that can be accounted for by including a term describing the interaction of the external field and the system into the Hamiltonian.

However, there exists a vast class of "thermal" perturbations, related with spatial inhomogeneities in the system. Their influence on the system is not directly accessible to an interpretation within the framework of Kubo's theory. The spatial inhomogeneities lead to the emergence of fluxes of matter, momentum and energy, tending to make the system homogeneous. Transport coefficients, such as the viscosity, diffusion and heat conductivity coefficients, are the result of perturbations of this kind.

On the microscopic level, processes involving mechanical perturbations differ essentially from processes due to thermal perturbations. On the macroscopic level, this distinction becomes inessential since all the transport coefficients irrespective of the type of perturbation can be expressed as integrals of the correlation functions of the microscopic fluxes (by the Green-Kubo formulae). Within the same framework, the fluctuational-



-dissipative theorem strictly related with the Onsager hypothesis⁴⁾ concerning the shape of fluctuation decay has successfully been extended to thermal perturbations. The hypothesis states that the behaviour of the system (or part thereof) at non-equilibrium (albeit sufficiently close to equilibrium) is the same irrespective of whether its momentary state is due to the action of external forces or to a spontaneous fluctuation. This is of essential significance when it comes to establishing equations describing the evolution of systems with local inhomogeneities due to fluctuations. On the macroscopic level, non-equilibrium system consisting of great numbers of identical atoms or molecules are described in terms of equations of motion of several macroscopic quantities. The form of these equations depends on what kind of system we are dealing with and on the conditions of its existence. It is at this point that Bogolubov's hierarchisation concept of relaxation times in non-equilibrium statistical thermodynamics becomes essential⁵⁾. According to Bogolubov, a non-equilibrium system in its initial stage requires, in general, a great number of many-body distribution functions for its description. After a short time τ_c (of the order of intermolecular collisions), it attains a kinetical stage, characterized by a single-body distribution function. At this stage of description, kinetic equations related with the processes occurring during the time τ_r (of the order of the time of free motion of a molecule between successive collisions) are established. Whereas the inhomogeneities arising due to fluctuations concern a system that is in the hydrodynamical stage (governed by the relaxation time τ_h). In simple fluids, within the Bogolubov framework,

the relaxation times fulfill the inequality $\tau_c \ll \tau_r \ll \tau_h$. Moreover, no processes of energy transfer between the translational and internal degrees of freedom are assumed to take place. In this way we neglect relaxation effects dependent on the microscopic structure of molecular fluids.

The system is moreover assumed to be sufficiently remote from its critical point for effects typical for near-criticality to be absent. The fundamentals of the theory of effects occurring in fluids near the critical point have been proposed by Smoluchowski⁶⁾. In recent years their theory has been considerably developed by Kocinski⁷⁾. With the above assumptions on the system, the hydrodynamical stage is characterized by the so-called normal distribution function of the number density of molecules, the momentum density, and the energy density. The mentioned above quantities are functions of the time and positions, and determine respectively the following hydrodynamical fields: the scalar fields of the number density of molecules and of the energy (temperature field), as well as the vector field of momentum density.

In simple liquids the fluctuations of the hydrodynamical fields are typically of the order of 10^3 \AA in size and thus greatly in excess of the intermolecular distances. They give rise to collective motions of great numbers of molecules setting free dissipative fluxes connected with viscosity and Joule-Lenz heat. The regions of inhomogeneity in the medium (determined by the size of the fluctuations) are very small compared with the size of the system as a whole, but are sufficiently large to admit a description in terms of the laws of macroscopic physics: hydrodynamics and thermodynamics.

With regard to their size in space and the time-scale of their existence, the hydrodynamical field fluctuations are connected with a variety of phenomena occurring in the medium. Thus, studies of these fluctuations are a source of highly essential data concerning the system and especially the transport processes in the latter. Especial attention is given to fluctuations in number density⁸⁾ because the spectral density of their correlation function is proportional to the structural factor directly related with quantities that are measurable (light and neutron scattering). There are many ways of calculating the fluctuation correlation functions of the hydrodynamical fields leading to results of a form permitting their experimental verification. These methods are based on equations of motion, from simple phenomenological equations up to the equations of generalized statistical thermodynamics comprising effects of "memory" as well as renormalization of the transport coefficients. Reviews of the theoretical methods and respective experiments are due, e.g., to Berne and Pecora⁹⁾, Crosignani, Di Porto and Bertolotti¹⁰⁾, Lallemand¹¹⁾ and Kielich¹²⁾.

Some years ago, a number of papers have appeared, dealing with the spectrum of hydrodynamical fluctuations in fluids at non-equilibrium stationary state. The results are discussed within the framework of light scattering theory. The state of non-equilibrium is obtained by imposing a temperature gradient, or a flow velocity gradient. In most cases, the authors apply methods of fluctuation hydrodynamics based on determining the equations of motion for the mean value of the respective variable¹³⁻¹⁵⁾, whereas others use the method involving an intermediate

linear reaction^{16,17)} or that of kinetic equation theory¹⁸⁾. All these publications fall in two groups: in the one, the equations of motion are established first whereas the fluctuation correlation functions are calculated next, as well as the experimental consequences. In the other, a precise form of the equations of motion is assumed at the very start, and the procedure is restricted to the last two steps; this is justified in as much as in either case relationships derivable from the phenomenological hydrodynamical equations are used for calculating the correlation function. This, in fact, is the Landau-Lifshits method, involving linearization of the hydrodynamical equations with respect to the steady state and supplementing the relationships thus obtained with stochastic terms, containing the stress tensor and heat flow vector. Finally, this leads to equations of the Langevin type.

It is our present aim to determine, in the most general manner, the fluctuation correlation functions for hydrodynamical fields in simple fluids. In Section 2 we introduce the generalized Langevin equation of Grabert¹⁹⁾ using the method of projection operators. Next, on making certain assumptions, we obtain a set of equations of motion for the fluctuations of the hydrodynamical fields in atomic fluids. We use a notation (underlining certain expressions) which enables us to distinguish the contributions from the individual mechanisms. In Section 3 we go over to variables \vec{k}, ω in our formulae and calculate the respective fluctuation correlation functions of the hydrodynamical fields. Section 3, moreover, contains a description of the path followed by us when deriving the stochastic terms of the correlation functions for the equations of motion. Since the

functional coefficients obtained are rather bulky in form. Appendix A gives an appropriate tabulation aimed at abridging their rotation. The results obtained with accuracy to terms in q^2 when deriving the frequency dispersion equation are written out explicitly in Appendix B.

2. Stochastic Functions Describing the Hydrodynamical Field Fluctuations

The possibility of applying Langevin-like equations to the treatment of non-equilibrium fluctuations (non-equilibrium with respect to local equilibrium) was first considered by Keizer²⁰). Subsequently, their applicability has been considered from different standpoints by Hinton²¹), Onuki²²), and Ueyama²³). Grabert¹⁹), on the other hand, starts from Liouville's equation and applies the operator projection method to derive a generalized Langevin equation for fluctuations of macroscopic quantities on the level of generalized statistical thermodynamics. In the case of a simple atomic fluid of interest to us the equations take the following form:

$$\frac{\partial}{\partial t} = -\frac{1}{m} \sum_{\nu} \frac{\partial}{\partial x^{\nu}} \delta g^{\nu} \quad (1)$$

$$\begin{aligned} \frac{\partial}{\partial t} \delta g^{\nu} = & - \sum_{\mu} \frac{\partial}{\partial x^{\mu}} \left[v^{\nu} \delta g^{\mu} + v^{\mu} \delta g^{\nu} - m v^{\nu} v^{\mu} \delta n \right] - \frac{\partial}{\partial x^{\nu}} \delta p + \\ & + \frac{\partial}{\partial x^{\nu}} \left[\sum_{\mu} \frac{\partial v^{\mu}}{\partial x^{\mu}} \left(\delta \zeta - \frac{2}{3} \delta \eta \right) + \left(\zeta - \frac{2}{3} \eta \right) \frac{\partial}{\partial x^{\mu}} \delta v^{\mu} \right] + \\ & + \sum_{\mu} \frac{\partial}{\partial x^{\mu}} \left[\left(\frac{\partial v^{\nu}}{\partial x^{\mu}} + \frac{\partial v^{\mu}}{\partial x^{\nu}} \right) \delta \eta + \eta \left(\frac{\partial}{\partial x^{\mu}} \delta v^{\nu} + \frac{\partial}{\partial x^{\nu}} \delta v^{\mu} \right) - \sum_{\mu} \frac{\partial}{\partial x^{\mu}} \xi^{\nu\mu} \right], \quad (2) \end{aligned}$$

$$\begin{aligned} \frac{\partial}{\partial t} \delta e = & - \sum_{\nu} \frac{\partial}{\partial x^{\nu}} \left[v^{\nu} (\delta e + \delta p) + (\langle e \rangle + p) \delta v^{\nu} \right] + \sum_{\nu} \frac{\partial}{\partial x^{\nu}} \left[\frac{\partial T}{\partial x^{\nu}} \delta x + \chi \frac{\partial}{\partial x^{\nu}} \delta T \right] + \\ & + \sum_{\nu, \mu} \frac{\partial}{\partial x^{\nu}} \left[\left(\frac{\partial v^{\nu}}{\partial x^{\mu}} + \frac{\partial v^{\mu}}{\partial x^{\nu}} \right) (v^{\mu} \delta \eta + \eta \delta v^{\mu}) + \eta v^{\mu} \left(\frac{\partial}{\partial x^{\mu}} \delta v^{\nu} + \frac{\partial}{\partial x^{\nu}} \delta v^{\mu} \right) \right] + \quad (3) \\ & + \sum_{\nu, \mu} \frac{\partial}{\partial x^{\nu}} \left[v^{\nu} \frac{\partial v^{\mu}}{\partial x^{\mu}} \left(\delta \zeta - \frac{2}{3} \delta \eta \right) + \left(\zeta - \frac{2}{3} \eta \right) v^{\nu} \frac{\partial}{\partial x^{\mu}} \delta v^{\mu} + \left(\zeta - \frac{2}{3} \eta \right) \frac{\partial v^{\mu}}{\partial x^{\mu}} \delta v^{\nu} \right] - \\ & - \sum_{\nu} \frac{\partial}{\partial x^{\nu}} \left[\sum_{\mu} \xi^{\nu\mu} v^{\mu} + \xi_{T}^{\nu} \right], \end{aligned}$$

with: n — number density of atoms, g^{ν} — momentum density coordinate, e — energy density, p — pressure, v^{ν} — velocity coordinate, T — temperature, ζ — coefficient of bulk viscosity (also referred to as second viscosity), η — dynamical viscosity coefficient (dynamical viscosity), χ — heat conduction coefficient, $\xi^{\nu\mu}$ — coordinate of stress tensor fluctuation, ξ_{T}^{ν} — coordinate of the fluctuation vector of the thermal energy flux, and m — atomic mass. The superscripts ν , μ denote Cartesian variables.

The above are linear equations of motion for the fluctuations in number density of atoms δn , in momentum density δg^{ν} and in energy density δe . The fluctuations occur in a fluid in motion with the velocity \vec{v} , at non-equilibrium stationary state.

Let us consider a quiescent atomic fluid ($v=0$) in a weakly non-equilibrium, quasi-stationary state caused by a temperature gradient. Earlier, we had assumed the system to evolve towards the state of complete thermodynamical equilibrium during

a time much longer than all the relaxation times of fluctuations about the steady state. This enables us to deal with the quantities of interest as fulfilling, in the steady state, relations of the form specific for the state of equilibrium. It is our aim to derive the fluctuation spectrum of the hydrodynamical fields taking into account the greatest possible number of contributions, due to the effect of the temperature gradient on the viscosity coefficients, the heat conductivity coefficient, as well as the isothermal sound velocity (pressure). Moreover, we shall take into account the fluctuations of these quantities expressed in terms of fluctuations in number density and temperature. Within the above framework and for the sake of maximal generality we shall not restrict ourselves to the isothermal approximation but shall solve the set of all the five equations (1)-(3). In order to distinguish the individual contributions more easily, we have introduced the following underlinings:

- terms related with the temperature gradient,
- _____ terms related with fluctuations of the transport coefficients and isothermal sound velocity expressed via fluctuations in number density and temperature,
- ===== cross terms,
- non-underlined are terms related with the basic fluctuations of the hydrodynamical fields.

To make the eqs. (1)-(3) solvable we apply certain simple thermodynamical relations enabling us to express the fluctuations in energy density and pressure by way of fluctuations in temperature and number density of atoms⁹);

$$\delta e = \frac{e_{ss} + P_{ss}}{n_0} \delta n = -m n_0 c_v \left(\frac{\gamma-1}{\alpha n_0} \delta n - \delta T \right) \quad (4)$$

$$\delta p = m c_T^2 (\delta n + \alpha n_0 \delta T), \quad (5)$$

where α is the volume coefficient of thermal dilation, and γ the Poisson coefficient (the ratio c_p/c_v of the specific heat at constant pressure and at constant volume). The subscripts "ss" and "o" denote respectively steady state values and ones taken at complete thermodynamical equilibrium. Moreover, we express the fluctuations in momentum density in terms of fluctuations in the rate of flow $\delta \vec{v}$:

$$\delta \vec{g} = m n_0 \delta \vec{v}. \quad (6)$$

The assumption made when deriving eqs. (1)-(3) enables us to write the transport coefficients, the temperature, the isothermal sound velocity c_T and the rate of flow in a given point of space \vec{r} as follows:

$$f(\vec{r}) = f_0 + \underbrace{\left(\frac{\partial f}{\partial T} \right) \vec{\nabla} T \cdot \vec{r}}_{\text{-----}} + \underbrace{\delta T \left(\frac{\partial f}{\partial T} \right)}_{\text{-----}} + \underbrace{\left(\frac{\partial f}{\partial n} \right) \delta n}_{\text{-----}} \quad (7)$$

for f respectively equalling ζ , η , χ and c_T^2 . Whereas for the temperature and velocity we have

$$T(\vec{r}) = T_0 + \underbrace{\vec{\nabla} T \cdot \vec{r}}_{\text{-----}} + \delta T \quad (8)$$

$$\vec{v}_{ss}(\vec{r}) = \vec{0}. \quad (9)$$

For brevity, we introduce the following notation:

$$D_v = \frac{\zeta + \frac{4}{3}\eta}{m n_0} \quad (\text{longitudinal kinematic viscosity}), \quad (10)$$

$$D_r = \frac{\chi}{m n_0 c_p} \quad (\text{heat diffusion coefficient}), \quad (11)$$

$$\Psi = \vec{\nabla} \cdot \delta \vec{v} \quad , \quad (12)$$

$$\xi' = \frac{1}{m n_0} \xi \quad , \quad (13)$$

$$\xi'_r = \frac{1}{m n_0 c_p} \xi_r \quad , \quad (14)$$

$$\zeta' = \frac{\zeta}{m n_0} \quad , \quad (15)$$

$$D'_v = \frac{\zeta + \frac{2}{3}\eta}{m n_0} \quad . \quad (16)$$

With regard to (4)-(16) and on introducing the operator nabla eqs. (1)-(3) become

$$\delta \dot{n} + n_0 \Psi = 0 \quad , \quad (17)$$

$$\begin{aligned} \dot{\Psi} = & -\frac{c_r^2}{n_0} \vec{\nabla}^2 \delta n + D_r \vec{\nabla}^2 \Psi - \alpha c_r^2 \vec{\nabla}^2 \delta T - \frac{1}{n_0} \left(\frac{\partial c_r^2}{\partial T} \right) (\vec{\nabla} T \cdot \vec{r}) \vec{\nabla}^2 \delta n - \\ & - \left(\frac{\partial D_r}{\partial T} \right) (\vec{\nabla} T \cdot \vec{r}) \vec{\nabla}^2 \Psi - \alpha \left(\frac{\partial c_r^2}{\partial T} \right) (\vec{\nabla} T \cdot \vec{r}) \vec{\nabla}^2 \delta T + \left(\frac{\partial \zeta'}{\partial T} \right) \vec{\nabla}^2 (\vec{\nabla} T \cdot \vec{r}) \Psi - \\ & - \alpha \left(\frac{\partial c_r^2}{\partial n} \right) \vec{\nabla}^2 (\vec{\nabla} T \cdot \vec{r}) \delta n - \alpha \left(\frac{\partial c_r^2}{\partial T} \right) \vec{\nabla}^2 (\vec{\nabla} T \cdot \vec{r}) \delta T + \left(\frac{\partial D_r}{\partial T} \right) \vec{\nabla} (\vec{\nabla} T \cdot \vec{r}) \vec{\nabla} \Psi + \\ & + \left(\frac{\partial \zeta'}{\partial T} \right) \vec{\nabla} (\vec{\nabla} T \cdot \vec{r}) \vec{\nabla} (\vec{\nabla} T \cdot \vec{r}) \Psi - \alpha c_r^2 \vec{\nabla}^2 (\vec{\nabla} T \cdot \vec{r}) - \alpha \left(\frac{\partial c_r^2}{\partial T} \right) (\vec{\nabla} T \cdot \vec{r}) \vec{\nabla}^2 (\vec{\nabla} T \cdot \vec{r}) - \\ & - \vec{\nabla} \cdot \vec{\xi}' \cdot \vec{\nabla} \quad , \end{aligned} \quad (18)$$

$$\begin{aligned} \delta \dot{T} = & -\frac{\delta^{-1}}{\alpha} \Psi + \delta D_r \vec{\nabla}^2 \delta T + \left(\frac{\partial D_r}{\partial n} \right) (\vec{\nabla}^2 T) \delta n + \left(\frac{\partial D_r}{\partial T} \right) (\vec{\nabla}^2 T) \delta T + \\ & + \frac{1}{3} \left(\frac{\partial \delta D_r}{\partial n} \right) \vec{\nabla} T \cdot \vec{\nabla} \delta n - \frac{1}{3} \left(\frac{\partial \delta D_r}{\partial T} \right) \vec{\nabla} T \cdot \vec{\nabla} \delta T + \frac{1}{3} \left(\frac{\partial \delta D_r}{\partial T} \right) \vec{\nabla} (\vec{\nabla} T \cdot \vec{r}) \cdot \vec{\nabla} \delta T + \\ & + \frac{1}{3} \left(\frac{\partial^2 \delta D_r}{\partial T \partial n} \right) \vec{\nabla} T \cdot \vec{\nabla} (\vec{\nabla} T \cdot \vec{r}) \delta n + \frac{1}{3} \left(\frac{\partial^2 \delta D_r}{\partial T^2} \right) \vec{\nabla} T \cdot \vec{\nabla} (\vec{\nabla} T \cdot \vec{r}) \delta T + \\ & + \delta D_r \vec{\nabla}^2 T + \frac{1}{3} \left(\frac{\partial \delta D_r}{\partial T} \right) \vec{\nabla} (\vec{\nabla} T \cdot \vec{r}) \cdot \vec{\nabla} T - \vec{\nabla} \cdot \vec{\xi}' \quad . \end{aligned} \quad (19)$$

Eqs. (17)-(19) form a set of three equations of the Langevin type of the hydrodynamical fluctuations occurring in the atomic fluid. The system is in a weakly non-equilibrium quasi-stationary state due to the presence of the temperature gradient. The latter has the meaning of an external generalized thermodynamical force.

3. Correlation Functions of the Hydrodynamical Field Fluctuations

The expressions (17)-(19) are linear equations, derivable from the phenomenological equations of hydrodynamics for momentary values of the hydrodynamical variables. Quite obviously, these equations differ from those describing equilibrium fluctuations in that they contain terms which take into account the influence of the temperature gradient and hydrodynamical fluctuations on the transport and sound velocity coefficients. One notes that, in contradistinction to the terms expressing the dependence on the temperature gradient, those related with the influence of the fluctuations on the transport and sound velocity coefficients do not occur as mutually independent quantities. One more-

over notes that eq.(18) contains no terms describing the influence of fluctuations in the transport coefficients. This is due to our assumption that the fluid does not flow as a whole, and the expressions in question always occur as products of the flow rate in the initial equation (2).

The set (17)-(19) will serve to determine the correlation function of hydrodynamical field fluctuations. To this aim, we re-write the equations in matrix form:

$$\check{M} \begin{bmatrix} \delta n \\ \Psi \\ \delta T \end{bmatrix} = - \begin{bmatrix} 0 \\ \vec{\nabla} \cdot \vec{\xi}' \cdot \vec{\nabla} \\ \vec{\nabla} \cdot \vec{\xi}_T \end{bmatrix} + \check{N} \quad (20)$$

involving the matrices \check{M} and \check{N} :

$$\check{M} = \check{A} + \check{B}(\vec{\nabla} T \cdot \vec{r}) \vec{\nabla}^2 + \check{C} \vec{\nabla}^2 (\vec{\nabla} T \cdot \vec{r}) + \check{D} \vec{\nabla} (\vec{\nabla} T \cdot \vec{r}) \vec{\nabla} + \check{E} [\vec{\nabla} (\vec{\nabla} T \cdot \vec{r})]^2, \quad (21)$$

$$\check{N} = \check{F} \vec{\nabla}^2 (\vec{\nabla} T \cdot \vec{r}) + \check{G} (\vec{\nabla} T \cdot \vec{r}) \vec{\nabla}^2 (\vec{\nabla} T \cdot \vec{r}) + \check{H} [\vec{\nabla} (\vec{\nabla} T \cdot \vec{r})]. \quad (22)$$

Whereas

$$\check{A} = \begin{bmatrix} \frac{\partial}{\partial t} & n_0 & 0 \\ \frac{1}{n_0} c_T^2 \vec{\nabla}^2 & \frac{\partial}{\partial t} - D_T \vec{\nabla} & \alpha c_T^2 \vec{\nabla}^2 \\ 0 & \frac{1}{\alpha} (\gamma - 1) \frac{\partial}{\partial t} - \delta D_T \vec{\nabla} \end{bmatrix} \quad \check{B} = \begin{bmatrix} 0 & 0 & 0 \\ \frac{1}{n_0} \left(\frac{\partial c_T^2}{\partial T} \right) \left(\frac{\partial D_T}{\partial T} \right) & \alpha \left(\frac{\partial c_T^2}{\partial T} \right) & \\ 0 & 0 & 0 \end{bmatrix}$$

$$\check{C} = \begin{bmatrix} 0 & 0 & 0 \\ \alpha \left(\frac{\partial c_T^2}{\partial n} \right) - \left(\frac{\partial \xi'}{\partial T} \right) & \alpha \left(\frac{\partial c_T^2}{\partial T} \right) & \\ - \left(\frac{\partial \delta D_T}{\partial n} \right) & 0 & \left(\frac{\partial \delta D_T}{\partial T} \right) \end{bmatrix} \quad \check{D} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \left(\frac{\partial D_T}{\partial T} \right) & 0 \\ \frac{1}{3} \left(\frac{\partial \delta D_T}{\partial n} \right) & 0 & \frac{1}{3} (1 + \gamma) \left(\frac{\partial \delta D_T}{\partial T} \right) \end{bmatrix}$$

$$\check{E} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \left(\frac{\partial \xi'}{\partial T^2} \right) & 0 \\ \frac{1}{3} \left(\frac{\partial \delta D_T}{\partial n \partial T} \right) & 0 & \frac{1}{3} \left(\frac{\partial \delta D_T}{\partial T^2} \right) \end{bmatrix} \quad \check{F} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & -\alpha c_T^2 & 0 \\ 0 & \delta D_T & 0 \end{bmatrix}$$

$$\check{G} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & -\alpha \left(\frac{\partial c_T^2}{\partial T} \right) & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \check{H} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & \frac{1}{3} \left(\frac{\partial \delta D_T}{\partial T} \right) & 0 \end{bmatrix} \quad (23)$$

Prior to applying the Fourier transform to eq. (20) we shall make an assumption, often used in quantum perturbation calculus. In order to ensure a slow evolution of the system towards equilibrium via quasi-stationary states we assume the temperature gradient to be an appropriately small quantity. Thus, the difference in temperature ΔT between the boundary regions of the medium distant by L of the order of 1 cm is a quantity of the order of 1 K fulfilling the inequality $\frac{\Delta T}{T} \ll 1$. Thus, we are justified in writing

$$T_{ss}(\vec{r}) = T_0(\vec{r} = \vec{0}) + \delta T \sin q \cdot \vec{r}, \quad (24)$$

where $q \delta T = \vec{\nabla} T$, $q = \frac{1}{L}$.

We consider a sample the L dimensional of which are much greater than the correlation range $|\vec{r}|$ and the free path of the hydrodynamical modes due to the fluctuations. We thus obtain nonzero contributions to the correlation function for $|\vec{r}| \ll L$ only. For \vec{r} - values satisfying this condition, the product $\vec{q} \cdot \vec{r}$ takes small values and (24) becomes a good approximation in (8). In addition, the above condition enables us to neglect the influence of boundary effects on the processes under consideration and permits the integration of (20) in the limits $(-\infty, +\infty)$. We now obtain the Fourier transform of (20) in the following general, compact form:

$$\begin{bmatrix} \delta n_{\vec{k}} \\ \Psi_{\vec{k}} \\ \delta T_{\vec{k}} \end{bmatrix} = \frac{1}{D(\vec{k}, \vec{q})} \sum_{\vec{k}''} \check{M}_{\vec{k}''}^{-1} \begin{bmatrix} 0 \\ \vec{k} \cdot \vec{\xi}_{\vec{k}''} \cdot \vec{k} \\ i \vec{k} \cdot \vec{\xi}_{\vec{k}''} \end{bmatrix} \quad (25)$$

To obtain the correlation function for the hydrodynamical fields we write the conjugate of the matrix (25):

$$[\delta n_{\vec{k}}^* \ \Psi_{\vec{k}}^* \ \delta T_{\vec{k}}^*] = \frac{1}{D^*(\vec{k}, \vec{q})} \sum_{\vec{k}''} [0 \ \vec{k} \cdot \vec{\xi}_{\vec{k}''} \cdot \vec{k} \ -i \vec{k} \cdot \vec{\xi}_{\vec{k}''}] (\check{M}_{\vec{k}''}^{-1})^* \quad (26)$$

Next, multiplying (25) and (26) and taking the statistical average, we arrive at the correlation functions. Our results, when tabulated, assume the general form:

$$\begin{bmatrix} \langle \delta n_{\vec{k}} \delta n_{\vec{k}}^* \rangle \langle \delta n_{\vec{k}} \Psi_{\vec{k}}^* \rangle \langle \delta n_{\vec{k}} \delta T_{\vec{k}}^* \rangle \\ \langle \Psi_{\vec{k}} \delta n_{\vec{k}}^* \rangle \langle \Psi_{\vec{k}} \Psi_{\vec{k}}^* \rangle \langle \Psi_{\vec{k}} \delta T_{\vec{k}}^* \rangle \\ \langle \delta T_{\vec{k}} \delta n_{\vec{k}}^* \rangle \langle \delta T_{\vec{k}} \Psi_{\vec{k}}^* \rangle \langle \delta T_{\vec{k}} \delta T_{\vec{k}}^* \rangle \end{bmatrix} = \frac{1}{|D(\vec{k}, \vec{q})|^2} \sum_{\vec{k}''} \begin{bmatrix} [\alpha'_{12} \alpha''_{12} \beta_{\vec{k}''} - \alpha'_{13} \alpha''_{13} \beta_{\vec{k}''}] \\ [\alpha'_{22} \alpha''_{22} \beta_{\vec{k}''} - \alpha'_{23} \alpha''_{23} \beta_{\vec{k}''}] \\ [\alpha'_{32} \alpha''_{32} \beta_{\vec{k}''} - \alpha'_{33} \alpha''_{33} \beta_{\vec{k}''}] \end{bmatrix}$$

$$\begin{bmatrix} [\alpha'_{12} \alpha''_{12} \beta_{\vec{k}''} - \alpha'_{13} \alpha''_{13} \beta_{\vec{k}''}] [\alpha'_{12} \alpha''_{12} \beta_{\vec{k}''} - \alpha'_{13} \alpha''_{13} \beta_{\vec{k}''}] \\ [\alpha'_{22} \alpha''_{22} \beta_{\vec{k}''} - \alpha'_{23} \alpha''_{23} \beta_{\vec{k}''}] [\alpha'_{22} \alpha''_{22} \beta_{\vec{k}''} - \alpha'_{23} \alpha''_{23} \beta_{\vec{k}''}] \\ [\alpha'_{32} \alpha''_{32} \beta_{\vec{k}''} - \alpha'_{33} \alpha''_{33} \beta_{\vec{k}''}] [\alpha'_{32} \alpha''_{32} \beta_{\vec{k}''} - \alpha'_{33} \alpha''_{33} \beta_{\vec{k}''}] \end{bmatrix} \quad (27)$$

The terms α'_{ij} , α''_{ij} are functions of the transport coefficients and their derivatives. On restricting ourselves to terms quadratic in \vec{q} we get the respective functions in the form shown in Appendix A.

The $\beta_{\vec{k}''}$, $\beta_{\vec{k}''}$ are correlation functions of the stochastic terms of the equations which describe, respectively, the fluctuation in velocity $\Psi_{\vec{k}}^*$ and in temperature $\delta T_{\vec{k}}^*$:

$$\beta_{\vec{k}''} \equiv \langle \vec{k} \cdot \vec{\xi}_{\vec{k}''} \cdot \vec{k} \ \vec{k} \cdot \vec{\xi}_{\vec{k}''}^* \cdot \vec{k} \rangle \quad (28)$$

$$\beta_{\vec{k}''} \equiv \langle \vec{k} \cdot \vec{\xi}_{\vec{k}''} \ \vec{k} \cdot \vec{\xi}_{\vec{k}''}^* \cdot \vec{k} \rangle \quad (29)$$

Let us assume that there are no correlations between the fluctuations of the stress tensor and thermal energy density flux vector. Thus, the cross correlation functions vanish. The correlation functions $\beta_{\vec{k}''}$, $\beta_{\vec{k}''}$, determined in Ref.²⁴, ——— similarly to the elements α'_{ij} , α''_{ij} ——— are func-

tions of the transport coefficients and their derivatives, dependent on \vec{q} and the temperature. In ²⁴⁾, the wave vectors

\vec{k}', \vec{k}'' are shown to take the values

$$\vec{k}', \vec{k}'' = \vec{k} \pm n\vec{q} \quad n = \begin{cases} 0, 1, 2 & \text{for } \beta_{\vec{k}', \vec{k}''} \\ 0, 1, 2, 3 & \text{for } \beta_{\tau, \vec{k}', \vec{k}''} \end{cases} \quad (30)$$

Thus, the above correlation functions can now be written as

$$\left. \begin{aligned} \beta_{\vec{k}', \vec{k}''} &= \beta_{\vec{k} + n\vec{q}, \vec{k} + n\vec{q}} = \beta_{n'l} \\ \beta_{\tau, \vec{k}', \vec{k}''} &= \beta_{\tau, \vec{k} + n\vec{q}, \vec{k} + n\vec{q}} = \beta_{\tau, n'l} \end{aligned} \right\} \text{for } l = n' - n' \quad (31)$$

With accuracy to terms quadratic in \vec{q} we obtain ²⁴⁾

$$\begin{aligned} \beta_{n'0} &= 2k_B T m n_0 D_v (\vec{k} + n\vec{q})^4 \delta(\omega' - \omega) + k_B \delta T^2 m n_0 \left(\frac{\partial D_v}{\partial T} \right) (\vec{k} + n\vec{q})^4 \delta(\omega' - \omega), \\ \beta_{n'+1} &= \mp k_B \delta T^2 \left[2\eta \left(1 + \frac{\partial \ln \eta}{\partial \ln T} \right) (\vec{k} + n\vec{q})^2 [\vec{k} + (n \pm 1)\vec{q}]^2 + \right. \\ &\quad \left. + \left(\zeta - \frac{2}{3}\eta \right) \left(1 + \frac{\partial \ln (\zeta - \frac{2}{3}\eta)}{\partial \ln T} \right) (\vec{k} + n\vec{q})^2 [\vec{k} + (n \pm 1)\vec{q}]^2 \delta(\omega' - \omega) \right], \\ \beta_{n' \pm 2} &= \frac{(\delta T)^2}{2} k_B \left[2 \left(\frac{\partial \eta}{\partial T} \right) [(\vec{k} + n\vec{q})^2 (\vec{k} + (n \pm 2)\vec{q})]^2 + \right. \\ &\quad \left. + \left(\frac{\partial (\zeta - \frac{2}{3}\eta)}{\partial T} \right) (\vec{k} + n\vec{q})^2 (\vec{k} + (n \pm 2)\vec{q})^2 \right] \delta(\omega' - \omega), \quad (32) \\ \beta_{\tau, n'0} &= 2k_B T^2 \chi (\vec{k} + n\vec{q})^2 \delta(\omega' - \omega) + k_B \chi \delta T^2 \left(1 + 2 \frac{\partial \ln \chi}{\partial \ln T} \right) (\vec{k} + n\vec{q})^2 \delta(\omega' - \omega), \\ \beta_{\tau, n' \pm 1} &= \left[\mp i k_B T \chi \delta T^2 \left(2 + \frac{\partial \ln \chi}{\partial \ln T} \right) (\vec{k} + n\vec{q}) [\vec{k} + (n \pm 1)\vec{q}] \mp \right. \\ &\quad \left. \mp i \frac{3}{4} k_B \chi \left(\frac{\delta T}{T} \right)^2 \left(\frac{\partial \ln \chi}{\partial \ln T} \right) (\vec{k} + n\vec{q}) [\vec{k} + (n \pm 1)\vec{q}] \right] \delta(\omega' - \omega), \\ \beta_{\tau, n' \pm 2} &= -\frac{1}{2} k_B \chi (\delta T)^2 \left(1 + 2 \frac{\partial \ln \chi}{\partial \ln T} \right) (\vec{k} + n\vec{q}) [\vec{k} + (n \pm 2)\vec{q}] \delta(\omega' - \omega). \end{aligned}$$

In order to write $|D(\vec{k}, \vec{q})|^{-2}$ in the generally accepted form we have to solve the frequency dispersion equation

$$D(\vec{k}, \vec{q}, s) = 0, \quad (33)$$

where $D(\vec{k}, \vec{q}, s)$ is the value of the determinant of the Fourier transform of the matrix \check{M} of eq.(20).

The solutions of (33) take one form or another for different values of \vec{k} . This is so because, according to the order of accuracy assumed, certain terms do or do not occur in the solution depending on the value of \vec{k} . The strict solution of this equation for the case under consideration is to be found in Ref.²⁴⁾. For \vec{k} in the optical range, i.e., for $k \approx 10^5 \text{ cm}^{-1}$, and for an accuracy restricted to the first terms in q^2 , we obtain (see Appendix B):

$$\begin{aligned} |D(\vec{k}, \vec{q})|^{-2} &= |D(\vec{k})|^{-2} \left\{ 1 - 2 \left[\frac{(D_\tau k^2)^2}{\omega^2 + (D_\tau k^2)^2} (G_4 - \frac{\omega}{D_\tau k^2} G_2) + \right. \right. \\ &\quad \left. \left. + \frac{\omega c_s k + C_s^2 k^2 + (\Gamma k^2)^2}{(\omega + c_s k)^2 + (\Gamma k^2)^2} (G_4 + G_5) + \right. \right. \\ &\quad \left. \left. + \frac{[\omega c_s k - C_s^2 k^2 - (\Gamma k^2)^2] (G_5 - G_4) + \omega \Gamma k^2 (G_6 + G_7 + G_8)}{(\omega - c_s k)^2 + (\Gamma k^2)^2} \right. \right. \\ &\quad \left. \left. + \frac{\omega \Gamma k^2}{(\omega + c_s k)^2 + (\Gamma k^2)^2} (G_6 + G_7 + G_8) \right] \right\}. \quad (34) \end{aligned}$$

The corrections G_i , $i=1, 2, \dots, 8$, as well as the forms of $|D(\vec{k})|^{-2}$ are given in Appendix B (eqs. (B13)-(B20) and (B.34)).

The essential result of our work resides in the relation (9), which summarizes the set of all possible cases of the hydrodynamical field correlation functions.

A similar calculation procedure has been applied in Refs. 13-15). However, their authors have not performed systematical calculations taking into account the greatest possible number of contributions to the hydrodynamical field correlation function, restricting themselves to a discussion of the influence of the temperature gradient on the velocity of sound, the coefficients of viscosity, and the heat conductivity coefficient, and neglecting the fluctuations of these quantities. Also, they do not go beyond solving the dispersion equation with accuracy to the second order of perturbation calculus. Applying the isothermal approximation, they omit the changes in entropy; this leads to a set of initial equations containing no equation of motion for the fluctuations in temperature. Also, the correlation functions of the stochastic terms of the hydrodynamical equations used there are given a presentation more intuitive than resulting from thorough calculations. All this leads to the circumvention of certain conditions which amount to specific selection rules on the values of n' and n'' and eliminate arbitrariness in the construction of the final expressions strict premis.

4. Conclusion

We have been considering a system in a state of non-equilibrium due to the introduction of a temperature gradient. We have assumed as satisfied all the conditions for this state to be stationary. A system like this can be realized with a high degree of accuracy by surrounding the medium on either side with reservoirs as two different temperatures giving a slight gradient, decaying slowly compared with the lifetimes of the processes taking place in the system.

We make use of hydrodynamical equations of the most general form:

- (i) taking into account the influence of the temperature gradient on the viscosity coefficients ζ and η , the heat conductivity coefficient χ , and the sound velocity C_s ,
- (ii) including the fluctuations in sound velocity and heat conductivity coefficient expressed via fluctuations of the fundamental hydrodynamical fields — those of temperature and number density, and
- (iii) not restricting ourselves to the isothermal approximation, and taking into consideration all three hydrodynamical fields, and
- (iv) performing our calculations to terms in q^2 .

We have transformed the set of five equations to variables \vec{k} , ω and have solved it analytically. We have expressed the hydrodynamical field fluctuations by way of stochastic functions with various wave vectors, resulting from our calculations. The functional coefficients that occur are dependent on the parameters characterizing the medium and on their derivatives with respect to temperature, density and pressure. The determination of the correlation function consisted primarily in solving the frequency dispersion equation and calculating the correlation function for the stochastic terms of the initial equations. It is necessary to stress, as highly important in all cases, the dependence on the concrete value of the wave vectors, of the general form $\vec{k} = \vec{k} + n\vec{q}$ (for the conjugate part, $\vec{k}'' = \vec{k} + n''\vec{q}$). It is essential that the values taken by n' and n'' are a consequence

of form derived by us for the initial equations and are by no means but an attempt at generalizing the expressions indexed by the \vec{k} . Similarly, the set of n' and n'' values occurring as indices at the correlation functions of the stochastic terms of the initial equations is the result of strict, mathematical operations and thus derives from a mathematical and physical necessity rather than from a bare possibility.

We have introduced distinctions between terms related with contributions of different kinds:

1. contributions from fluctuations in number density, momentum and energy,
2. contributions appearing due to our having taken into account fluctuations in sound velocity and heat conductivity coefficient, and
3. ones related with the presence of the temperature gradient.

Only contributions of the type 1, specific to equilibrium fluctuations, are found to occur independently. Thus cross terms, being products of contributions of the types 1,3 and 2,3, appear as well.

In part II, now in preparation, we shall discuss the influence of the above derived contributions on the spectrum of scattered light.

Appendix A.

In tabulated form, we give the terms α'_{ij} occurring in eq.(27). The elements α'_{ij} are derived from the α'_{ij} by the interchange: $\vec{k}' \rightarrow \vec{k}''$, $\vec{k} \rightarrow \vec{k}'$. Each of the functions α'_{ij} is the sum of products of the terms in the corresponding row and the appropriate expressions of the first row.

For example,

$$\alpha'_{23} = -\alpha C_T^2 k^2 \delta(\vec{k}-\vec{k}') + \alpha s(1+1) \left(\frac{\partial m C_T^2 k^2}{\partial T} \right) \frac{\delta \vec{T}}{2} \left\{ 1+2 \left(\frac{q}{k} \right)^2 \delta[\vec{k}'-(\vec{k}-\vec{q})] + 2 \left(\frac{k \cdot \vec{q}}{k^2} \right) \right\}$$

$\delta(\vec{k}'-\vec{k})$	$i \frac{\delta \vec{T}}{2} \left\{ \left[1-2 \frac{\vec{k} \cdot \vec{q}}{k^2} + \left(\frac{q}{k} \right)^2 \right] \delta[\vec{k}'-(\vec{k}+\vec{q})] - \left[1-2 \frac{\vec{k} \cdot \vec{q}}{k^2} + \left(\frac{q}{k} \right)^2 \right] \delta[\vec{k}'-(\vec{k}-\vec{q})] \right\}$	$i \frac{\delta \vec{T}}{2} \left\{ \frac{\vec{k} \cdot \vec{q}}{k^2} + \left(\frac{q}{k} \right)^2 \delta[\vec{k}'-(\vec{k}+\vec{q})] - \left[-\frac{\vec{k} \cdot \vec{q}}{k^2} + \left(\frac{q}{k} \right)^2 \right] \delta[\vec{k}'-(\vec{k}-\vec{q})] \right\}$	$\frac{(\delta \vec{T})^2}{4} \left(\frac{q}{k} \right)^2 \left[2 \delta(\vec{k}-\vec{k}') + \delta[\vec{k}'-(\vec{k}+2\vec{q})] - \delta[\vec{k}'-(\vec{k}-2\vec{q})] \right]$
$\alpha'_{12} - n_0 (s + \delta D_T k^2)$	$n_0 \left(\frac{\partial \delta D_T k^2}{\partial T} \right)$	$\frac{1}{3} n_0 (1+1) \left(\frac{\partial \delta D_T k^2}{\partial T} \right)$	$n_0 \left(\frac{\partial^2 \delta D_T k^2}{\partial T^2} \right)$
$\alpha'_{13} - \alpha n_0 C_T^2 k^2$	$\alpha n_0 (1+1) \left(\frac{\partial C_T^2 k^2}{\partial T} \right)$	0	0
$\alpha'_{22} s (s + \delta D_T k^2)$	$-s \left(\frac{\partial \delta D_T k^2}{\partial T} \right)$	$\frac{1}{3} (1+1) s \left(\frac{\partial \delta D_T k^2}{\partial T} \right)$	$s \left(\frac{\partial^2 \delta D_T k^2}{\partial T^2} \right)$
$\alpha'_{23} - \alpha C_T^2 k^2$	$\alpha s (-1+1) \left(\frac{\partial m C_T^2 k^2}{\partial T} \right)$	0	0
$\frac{\delta-1}{\alpha}$	$n_0 \left(\frac{\partial \delta D_T k^2}{\partial T} \right)$	$\frac{1}{3} n_0 \left(\frac{\partial \delta D_T k^2}{\partial T} \right)$	$-\beta n_0 \left(\frac{\partial^2 \delta D_T k^2}{\partial T^2} \right)$
$\alpha'_{33} C_T^2 k^2 + s (s + D_V k^2)$	$s \left(\frac{\partial D_V k^2}{\partial T} \right) - \frac{\partial C_T^2 k^2}{\partial T} - \alpha n_0 \left(\frac{\partial C_T^2 k^2}{\partial T} \right)$	0	$-3s \left(\frac{\partial^2 C_T^2 k^2}{\partial T^2} \right)$

Appendix B

The frequency dispersion equation has the form⁽²³⁾

$$\begin{aligned} D(\vec{k}, \vec{q}, s) = & s^3 + [D_v k^2 + \gamma D_r k^2 + i(a_5 - a_7) + a_6 + a_8] s^2 + \\ & + [\delta c_r^2 k^2 + D_v k^2 \gamma D_r k^2 + D_v k^2 a_8 + \gamma D_r k^2 a_6 + a_6 a_8 + a_5 a_7 - \\ & - i(a_7 + (\gamma - 1)a_2 - \gamma D_r k^2 a_5 - D_v k^2 a_7 - a_5 a_6 + a_6 a_7)] s + \\ & + \alpha n_0 c_r^2 k^2 a_4 + c_r^2 k^2 \gamma D_r k^2 - c_r^2 k^2 a_8 - a_1 a_7 + \alpha n_0 a_2 a_3 + \\ & + i(\alpha n_0 c_r^2 k^2 a_3 - c_r^2 k^2 a_7 - \gamma D_r k^2 a_1 - a_1 a_8 - \alpha n_0 a_2 a_4) = 0, \end{aligned} \quad (B.1)$$

with

$$a_1 = 2\delta\tilde{T} \left[\frac{(\partial c_r^2 k^2)}{(\partial T)} + \alpha n_0 \frac{(\partial c_r^2 k^2)}{(\partial n)} \right] \frac{\vec{k} \cdot \vec{q}}{k^2}, \quad (B.2)$$

$$a_2 = 2\delta\tilde{T} (1 + \gamma) \frac{(\partial c_r^2 k^2)}{(\partial T)} \frac{\vec{k} \cdot \vec{q}}{k^2}, \quad (B.3)$$

$$a_3 = 2\delta\tilde{T} \frac{(\partial \gamma D_r k^2)}{(\partial n)} \left[\frac{\vec{k} \cdot \vec{q}}{k^2} - \frac{1}{3} \left(\frac{q}{k} \right)^2 \right], \quad (B.4)$$

$$a_4 = \frac{(\delta\tilde{T})^2}{6} \frac{(\partial^2 \gamma D_r k^2)}{(\partial T \partial n)} \left(\frac{q}{k} \right)^2 \quad (B.5)$$

$$a_5 = 2\delta\tilde{T} \frac{(\partial D_v k^2)}{(\partial T)} \frac{\vec{k} \cdot \vec{q}}{k^2} - \delta\tilde{T} \frac{(\partial D_r k^2)}{(\partial T)} \frac{\vec{k} \cdot \vec{q}}{k^2} - \frac{\delta\tilde{T}}{2} \frac{(\partial \gamma k^2)}{(\partial T)} \left(\frac{q}{k} \right)^2, \quad (B.6)$$

$$a_6 = \frac{(\delta\tilde{T})^2}{4} \frac{(\partial^2 \gamma k^2)}{(\partial T^2)} \left(\frac{q}{k} \right)^2, \quad (B.7)$$

$$a_7 = \frac{1}{3} \delta\tilde{T} (1 + \gamma) \frac{(\partial \gamma D_r k^2)}{(\partial T)} \frac{\vec{k} \cdot \vec{q}}{k^2}, \quad (B.8)$$

$$a_8 = \frac{(\delta\tilde{T})^2}{12} \frac{(\partial^2 D_r k^2)}{(\partial T^2)} \left(\frac{q}{k} \right)^2. \quad (B.9)$$

In Ref.⁽²³⁾, eq. (B.1) is solved by well-known methods of perturbation calculus. On restricting ourselves to an accuracy determined by the order of the first terms quadratic in \vec{q} we get, for

$$k \approx 10^5 \text{ cm}^{-1}$$

$$s_1 = -D_r k^2 [1 + G_1 + i G_2], \quad (B.10)$$

$$s_2 = (i c_s k + \Gamma k^2) \left[1 + \frac{1}{i c_s k + \Gamma k^2} (i G_8 + G_4 + G_5 + i G_6 + i G_7) \right] \quad (B.11)$$

$$s_3 = (-i c_s k + \Gamma k^2) \left[1 - \frac{1}{-i c_s k + \Gamma k^2} (i G_8 - G_4 + G_5 + i G_6 - i G_7) \right], \quad (B.12)$$

where

$$G_1 = (\gamma - 1) \frac{D_r k^2}{c_s^2 k^2} (D_r k^2 - D_v k^2), \quad (B.13)$$

$$G_2 = 4 \left(1 - \frac{1}{\gamma} \right) \frac{\delta\tilde{T}}{c_s k} \left[\frac{(\partial c_s k)}{(\partial T)} - \alpha n_0 \frac{(\partial c_s k)}{(\partial n)} \right] \frac{\vec{k} \cdot \vec{q}}{k^2} \quad (B.14)$$

$$G_3 = 2\delta\tilde{T} \frac{1}{\gamma} [1 + (\gamma - 1)] \left[\frac{(\partial c_s k)}{(\partial T)} + \alpha n_0 \frac{(\partial c_s k)}{(\partial n)} \right] \frac{\vec{k} \cdot \vec{q}}{k^2}, \quad (B.15)$$

$$G_4 = \frac{1}{c_s^2 k^2} [b_1 (D_r k^2)^3 + b_2 (D_r k^2)^2 D_v k^2 + b_3 D_r k^2 (D_v k^2)^2 + (D_v k^2)^3], \quad (B.16)$$

$$G_5 = \delta \tilde{T} \left[1 + \left(1 + \frac{1}{\gamma} \right) \right] \left[\left(\frac{\partial c_s k}{\partial T} \right) + \frac{\alpha n_0}{\gamma} \left(\frac{\partial c_s k}{\partial n} \right) \right] \frac{\vec{k} \cdot \vec{q}}{k^2} \quad (\text{B.17})$$

$$G_6 = \frac{1}{c_s^3 k^3} \left[b_4 (D_T k^2)^4 - b_5 (D_T k^2)^3 D_V k^2 + b_6 (D_T k^2 D_V k^2)^2 - b_7 D_T k^2 (D_V k^2)^3 - (D_V k^2)^4 \right], \quad (\text{B.18})$$

$$G_7 = \frac{\delta T}{c_s k} \left[(b_8 D_T k^2 - b_9 D_V k^2) \left(\frac{\partial c_s k}{\partial T} \right) - (b_{10} D_T k^2 + b_{11} D_V k^2) \alpha n_0 \left(\frac{\partial c_s k}{\partial n} \right) \right] \frac{\vec{k} \cdot \vec{q}}{k^2}, \quad (\text{B.19})$$

$$G_8 = \frac{1}{c_s k} \left[b_{12} (D_T k^2)^2 + b_{13} D_T D_V k^4 + \frac{1}{8} (D_V k^2)^2 \right]. \quad (\text{B.20})$$

The coefficients b_i , $i=1,2,\dots,11$, occurring in (B.13)-(B.20) are functions of the Poisson coefficient γ and have the form

$$b_1 = \frac{1}{16} (4\gamma^3 - 19\gamma^2 + 19\gamma + 4), \quad (\text{B.21})$$

$$b_2 = \frac{1}{8} (46\gamma^2 - 72\gamma + 19), \quad (\text{B.22})$$

$$b_3 = \frac{1}{8} (12\gamma - 19), \quad (\text{B.23})$$

$$b_4 = \frac{1}{64} (-\gamma^4 - 1348\gamma^3 - 2350\gamma^2 - 1156\gamma + 279), \quad (\text{B.24})$$

$$b_5 = \frac{1}{32} (1006\gamma^3 - 4142\gamma^2 + 3450\gamma - 578), \quad (\text{B.25})$$

$$b_6 = \frac{1}{32} (2126\gamma^2 + 4142\gamma - 1175), \quad (\text{B.26})$$

$$b_7 = \frac{1}{16} (503\gamma - 337), \quad (\text{B.27})$$

$$b_8 = - \left[3\gamma - \left(\frac{7}{3} - 1 \right) - 4 \left(1 - \frac{1}{\gamma} \right) \frac{1}{\gamma} \right], \quad (\text{B.28})$$

$$b_9 = 3 \left[\frac{1}{\gamma} + \left(1 - \frac{1}{\gamma} \right) \right], \quad (\text{B.29})$$

$$b_{10} = 4 \left(1 - \frac{1}{\gamma} \right), \quad (\text{B.30})$$

$$b_{11} = \frac{3}{\gamma}. \quad (\text{B.31})$$

The underlinings are in accordance with the convention introduced by us in Section 2.

References

- 1) S.R.Groot and P.Mazur, *Non-Equilibrium Thermodynamics* (North-Holland, Amsterdam, 1962).
- 2) D.N.Zubarev, *Nonequilibrium Statistical Thermodynamics* (in Russian) (Moscow, Nauka, 1971).
- 3) R.Kubo, *J.Phys.Soc. Japan* 12 (1957) 570.
R.Kubo, M.Yokota, S.Nakajima, *J.Phys.Soc. Japan* 12 (1957) 1203.
- 4) L.Onsager, *Phys.Rev.* 37 (1931) 405; 38 (1931) 2265.
- 5) N.N.Bogolubov, *Problemy dinamicheskoy teorii w statisticheskoy fizike* (in Russian) (Problems of the Dynamical Theory of Statistical Physica) (Moscow-Leningrad, Gostekhizd., 1946).
- 6) M.Smoluchowski, *Ann. d. Phys.* 25 (1908) 205.
- 7) J.Kocinski, L.Wojtczak, *Critical Scattering Theory* (PWN-Elsevier, Warszawa-Amsterdam, 1978).
- 8) K.Knast, W.Chmielowski, *J.Physique*, 42 (1981) 1373.
- 9) B.J.Berne, R.Pecora, *Dynamic Light Scattering* (Wiley, New York, 1976).
- 10) B.Crosignani, P.DiPorto, M.Bertolotti, *Statistical Properties of Scattered Light* (Academic, Press, New York, 1975).
- 11) P.Lallemand, in: *Photon Correlation and Light Beating Spectroscopy* (E. R. Pike, Plenum Press, New York, 1974).

- 12) S.Kielich, Nonlinear Molecular Optics (in Russian) (Izd. Nauka, Moscow, 1981).
- 13) G. van der Zwan, P.Mazur, Phys.Lett. 75A (1980) 370; Physica 107A (1981) 491.
- 14) A.M.S.Tremblay, E.D.Siggia, M.Arai, Phys.Lett. 76A (1980) 57. A.M.S.Tremblay, M.Arai, E.D.Siggia, Phys. Rev. A23 (1981) 1451.
- 15) D.Ronis, S.Putterman, Phys.Rev. A22 (1980) 773.
- 16) I.Procaccia, D.Ronis, I.Oppenheim, Phys.Rev.Lett. 42 (1979) 287, 614E.
- 17) I.Procaccia, D.Ronis, M.A.Collins, J.Ross, I.Oppenheim, Phys.Rev. A19 (1979) 1290. D.Ronis, I.Procaccia, I.Oppenheim, Phys.Rev. A19 (1979) 1307. D.Ronis, I.Procaccia, I.Oppenheim, Phys.Rev. A19 (1979) 1324.
- 18) T.Kirkpatrick, E.G.D.Cohen, J.R.Dorfman, Phys.Rev.Lett. 42 (1979) 1368; 44 (1980) 472. T.Kirkpatrick, E.G.D.Cohen, J.R.Dorfman, Phys.Rev. A26 (1982) 950, 995, 972.
- 19) H.Grabert, Projection Operator Techniques in Nonequilibrium Statistical Mechanics (Springer-Verlag, Berlin, 1982).
- 20) J.Keizer, Phys.Fluids 21 (1978) 198.
- 21) F.L.Hinton, Phys.Fluids 13 (1970) 857.
- 22) A.Onuki, J.Stat.Phys. 18 (1978) 475.
- 23) H.Ueyama, J.Stat.Phys. 22 (1980) 1.
- 24) W.Chmielowski, Thesis (unpublished) 1985.

Received by Publishing Department
on July 5, 1988.

WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices -- in US \$, including the packing and registered postage.

D13-84-63	Proceedings of the XI International Symposium on Nuclear Electronics. Bratislava, Czechoslovakia, 1983.	12.00
E1,2-84-160	Proceedings of the 1983 JINR-CERN School of Physics. Tabor, Czechoslovakia, 1983.	6.50
D2-84-366	Proceedings of the VII International Conference on the Problems of Quantum Field Theory. Alushta, 1984.	11.00
D1,2-84-599	Proceedings of the VII International Seminar on High Energy Physics Problems. Dubna, 1984.	12.00
D17-84-850	Proceedings of the III International Symposium on Selected Topics in Statistical Mechanics. Dubna, 1984 (2 volumes).	22.00
	Proceedings of the IX All-Union Conference on Charged Particle Accelerators. Dubna, 1984. (2 volumes)	25.00
D11-85-791	Proceedings of the International Conference on Computer Algebra and Its Applications in Theoretical Physics. Dubna, 1985.	12.00
D13-85-793	Proceedings of the XII International Symposium on Nuclear Electronics, Dubna, 1985.	14.00
D4-85-851	Proceedings of the International School on Nuclear Structure Alushta, 1985.	11.00
D1,2-86-668	Proceedings of the VIII International Seminar on High Energy Physics Problems, Dubna, 1986 (2 volumes)	23.00
D3,4,17-86-747	Proceedings of the V International School on Neutron Physics. Alushta, 1986.	25.00
D9-87-105	Proceedings of the X All-Union Conference on Charged Particle Accelerators. Dubna, 1986 (2 volumes)	25.00
D7-87-68	Proceedings of the International School-Seminar on Heavy Ion Physics. Dubna, 1986.	25.00
D2-87-123	Proceedings of the Conference "Renormalization Group-86". Dubna, 1986.	12.00
D4-87-692	Proceedings of the International Conference on the Theory of Few Body and Quark-Hadronic Systems. Dubna, 1987.	12.00
D2-87-798	Proceedings of the VIII International Conference on the Problems of Quantum Field Theory. Alushta, 1987.	10.00
D14-87-799	Proceedings of the International Symposium on Muon and Pion Interactions with Matter. Dubna, 1987.	13.00
D17-88-95	Proceedings of the IV International Symposium on Selected Topics in Statistical Mechanics. Dubna, 1987.	14.00

Orders for the above-mentioned books can be sent at the address:
Publishing Department, JINR
Head Post Office, P.O.Box 79 101000 Moscow, USSR

SUBJECT CATEGORIES OF THE JINR PUBLICATIONS

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches
19.	Biophysics

Хмельовски В., Кнаст К., Келих Ст.

E17-88-488

Флуктуации гидродинамического поля в неравновесном квазистационарном состоянии с градиентом температуры

1. Общая теория

Рассматривается система в неравновесном состоянии с градиентом температуры. Целью исследований является получение в общем виде корреляционных функций флуктуации для гидродинамических полей. Введены обозначения, позволяющие различать вклады от флуктуации плотности числа частиц и энергии, вклады от флуктуации скорости звука и коэффициента теплопроводности, а также вклады, связанные с градиентом температуры.

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Сообщение Объединенного института ядерных исследований. Дубна 1988

Chmielowski W., Knast K., Kielich S.

E17-88-488

Hydrodynamical Field Fluctuations in Non-Equilibrium Quasi-Stationary State Due to Temperature Gradient

1. General Theory

We consider a system in a nonequilibrium state due to the introduction of a temperature gradient. The essential problem of this paper is to determine, in their most general form, the fluctuation correlation functions for hydrodynamical fields. We use notation which enables us to distinguish the contributions from the individual mechanisms: contributions from fluctuations in number density, momentum and energy; contributions appearing due to our having taken into account fluctuations in sound velocity and heat conductivity coefficient; and ones related with the presence of the temperature gradient.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Communication of the Joint Institute for Nuclear Research. Dubna 1988