


89-100



**ОБЪЕДИНЕННЫЙ  
ИНСТИТУТ  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ  
ДУБНА**

S 38

E17-89-100

R.Schumann

**ON THE FUNCTIONAL  
INTEGRAL METHOD APPLIED TO MODELS  
WITH HUBBARD TYPE INTERACTION**

Submitted to "Physics Letters A"


**1989**

The Hamiltonian

$$H = \sum_{\mathbf{k}} \sum_{\sigma} \varepsilon_{\mathbf{k}} c_{\mathbf{k}\sigma}^{\dagger} c_{\mathbf{k}\sigma} + U \sum_i n_{i\uparrow} n_{i\downarrow} \quad (1)$$

with  $c_{i\sigma}^{\dagger}$  and  $c_{\mathbf{k}\sigma}^{\dagger}$  being creation operators of fermions in a Bloch and Wannier state resp., is usually called Hubbard model<sup>1/</sup>, if one assumes  $U$  to be positive, since it is the on-site matrix element of the electron-electron interaction. Otherwise, if  $U$  is thought to describe a kind of effective attraction of electrons like in the simplest version of a model of superconductivity, it can be negative. Unfortunately the two cases, i.e.  $U > 0$  and  $U < 0$  resp., are usually treated differently, due to the differing goals of describing magnetism or superconductivity<sup>2,3/</sup>. In the following it will be shown, that by generalization of the functional integral method outworked mainly for magnetic purposes, it is possible to handle both the attractive and the repulsive version of the model (1) on the same footing.

The functional integral technique starts with rewriting the interaction term of the Hamiltonian as quadratic form. There are several different transformations known within the context of magnetism, which are all operator identities and formal equivalent - till the moment, when approximations are introduced. However, employing a special approximation scheme usually destroys this equivalence yielding different results to the same physical questions. Due to the lack of a criterion which transformation is to prefer the method is often cried arbitrary, since the choice of the quadratic form pre-determines the results. To avoid this disadvantage the authors argued, that dealing with the model (1) one should not restrict the various possibilities of breaking symmetries, what is usually done by adopting a special quadratic form, rather the model should "have the choice" by minimising its thermodynamical potential<sup>4/</sup>.



Working out this idea we start by rewriting the interaction term

$$n_{i\uparrow} n_{i\downarrow} = \frac{1}{4} (1 - \alpha_i) + R_{i\bar{z}} + (\alpha_i - 1) \frac{1}{3} \underline{S}_i^2 + (\alpha_i + 1) \frac{1}{3} \underline{R}_i^2 \quad (2)$$

with  $\underline{S}_i$  being the spin vector

$$S_{ix} = \frac{1}{2} (c_{i\uparrow}^{\dagger} c_{i\downarrow} + c_{i\downarrow}^{\dagger} c_{i\uparrow}); S_{iy} = \frac{1}{2i} (c_{i\uparrow}^{\dagger} c_{i\downarrow} - c_{i\downarrow}^{\dagger} c_{i\uparrow}); S_{iz} = \frac{1}{2} (n_{i\uparrow} - n_{i\downarrow}) \quad (3)$$

and  $\underline{R}_i$  being the quasispin vector

$$R_{ix} = \frac{1}{2} (c_{i\uparrow}^{\dagger} c_{i\downarrow}^{\dagger} + c_{i\downarrow} c_{i\uparrow}); R_{iy} = \frac{1}{2i} (c_{i\uparrow}^{\dagger} c_{i\downarrow}^{\dagger} - c_{i\downarrow} c_{i\uparrow}); R_{iz} = \frac{1}{2} (n_{i\uparrow} + n_{i\downarrow} - 1). \quad (4)$$

This transformation was shown to be a generalization of quite all quadratic forms used in the context of magnetism <sup>/4/</sup>. However, due to the containment of both spins and quasispins in (2) it is possible to look for magnetism ( $\langle \underline{S}_i \rangle \neq 0$ ), charge ordering ( $\langle R_{i\bar{z}} \rangle \neq 0$ ), or superconduction ( $\langle R_{ix} \rangle \neq 0$ ) in a unified manner. Inserting eq. (2) into eq. (1) and applying the time ordering trick the partition sum becomes

$$\mathcal{Z}_\alpha = e^{-\frac{\beta U}{4} \sum_i (1 + \alpha_i)} \text{tr} \left\{ T_\tau e^{-\beta \mathcal{H}_0} e^{-\beta \int_0^1 d\tau H_A(\tau)} \right\} \quad (5)$$

Here was abbreviated

$$\mathcal{H}_0 = \sum_{\underline{k}} \sum_{\sigma} (\epsilon_{\underline{k}} - \mu + \frac{U}{2} - \sigma h_{ex}) n_{\underline{k}\sigma} \quad (6)$$

$$H_A(\tau) = -\frac{U}{3} \sum_i (1 - \alpha_i) \underline{S}_i^2(\tau) + \frac{U}{3} \sum_i (1 + \alpha_i) \underline{R}_i^2(\tau) \quad (7)$$

$$A(\tau) = e^{\tau \beta \mathcal{H}_0} A e^{-\tau \beta \mathcal{H}_0} \quad (8)$$

with  $h_{ex}$  being the external magnetic field applied in  $z$ -direction and measured in energy units. One can immediately see that the chemical potential  $\mu$  which is known to be  $U/2$  when neutrality and electron hole symmetry is assumed, acts as symmetry breaking field with respect to  $R_{i\bar{z}}$  in the same manner as  $h_{ex}$  does with respect to  $S_{iz}$ . The latter may be of interest when (1) is applied to describe substitution effects in magnets or superconductors. Now, the Hubbard-Stratonovic transformation

$$e^{\alpha A^2} = \int_{-\infty}^{+\infty} dx e^{-\pi x^2 - 2\sqrt{\pi\alpha} x A} \quad (9)$$

is used at every lattice site and at each instant of "imaginary time" to rewrite the partition sum as functional integral

$$\mathcal{Z}_\alpha = \int_{-\infty}^{+\infty} \prod_{i=1}^N \mathcal{D}^3 \underline{x}_i(\tau) \mathcal{D}^3 \underline{y}_i(\tau) e^{-\Omega_\alpha[\dots \underline{x}_i(\tau), \dots, \underline{y}_i(\tau), \dots]} \quad (10)$$

with  $\Omega_\alpha$  being the functional

$$\Omega_\alpha = \pi \sum_i \int_0^1 d\tau (\underline{x}_i^2(\tau) + \underline{y}_i^2(\tau)) + \ln Z_\alpha[\dots \underline{x}_i(\tau), \dots, \underline{y}_i(\tau), \dots], \quad (11)$$

where

$$Z_\alpha = \text{tr} \left\{ T_\tau e^{-\beta \mathcal{H}_0} e^{-\sqrt{\frac{4\pi}{3}} \beta U \sum_i \int_0^1 d\tau (\sqrt{1-\alpha_i} \underline{x}_i \underline{S}_i + i \sqrt{1+\alpha_i} \underline{y}_i \underline{R}_i)} \right\} \quad (12)$$

By substituting

$$\underline{X}_i(\tau) = \sqrt{1-\alpha_i} \underline{x}_i(\tau); \quad \underline{Y}_i(\tau) = \sqrt{1+\alpha_i} \underline{y}_i(\tau) \quad (13)$$

one can shift the  $\alpha_i$ -dependence to the Gaussian measure

$$\mathcal{Z}_\alpha = \int_{-\infty}^{+\infty} \prod_{i=1}^N \frac{\mathcal{D}^3 \underline{X}_i(\tau) \mathcal{D}^3 \underline{Y}_i(\tau)}{(1-\alpha_i^2)^{3/2}} e^{-\Omega[\alpha; \dots \underline{X}_i(\tau), \dots, \underline{Y}_i(\tau), \dots]} \quad (14)$$

with

$$\Omega[\alpha; \dots] = \pi \sum_i \int_0^1 d\tau \frac{\underline{X}_i^2(\tau)}{1-\alpha_i} + \frac{\underline{Y}_i^2(\tau)}{1+\alpha_i} - \ln Z[\dots \underline{X}_i(\tau), \dots, \underline{Y}_i(\tau), \dots] \quad (15)$$

$$Z[\dots] = \text{tr} \left\{ T_\tau e^{-\beta \mathcal{H}_0} e^{-\sqrt{\frac{4\pi}{3}} \beta U \sum_i \int_0^1 d\tau (\underline{X}_i(\tau) \underline{S}_i(\tau) + i \underline{Y}_i(\tau) \underline{R}_i(\tau))} \right\} \quad (16)$$

Therefore the functional  $Z$  is no longer explicit dependent on the  $\alpha_i$ 's. Now, one can proceed in the usual way, i.e. applying coupling constant trick, writing down the associated Dyson equation, discuss the different approximation schemes developed to calculate  $Z$ , etc., as done, e.g., in <sup>/3/</sup> for the two field scheme, which is a special case of eq. (2). Of course, the latter is out of the scope of this letter and will be published elsewhere. However, without doing any special calculation one immediately recognizes that for each appro-

ximation the dependence of the  $\alpha_i$  will be different. Therefore the best choice of the  $\alpha_i$  is determined by the approximation employed to calculate the partition sum and the related thermodynamical potential. By minimizing the latter one finds the equation determining the  $\alpha_i$  straightforwardly.

$$-\frac{1}{\beta} \frac{\partial}{\partial \alpha_i} \ln Z = 0 \quad ; \text{ for all } \alpha_i \quad (17)$$

This criterion holds independent of the method used to evaluate the functional integral (14). However, usually the partition sum is calculated by means of the saddle point approximation, i.e. only the extremizing "paths", hereafter assigned as  $X_i^e(\tau)$  and  $Y_i^e(\tau)$  resp., are of interest. By minimizing the functional  $\Omega$  with respect to the fields one finds

$$X_i^e(\tau) = \frac{1-\alpha_i}{2\pi} \frac{\delta}{\delta X_i(\tau)} \ln Z \Big|_{X_i(\tau)=X_i^e(\tau); Y_i(\tau)=Y_i^e(\tau)} \quad (18)$$

$$Y_i^e(\tau) = \frac{1+\alpha_i}{2\pi} \frac{\delta}{\delta Y_i(\tau)} \ln Z \Big|_{X_i(\tau)=X_i^e(\tau); Y_i(\tau)=Y_i^e(\tau)} \quad (19)$$

Within the saddle point approximation the thermodynamical potential becomes a functional of the extremizing paths, which are functions of the  $\alpha_i$  via eqs. (18), (19). One has

$$-\frac{1}{\beta} \ln Z_{\alpha}^{sp} = \frac{U}{4} \sum_i (1+\alpha_i) + \frac{3}{2\beta} \sum_i \ln(1-\alpha_i^2) + \frac{1}{\beta} \Omega[\alpha; \dots X_i^e(\tau); \dots Y_i^e(\tau)] \quad (20)$$

From eq. (17) one finds

$$0 = \frac{1}{\beta} \frac{\partial \Omega}{\partial \alpha_i} + \frac{1}{\beta} \left( \frac{\delta \Omega}{\delta X_i^e} \right) \frac{dX_i^e}{d\alpha_i} + \frac{1}{\beta} \left( \frac{\delta \Omega}{\delta Y_i^e} \right) \frac{dY_i^e}{d\alpha_i} - \frac{3\alpha_i}{\beta(1-\alpha_i^2)} + \frac{U}{4} \quad (21)$$

The second and the third term in eq.(21) vanish, due to the saddle point approximation. Since  $Z$  depends not explicitly on  $\alpha_i$  the resulting expression looks like

$$0 = \frac{\beta U}{4} - \frac{3\alpha_i}{1-\alpha_i^2} + \frac{\pi}{(1-\alpha_i)^2} \int_0^1 d\tau X_i^{e2}(\tau) - \frac{\pi}{(1+\alpha_i)^2} \int_0^1 d\tau Y_i^{e2}(\tau) \quad (22)$$

The fields can be expressed with the help of mean values of the local spins and quasispins. From variation of the functional  $\ln Z$  in eqs. (18) and (19) one finds

$$X_i^e(\tau) = -\frac{1-\alpha_i}{2\pi} \sqrt{\frac{4\pi}{3} \beta U} \langle S_i(\tau) \rangle_e \quad (23)$$

$$Y_i^e(\tau) = -i \frac{1+\alpha_i}{2\pi} \sqrt{\frac{4\pi}{3} \beta U} \langle R_i(\tau) \rangle_e \quad (24)$$

Here was introduced

$$\langle A(\tau) \rangle_e = \frac{\text{tr} \{ T_{\tau} \rho_e A(\tau) \}}{\text{tr} \{ T_{\tau} \rho_e \}} \quad (25)$$

and

$$\rho_e = e^{-\beta \mathcal{H}_0} e^{-\sqrt{\frac{4\pi}{3} \beta U} \sum_i \int_0^1 d\tau (X_i^e(\tau) S_i(\tau) + Y_i^e(\tau) R_i(\tau))} \quad (26)$$

Inserting eqs. (23) and (24) into eq. (22) yields

$$1-\alpha_i^2 = \frac{\beta U}{12} \left( 1 + \frac{4}{3} \int_0^1 d\tau \langle S_i(\tau) \rangle_e^2 + \frac{4}{3} \int_0^1 d\tau \langle R_i(\tau) \rangle_e^2 \right) =: B \quad (27)$$

which gives the two solutions

$$\alpha_{i,1,2} = -\frac{1}{2B} (1 \pm \sqrt{1+4B^2}) \quad (28)$$

The second term in eq. (20) demands

$$1-\alpha_i^2 = \frac{\alpha_i}{B} > 0 \quad (29)$$

This selects

$$\alpha_i = -\frac{1}{2B} (1 - \sqrt{1+4B^2}) \quad (30)$$

By eq. (30) the  $\alpha_i$  are completely determined and therefore, no "arbitrariness" remains. However, since the  $\alpha_i$  depend on the mean values of the local spins and quasispins, i.e. the quadratic form itself depends on the results, which should be calculated

from it, one has to solve a self consistent problem. The remaining task is to determine the functional  $Z$  in a more or less advanced approximation scheme. This is beyond the scope of this letter and will be published elsewhere.

The idea of the method presented above can also be applied if the functional integration is carried out by expansion around the extremizing paths to the second order. Furthermore, what said above is independent of the special kind of  $\mathcal{H}_0$ , since it is related to the Hubbard interaction term only, and therefore other models, e.g. the (periodic) Anderson model, may be treated in complete analogy.

The author wants to thank E.Heiner for discussion and critical remarks.

#### References

1. J.Hubbard, Proc. R.Soc.London, A276 (1963) 238.
2. e.g. A.V.Svidsinskij, Prostranstvenno-Njeodnorodnije Sadaci Teorii Sverkhprovodimosti, Nauka, Moscow, 1982.
3. T.Moriya, Spin Fluctuations in Itinerant Electron Magnetism, Springer, Berlin, 1985.
4. R.Schumann and E.Heiner, in: P.Ziesche (ed): Proc. 18th Symp. Electronic Structure, TU Dresden, 1988, p. 22; Physics Lett. A, 134 (1988) 202.
5. R.L.Stratonovich, Dokl.Akad.Nauk SSSR, 115 (1957) 1094.  
J.Hubbard, Phys.Rev.Lett., 3 (1959) 77.

Received by Publishing Department  
on February 17, 1989.