

СООБЩЕНИЯ Объединенного института ядерных исследований дубна

H41

E17-89-654

# R.Hayn\*

THE EXACT INSTANTON SOLUTION FOR THE CONDUCTIVITY OF A DISORDERED SOLID

\*On leave from: Technical University of Dresden, GDR

1989

## 1.Introduction

One can study the physical properties of disordered systems for Fermi-energies deep in the localized tail in a systematic way. This was done first for the density of states by Lifshitz [1]. Mott and Davis [2] have calculated the frequency-dependent conductivity  $\sigma(\omega)$  in the same limit. The conductivity is then determined by the probability of finding two deep potential wells with eigenvalues at  $E\pm\omega$ . Because of the repulsion of energy levels by tunneling these wells must be separated by a minimal distance proportional to  $ln \ \omega$ . In the one-dimensional case, to which we will restrict our considerations for simplicity, this implies that the conductivity will be proportional to  $\omega^2 ln^2 \omega$  [2]. These qualitative arguments can be made more explicit by calculating the semiclassical limit of a certain functional integral. The interesting results come from nontrivial saddle-points and cannot be reached by ordinary perturbation theory. For the density of states this has been done already in a number of papers [3-6]. However, to the authors knowledge, there exists only one analogous calculation of the conductivity [7]. In the present work we start from a supersymmetric formulation of the problem. As a first step we only present the exact solution of the saddle-point equation. The calculation of the fluctuations around it shall be published in a forthcoming publication.

The saddle-point equation corresponds to the classical motion of a particle in a two-dimensional nonlinear and

a and the second s

Service and the service of the servi

$$S_{2}(x) = \frac{1}{2\pi^{2}} \operatorname{Re} \left( \mathfrak{F}^{+}(x) - \mathfrak{F}^{-}(x) \right)$$
 (3)

where

$$\mathfrak{S}^{\alpha\beta}(\mathbf{x}) = \langle \mathbf{G}(\mathbf{E} - \frac{\omega}{2} + \alpha \mathbf{i}0^{+}; \mathbf{0}, \mathbf{x}) \mathbf{G}(\mathbf{E} + \frac{\omega}{2} + \beta \mathbf{i}0^{+}; \mathbf{x}, \mathbf{0}) \rangle . (4)$$

Next we write the Greens functions as functional integrals over commuting fields  $\phi_i$  and anticommuting fields  $\chi_i$  and carry out the average. This results in a supersymmetric field theory [15]. The product of advanced and retarded Greens functions for instance, is given by

$$\mathfrak{F}^{-+}(\mathbf{x}) = \frac{|\mathbf{E}|^2}{16\gamma^2} \int \mathcal{D}\{\Psi_1, \Psi_1^*\} \ \chi_1(0)\chi_1^*(\tilde{\mathbf{x}})\chi_2(\tilde{\mathbf{x}})\chi_2^*(0) \ e^{-S} \tag{5}$$

with  $\tilde{x}=x\sqrt{|E|}$  and the action

$$S = -\frac{1}{4} \varepsilon^{3/2} i \int dx \left\{ \left| \frac{d\Psi}{dx} \right|^{2} - \left| \frac{d\Psi}{dx} \right|^{2} + (1+\bar{\omega}) \left| \Psi_{1} \right|^{2} - (1-\bar{\omega}) \left| \Psi_{2} \right|^{2} + \frac{1}{4} \left( \left| \Psi_{1} \right|^{2} - \left| \Psi_{2} \right|^{2} \right)^{2} \right\} , \qquad (6)$$

where  $\Psi_i$  is a short-hand notation for  $(\phi_1, \phi_1, \chi_1, \chi_1)$ . In (6) we already introduced by a simple rescaling the dimensionless parameters of the theory, namely  $\varepsilon = |\mathbf{E}|/\gamma^{2/3}$ and  $\overline{\omega} = \omega/2|\mathbf{E}|$ . The relevant saddle-point of (6) lies in the complex plane. We go from  $\Psi_1 = 0$  in the direction of steepest descent by rotating the real and imaginary part of  $\Psi_1$  such, that  $|\Psi_1|^2 \rightarrow \mathbf{i}|\Psi_1|^2$  and  $|\Psi_2|^2 \rightarrow -\mathbf{i}|\Psi_2|^2$ . Thus we obtain a real action:

### 2.The saddle-point equation

We consider the model of a free electron in a white-noisepotential V(x) given by the Hamiltonian

$$H = -\frac{d^{2}}{dx^{2}} + V(x) , \quad \langle V(x)V(x') \rangle = 2\gamma \delta(x-x')$$
(1)

setting  $h/2\pi=2m=1$ . In general the dimension does not play a crucial role in the instanton approach [3-7]. We concentrate therefore on one dimension and express the conductivity  $\sigma$  with the help of a correlation function  $S_2(x)$  ([7]):

Re 
$$\sigma(\omega) = -\frac{e^2 \omega^2 \pi}{2} \int dx x^2 S_2(x)$$

(2)

The correlation function can be expressed in terms of averaged Greens functions:

2

$$S = \frac{1}{4} \varepsilon^{3/2} \int d\mathbf{x} \left( \left| \frac{d\Psi}{dx}_{1} \right|^{2} + \left| \frac{d\Psi}{dx}_{2} \right|^{2} + (1 + \bar{\omega}) \left| \Psi_{1} \right|^{2} + (1 - \bar{\omega}) \left| \Psi_{2} \right|^{2} - \frac{1}{4} \left( \left| \Psi_{1} \right|^{2} + \left| \Psi_{2} \right|^{2} \right)^{2} \right) \qquad (7)$$

In deriving the saddle-point equation we may neglect the anticommuting fields. Furthermore we consider only the real part of  $\phi_1$  as the action (7) is invariant with respect to rotations in the complex  $\phi_1$ - and  $\phi_2$ -plane separately. The saddle-point condition  $\delta S=0$  gives than the classical problem of motion in a wine bottle like potential

$$\{\frac{d^{2}}{dx^{2}} + \frac{1}{2}\vec{\phi}^{2} - 1 + \vec{\omega}\} \phi_{1/2} = 0 , \qquad (8)$$

known as the Garnier-problem [8,9,12]. The instanton solutions of (8) are the only one with finite action. They approach  $\vec{\phi}=0$  at the boundaries  $x \rightarrow \pm \infty$  and therefore must have zero energy. There exists a second constant of motion

$$K = \frac{1}{4} \left( \phi_1 p_2 - \phi_2 p_1 \right)^2 + \overline{\omega} \left( p_1^2 + \frac{\phi_1^4}{4} - p_2^2 - \frac{\phi_2^4}{4} - (1 + \overline{\omega}) \phi_1^2 - (1 - \overline{\omega}) \phi_2^2 \right)$$

with  $p_i = d\phi_i/dx$ , which has been found in [9]. It also must be zero for the instanton solutions, because of their boundary conditions. A complete set of instanton solutions depends therefore on two parameters.

A mathematical investigation of equations like (8) can be found in [9,11]. Here we look for the solution with finite action relevant to the conductivity problem. The usual arguments about two nearly degenerate potential wells lead us to expect that it looks like a two instanton solution. A first indication about it gives the approximate consideration in the next section.

# 3.Approximate solution

In close analogy to [2,7] we start with two isolated one instanton solutions:

$$g_{1/2} = \frac{2\sqrt{1\pm\eta}}{\cosh(\sqrt{1\pm\eta}(x\pm D/2))}$$
(9)

2

of the equations

11

$$\left(-\frac{d^{2}}{dx^{2}}+V_{1/2}\right) g_{1/2} = \left(-1+\eta\right) g_{1/2}, V_{1} = \frac{g_{1}}{2}.$$
(10)

We have written the instantons in this form in order to show the analogy with the quantum mechanical eigenvalues of a potential well. They are solutions of our problem (8) in the limit of an infinite separation D when  $\eta = \tilde{\omega}$ . To obtain a solution for a finite distance D we apply ordinary perturbation theory of two nearly degenerate eigenstates [16]. Due to the overlap matrix element t=4 exp(-D) the energy splitting increases. If we set it equal to  $\tilde{\omega}$ :

$$\bar{\omega} = \sqrt{\eta^2 + t^2}$$
(11)

we get an approximate solution of (8):

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} g_1 + \begin{pmatrix} \sin \theta \\ -\cos \theta \end{pmatrix} g_2 \quad .$$
 (12)

5

The angle  $\theta$  is determined by

(13)One can see from (11), that there exists a minimal distance equal  $ln 4/\tilde{\omega}$  where both potential wells have the same depth and the angle  $\theta$  equals  $\pi/4$ . It will be seen later, that the exact instanton-solution has the same gualitative behaviour as (12). But to calculate the fluctuations around the saddle point of (7) we need the exact instanton solution.

 $t = \tilde{\omega} \sin 2\theta$ ,  $n = \tilde{\omega} \cos 2\theta$ 

#### 4. The exact instanton solution

As already mentioned in the introduction, we have different possibilities to find an exact solution of (8). There exist a lot of similarities to the nonlinear Schrödinger equation for which the N-soliton solutions are given in [13.14]. We follow this very direct way to find the two instanton solution of (8).

We interpret  $-\vec{\phi}^2/2$  as potential V(x) and investigate the solutions of

$$L_{k} \phi(x,k) = \left(-\frac{d^{2}}{dx^{2}} + V(x) - k^{2}\right) \phi(x,k) = 0 \quad (14)$$

in dependence on k. It is known from [13] that the two instanton can be found with the ansatz

$$\phi(\mathbf{x},\mathbf{k}) = (\mathbf{k}^2 + \mathbf{a}_1(\mathbf{x})\mathbf{k} + \mathbf{a}_2(\mathbf{x})) e^{\mathbf{l}\mathbf{k}\mathbf{x}}$$
(15)

Comparing (14) with (8) we see, that we look for a potential V with two eigenvalues at the following values of k:

$$\kappa_{1/2} = i\sqrt{1\pm\bar{\omega}} = 2iu_{1/2} \qquad (16)$$

Because of  $L_{\kappa} = L_{\kappa}$  the relations

$$\phi(\mathbf{x}, \kappa_{1}^{\bullet}) + \alpha_{1} \phi(\mathbf{x}, \kappa_{1}) = 0 ; i=1,2$$
 (17)

must be fulfilled with some free parameters  $\alpha_i$ . On the contrary, already these  $\alpha_i$  determine uniquely the functions a (x) in (15). By inserting (15) in (14) the potential is given by V=2ida,/dx.

To fulfill the selfconsistency condition  $V=-\phi^2/2$ , it is very convenient [13] to define

$$\psi(\mathbf{x},\mathbf{k}) = \frac{\phi(\mathbf{x},\mathbf{k})}{\mathbf{R}(\mathbf{k})} = (1 + \sum_{n=1}^{2} \frac{\psi_{n}}{\mathbf{k} - \kappa_{n}} e^{-i\kappa_{n}\mathbf{x}} ) e^{i\mathbf{k}\mathbf{x}}$$
(18)

with  $R(k) = (k-\kappa_1)(k-\kappa_2)$ . The residua  $\psi_1$  of  $\psi(x,k)$  at  $\kappa_2$  are eigenfunctions of (14) at  $1\pm \overline{\omega}$ . They are determined by (17) which is now written as

$$\sum_{i=1}^{2} \left( \frac{e^{i(\kappa_{n}^{*}-\kappa_{1})x}}{\kappa_{n}^{*}-\kappa_{1}} + c_{i}\delta_{1n} \right) \psi_{1} = -e^{i\kappa_{n}^{*}}$$
(19)

with the new free parameters  $c = \alpha_{\rm c} R'(\kappa_{\rm c}) / R(\kappa_{\rm c})$  . Now we consider the function

$$\Omega(\mathbf{x},\mathbf{k}) = \psi(\mathbf{x},\mathbf{k})\psi^{\dagger}(\mathbf{x},\mathbf{k}^{\dagger})$$

The residuum at  $k \infty$  is equal to  $a_1 + a_1^{\dagger}$  and must vanish for real V. Comparing it with all the other residua gives the condition  $c_1 = -c_1^{\bullet}$ . By inserting an expansion of  $\psi(x,k)$  in powers of 1/k into (14), we find that the residuum of  $k\Omega(x,k)$  at  $k \infty$  is equal to V(x)/2. Once again we use the

fact that the sum over all residua of  $k\Omega$  vanishes and find

 $2\psi_{\mathbf{b}} = \phi_{\mathbf{b}}$ 

$$V(x)/2 = -\sum_{i=1}^{2} \psi_{i}^{*} \psi_{i} c_{i} (\kappa_{i}^{*} - \kappa_{i})$$
(20)

Now, we can fulfil the condition  $V=-\vec{\phi}^2/2$  if we set

and

$$c_{1} = |b_{1}|^{2} / (\kappa_{1}^{*} - \kappa_{1})$$

(21)

(22)

The general solution of (19,21,22) containes two parameters  $b_i$  as it must be. It represents the exact solution of (8):

$$\phi_{1/2}(x) = 8u_{1/2} \frac{e^{f_{1/2} \pm e^{2f_{2/1} + f_{1/2} - D_0}}}{1 + e^{2f_1} + e^{2f_2} + 4e^{2(f_1 + f_2 - D_0)}}, \qquad (23)$$

where we have introduced

$$f_i = 2u_i x - \ln|b_i|$$
 (24)

and the minimal distance  $D_{n}$  between the solitons, defined by

$$2e^{-b} = (u_1 - u_2) / (u_1 + u_2)$$
 (25)

The two degrees of freedom  $b_i$  correspond to a general shift of the x-coordinate which we will keep fixed in the following and to the distance between the instantons. To make the result more transparent and to show the simi-

larity with the approximate solution, we introduce instead of  $f_i$  the new parameters  $x_i$ :

$$x_{1} = f_{1} + \ln \sqrt{1 + exp(-2\Delta f(x))} = s(x) + D(x)/2$$

$$x_{2} = f_{2} - D_{0} + \ln 2 - \ln \sqrt{1 + exp(-2\Delta f(x))} = s(x) - D(x)/2$$
(26)

with  $\Delta f(x) = f_1 - f_2$ . We choose the  $b_1$  in (24) such that

$$\Delta f(x) = 2(u_1 - u_2)x + x_0$$
 (27)

and

$$s(x) = (u_1 + u_2)x$$
, (28)

....

where the new parameter  $x_0$  determines the spacing between the instantons. By introducing the angle  $\theta(x)$ 

$$\sin \theta(\mathbf{x}) = \frac{\exp(-\Delta f(\mathbf{x}))}{\sqrt{1 + \exp(-2\Delta f(\mathbf{x}))}}$$
(29)

and with the help of (26) we rewrite (23) as follows

$$\phi_{1}(x) = 8u_{1} \frac{\cos \theta(x)e^{-x_{1}} + \sin \theta(x)e^{x_{2}}}{1 + e^{-2x_{1}} + e^{2x_{2}}}$$

$$\phi_{2}(x) = 8u_{2} \frac{\sin \theta(x)e^{-x_{1}} - \cos \theta(x)e^{x_{2}}}{1 + e^{-2x_{1}} + e^{2x_{2}}}$$
(30)

It is interesting that the distance D(x), introduced in (25), depends on the angle  $\theta(x)$  in the same way as in the approximate solution:

$$exp(-D(x)) = exp(-D_{o}) \sin 2\theta(x)$$
<sup>(31)</sup>

The formulas (32) represent the final result of this paper, where the parameters  $x_1$ ,  $x_2$  and  $\theta$  are determined by subsequently inserting the expressions (16),(25-29) and (31). These form a family of zero-energy solutions parametrized by  $x_0$  corresponding to varying distances between the instantons. These solutions are similar in structure to the approximate solutions found in Sect.3. The only difference is that the angle  $\theta$  and the distance D now depend slightly on x, and the  $g_1$  have changed a little bit. One finds (12) to be a good approximation for small  $\bar{\omega}$  and

8

one has the advantage of a systematic way of improving it. One can see that the minimal distance between the instantons which is so important for the conductivity comes in very naturally without any additional constraint as in [7]. The exact solution (30) seems to be a good starting point to calculate the fluctuations around the extremum and to give an expression for the conductivity. This will be done in a subsequent publication.

The author is very indebted to Prof. John for many inspiring discussions and to Drs. Inosemzev and Kostov from JINR Dubna, who supplied him with some references to the mathematical literature on this problem.

#### <u>References</u>

- [1] Lifshitz, I.M., 1964, Adv. Phys. 13, 483
- [2] Mott, N.F., Davis, E.A., 1971, Electronic Processes in Non-crystalline Materials, Clarendon, Oxford, Sec.2.4
- [3] Cardy, J.L., 1978, J. Phys. <u>C11</u>, L321
- [4] Houghton, A., Schäfer, L.J., 1979, J. Phys. A12, 1309
- [5] Luttinger, J.M., Tao, R., 1983, Ann. Phys. <u>145</u>, 185
- [6] Luttinger, J.M., Waxler, R., 1987, Ann. Phys. 175, 319
- [7] Houghton, A., Schäfer, L.J., Wegner, F.J., 1980, Phys. Rev. <u>B22</u>, 3598
- [8] Garnier, R., 1919, Rend. Circ. Mat. Palermo 43, 155
- [9] Wojciechowski, S., 1984, Lett. Nuovo. Cim. 41, 361
- [10] Guilarte, J.M., 1988, Ann. Phys. 188, 307
- [11] Manakov, S.W., 1973, Zh. Eksp. Teor. Fiz. 65, 505

- [12] Kostov, N.A., 1989, Lett. Math. Phys. 17, 95
- [13] Dubrovin, B.A., Malanyuk, T.M., Krichever, I.M., Makhankov, V.G., 1988, Fis.el.častic i at. jadra <u>19</u>, 579
- [14] Nogami, Y., Warke, C.S., 1976, Phys. Lett. 59A, 251
- [15] Efetov, K.B., 1983, Adv. Phys. <u>32</u>, 53
- [16] Landau, L.D., Lifshitz, E.M., 1979, Lehrbuch der Theoretischen Physik, Band 3, Quantenmechanik, Berlin, Akademie-V.

Received by Publishing Department on September 14, 1989.

# Хайн Р.

E17-89-654

Точное инстантонное решение в задаче

о проводимости в неупорядоченном твердом теле

Методом функционального интегрирования с бозе- и фермиполями получено уравнение для седловой точки. Точное решение этого нелинейного уравнения получено с помощью метода, близкого методу обратной задачи рассеяния.

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Сообщение Объединенного института ядерных исследований. Дубна 1989

E17-89-654

The Exact Instanton Solution for the Conductivity of a Disordered Solid

Starting from a functional integral with Bose- and Fermi fields we derive the saddle-point-equation in the semiclassical limit. The exact solution of this nonlinear classical problem can be found, using a method similar to the inverse scattering method.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Communication of the Joint Institute for Nuclear Research. Dubna 1989

# SUBJECT CATEGORIES OF THE JINR PUBLICATIONS

Index	Subject	
1. High (	energy experimental physics	
2. High (	energy theoretical physics	
3. Low en	nergy experimental physics	Сообщение Об
4. Low e	nergy theoretical physics	
5. Mathe	matics	
6. Nuclea	ar spectroscopy and radiochemistry	
7. Heavy	ion physics	Hayn R.
8. Cryog	enics	The Exact In
9. Accel	erators	for the cond
10. Autom	atization of data processing	mi fields we
11. Compu	ting mathematics and technique	miclassical
12. Chemi	stry	classical pr
13. Exper	imental techniques and methods	the inverse
14. Solid	state physics. Liquids	The inves
15. Exper at lo	imental physics of nuclear reactions w energies	of mediecie
16. Healt	h physics. Shieldings	-
17. Theor	y of condenced matter	
18. Appli	ed researches	
19. Bioph	ysics	Communic