


89-813



СООБЩЕНИЯ  
ОБЪЕДИНЕННОГО  
ИНСТИТУТА  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ  
ДУБНА

M 75

E17-89-813

T.M.Mishonov<sup>1</sup>, B.K.Ghosh<sup>2</sup>

ACOUSTIC WAVES EMISSION  
BY TWO DIMENSIONAL PLASMA WAVES

---

<sup>1</sup> Permanent address: Theoretical Physics Division,  
University of Sofia, 5A. Ivanov Blvd., Sofia 1126,  
Bulgaria

<sup>2</sup> Department of Physics, Allahabad University,  
Allahabad 211002, India

1989

## 1. Introduction

The 2D plasmon is a well known excitation of 2DEG in Si inversion layers and  $\text{GaAs-AL Ga}_x\text{As}_{1-x}$  heterojunctions (see the review by Ando et al [1]). Recently there has been a considerable interest in the study of light scattering from free carriers in multiple quantum wells, heterostructures and heterojunction superlattices [2-5]. Investigations of the interaction between 2DEG and acoustic phonons is now another rich field of activity. A general review in this area is given by Challis et al [6].

The aim of the present paper is to investigate theoretically the phonon channel of the decay of plasmons excited by an externally applied e.m. field. This effect may be useful for the coherent phonon conversion of e.m. waves into acoustic waves. This method for coherent phonon generation was for the first time proposed by Krasheninnikov, Sultanov and Chaplik (KSC) [7] (c.f. [8]). KSC considered only the low frequency case for which the phonon wavelength  $\lambda_{ph}$  is larger than the thickness of the electron layer  $d$ . KSC concluded, that although this coherent conversion has a very low efficiency, one can observe it experimentally. Description of appropriate experimental techniques for phonon detection is given in [9,10].

In the following section 2, a brief description of the model structure is outlined for the present analysis. Section 3 present a formal theory of plasma waves. The acoustic emission is described in section 4, where an expression for the power conversion coefficient (PCC) is also obtained. Section 5 contains results and a brief discussion. Finally the main conclusions are also given in this section.

## 2. Model


We consider here a simplified model with only one filled 2D subzone. Fig.1 is shows the structure similar to the one described by Allen et al [11] and Batke et al [12]. In the GaAs layer the electrons move freely with an effective mass  $m$ . But in the z-direction they are confined by the large potential barrier of  $\text{Al}_x\text{Ga}_{1-x}\text{As}$ . The electron wave function is approximated in the flat-band model [13] by

$$\psi_0(z) = \begin{cases} (2/d)^{1/2} \cos(\pi z/d) & \text{for } -d/2 < z < d/2 \\ 0 & \text{for } |z| > d/2. \end{cases} \quad (1)$$

The instantaneous volume density of electrons is

$$n(r, t) = n(\rho, t) n_0(z), \quad (2)$$

where:  $n_0(z) = |\psi_0(z)|^2$ ,  $r \equiv (\rho, z)$ ,  $\rho \equiv (x, y)$ .



$$n(\rho, t) = n + \delta n(\rho, t). \quad (3)$$

Here  $n(\rho, t)$  is the instantaneous electronic density per unit area,  $n$  is the equilibrium electron number per unit area and  $\delta n(\rho, t)$  is the excess electron number per unit area induced by the external e.m. field. For simplification, we shall take the background dielectric constant  $\epsilon$  to be identical for GaAs and  $\text{Al}_{1-x}\text{Ga}_x\text{As}$ .




Figure 1. Transformation of the electromagnetic field into sound (schematically; not to scale). The e.m. field  $E_\omega$  receives momentum  $(2\pi\hbar/b)$  from the Al grating and excites plasmons which emit phonons [8].

### 3. 2D plasmons

We briefly describe below the dispersion relation and the decay of plasmons excited in 2DEG. The plasma waves are a self-consistent motion of electrons in which the fluctuation of electronic density creates variation of the electric potential, and the variation of the potential causes variation of the electronic density.

Within the linear response regime, the plane wave potential

$$\delta\phi(\rho, t) = \phi_{q, \omega} \exp i(\mathbf{q} \cdot \rho - \omega t) \quad (4)$$

induces plane wave variation of the electron density

$$\delta n(\rho, t) = n_{q, \omega} \exp i(\mathbf{q} \cdot \rho - \omega t), \quad (5)$$

where  $\mathbf{q} = (q_x, q_y)$  is the 2D wave vector,  $\rho = (x, y)$  is the 2D position vector,  $\omega$  is the frequency and  $t$  is the time. The Fourier coefficients of the electric potential and electron area density are related as

$$n_{q, \omega} = \Pi (e\phi_{q, \omega}) \quad (6)$$

where  $\Pi$  is the polarizability of 2DEG and  $e$  is the electron charge. The electron drift velocity  $\mathbf{v} = (v_x, v_y)$  is also a plane wave

$$\mathbf{v}(\rho, t) = \mathbf{v}_{q, \omega} \exp i(\mathbf{q} \cdot \rho - \omega t). \quad (7)$$

The conservation law

$$\partial \delta n(\rho, t) / \partial t = -n(\partial v_x / \partial x + \partial v_y / \partial y) \quad (8)$$

gives the relation

$$n_{q, \omega} = n \mathbf{q} \cdot \mathbf{v}_{q, \omega} / \omega. \quad (9)$$

This relation and the formula for the electric current  $\mathbf{j} = nev$  give the connection between  $\Pi$  and the conductivity tensor  $\underline{\sigma}$  (see for discussion [8])

$$\Pi = -i \mathbf{q} \cdot \underline{\sigma} \cdot \mathbf{q} / e^2 \omega. \quad (9)$$

Drude formula for conductivity

$$\sigma = \sigma_0 / (1 - i\omega\tau), \quad \sigma_0 = e^2 n \tau / m \quad (10)$$

goes to the Drude formula for the polarizability

$$\Pi = (nq^2 / m\omega^2) / (1 + i/\omega\tau). \quad (11)$$

Here  $\tau$  is relaxation time.

The Coulomb interaction between electrons renormalises the conductivity and polarizability as

$$\Pi^{ren} = \Pi / \kappa, \quad \sigma^{ren} = \sigma / \kappa, \quad (12)$$

where

$$\kappa \equiv 1 - U\Pi. \quad (13)$$

Here

$$U = 2\pi e^2 / \epsilon q, \quad (14)$$

is the 2D Fourier-transformation of the Coulomb potential  $e^2 / \epsilon |\rho|$ .

One can obtain plasmon dispersion relation under wave propagation condition  $\omega\tau \gg 1$  by solving the dispersion relation

$$\kappa(q, \omega) = 0. \quad (15)$$

Explicitly, this gives

$$\omega_{pl}(q) = \left( 2\pi e^2 n q / \epsilon m \right)^{1/2} - i/2\tau. \quad (16)$$

Here  $\omega_{pl}$  is the plasmon frequency and  $q = 2\pi/b$  (see fig.1) is the corresponding wave vector. Period of grating  $b$  determines the plasmon wavelength  $\lambda_{pl} = b \gg d$ .

The absorption coefficient  $A^*$  is defined as ratio of Ohmic dissipation power  $w_\sigma$  and the power of incident e.m. wave  $w_{em}$ :

$$A^* = w_{\sigma} / w_{em}, \quad (17)$$

$$w_{\sigma} = |E_{q\omega}|^2 \operatorname{Re} \sigma / 2,$$

$$w_{em} = c |E_{q\omega}|^2 / 8\pi.$$

Here  $E_{q\omega}$  is the amplitude of the e.m. wave, and the light velocity is denoted by  $c$ .

For the resonance frequency, i.e.  $\omega = \operatorname{Re} \omega_{pi}$  for the absorption coefficient we obtain

$$A^* = \begin{cases} 4\pi(\sigma_0 / c) & \text{In the Gaussian units} \\ (4\pi \cdot 10^{-7})c\sigma_0 & \text{In SI units} \end{cases} \quad (18)$$

this formula is applicable for  $A^* \ll 1$ , but for estimation of order of the effect, one can use it even if  $A^* \approx 1$ . For a complete investigation it is necessary to solve an electrodynamic problem for diffraction and refraction for the real grating structure. We suppose that  $A$  is of an order of relative change of the transmission coefficient [11], i.e.  $A^* \approx 10^{-2}$ .

#### 4. Acoustic emission

The variation of electron density caused by plasmons interacts with the lattice. This has been taken into account through a phenomenological deformation potential within the jellium model.

In the jellium model the Lagrangian density  $\mathcal{L}$  for the lattice motion including electron phonon interaction is given by

$$\mathcal{L} = \left[ \rho_0 \dot{u}^2 - K(\nabla \cdot u)^2 - \Xi n(r, t) \nabla \cdot u \right] / 2. \quad (19)$$

Here  $\Xi$  is the deformation potential,  $K$  is the elastic constant,  $\rho_0$  and  $u$  are the mass density and lattice displacement vector, respectively.

Using the principle of action, one obtains the equation of motion

$$\left[ \nabla^2 - (1/s^2) \partial^2 / \partial t^2 \right] u = (\Xi/K) \nabla n(r, t), \quad (20)$$

where  $s = (K/\rho_0)^{1/2}$  is the sound velocity.

For the electron density variation

$$\delta n(r, t) = n_{q, \omega} n_0(z) \cos(q \cdot \rho - \omega t) \quad (21)$$

the solution of the equation of motion is

$$u_z(r, t) = (\Xi/2K) n_{q, \omega} |\chi(\beta d)| \cos(q \cdot \rho + \beta |z| - \omega t), \quad (22)$$

$$u_x(r, t) = u_y(r, t) = 0,$$

where  $\beta = \omega/s$  is the wave vector of the acoustic phonon and  $\lambda_{ph} = 2\pi/\beta$  is the phonon wavelength. One can verify that the displacement  $u$  of the

lattice is practically normal to the 2DEG (x-y) layer for  $\omega/q \gg s$ ,  $\chi(\beta d)$  is the form-factor of electron density distribution in the z-direction perpendicular to the layer

$$\chi(\beta d) \equiv \int_{-\infty}^{+\infty} n_0(z) \exp(i\beta z) dz. \quad (23)$$

In the flat-band model (1) this is simplified to [14]

$$\chi(\gamma) = [\sin(\gamma)/\gamma] / [1 - (\gamma/\pi)^2], \quad (24)$$

where  $\gamma = \beta d/2 = \pi d/\lambda_{ph}$ .

The power per unit area of the emitted acoustic wave is given by

$$w_{ac} = 2 \langle \rho_0 \dot{u}^2 \rangle s, \quad (25)$$

where  $\langle \rangle$  denotes time-average. Substitution of (22) here gives

$$w_{ac} = (1/4) \rho_0 (\Xi/K)^2 |n_{q, \omega}|^2 |\chi(\beta d)|^2 \omega^2 s. \quad (26)$$

Here, a multiplication factor of 2 is included, which takes into account the two sides of the layer. Let us compare the acoustic power  $w_{ac}$  with ohmic dissipation per unit area  $w_{\sigma}$ . The ohmic power is given by

$$w_{\sigma} = \langle n F \cdot v \rangle, \quad (27)$$

where  $F = -(m/\tau)v$  is the friction force per electron, and  $v(\rho, t)$  is the drift velocity. Substituting here (9) and (21) we obtain for the final expression

$$w_{\sigma} = (m/\tau)(n/2)(\omega/qn)^2 |n_{q, \omega}|^2. \quad (28)$$

The power ratio  $P^*$  defined as

$$P^* = w_{ac} / w_{\sigma}, \quad (29)$$

is roughly speaking (for  $P^* \ll 1$ ) the probability of phonon decay of 2D plasmon. Formulae (26) and (28) give the final result

$$P^* = P_0 |\chi(\beta d)|^2, \quad P_0 = \tau n (\Xi/q)^2 / 2 s^3 \rho_0 m. \quad (30)$$

The power conversion coefficient (PCC)  $P^*$  of e.m. energy into acoustic wave is

$$C^* = A^* P^*. \quad (31)$$

(Probability of phonon generation is the probability of phonon absorption by the plasmon multiplied by phonon decay probability of the plasmon.)

#### 5. Results, discussion and conclusions

Let us take for estimation typical parameters for GaAs 2DEG (see [5]):

$$\begin{aligned}
n &= 7 \times 10^{11} \text{ cm}^{-2}, & b &= 1 \mu, & \mu &= \sigma_0 / ne = 5 \text{ m}^2 / \text{Vs}, \\
\lambda_{pl}^{-1} &= 10000 \text{ cm}^{-1}, & \epsilon &= 12.5, & \Xi &= 16 \text{ eV}, & d &= 260 \text{ \AA} \\
\rho_0 &= 5.3 \text{ g/cm}^3, & s &= 5 \text{ km/s}, & m &= 0.065 m_e, & & (32)
\end{aligned}$$

where  $m_e$  is the free electron mass. For these parameters we obtain:

$$\begin{aligned}
\tau &= 1.84 \times 10^{-12} \text{ s}, & \hbar/2\tau &= 0.179 \text{ meV}, & \omega_{pl} &= 6.1 \text{ meV}, \\
\omega\tau &= 17, & v_{pl} &\equiv \omega/q = 1480 \text{ km/s}, & Ma &\equiv v_{pl}/s = 300, \\
P_0 &= 2.14 \times 10^{-2}, & A^* &\approx 10^{-2}, & \lambda_{ph} &= 35 \text{ \AA}, \\
|\chi|^2 &\approx 10^{-5}, & \gamma &= 23, & C^* &\approx 10^{-8}. & & (33)
\end{aligned}$$

This estimation for PCC is in agreement with KSC.

In order to increase the PCC it is necessary to use layers with higher mobility ( $P_0 \propto \tau \propto \mu \propto \sigma_0$  (30)). There is remarkable success in this direction [15], GaAs structures with mobility of  $\mu = 5 \times 10^6 \text{ cm}^2/\text{V s}$  [16]. However the most important ingredient for PCC increasing is the 2DEG form-factor (24) (see also [6])

$$|\chi(\gamma)|^2 \propto (\lambda_{ph}/d)^6 \quad \text{for } \lambda_{ph} \ll d. \quad (34)$$

We hope that contemporary molecular beam epitaxy (MBE) can give narrow 2DEG with high mobility, for which the PCC can be comparable with the PCC = -64 db of the quartz piezoelectric surface (reported by Grill and Weis [17]).

Completely different can be the case of a thin ( $d = 500 \text{ \AA}$ ) high temperature superconducting layer [18]. In such layers with thickness lower than the London penetration depth  $\lambda_L$  it is also possible to propagate 2D plasma waves [19] if the frequency  $\omega_{pl}$  is lower than the superconductors gap  $2\Delta$ . In this case  $n_0(z) = 1/d$ ,  $n = dn^{3d}$  and

$$\chi(\beta d) = \sin(\gamma)/\gamma, \quad (35)$$

$$|\chi(\gamma)|^2 \propto (\lambda_{ph}/d)^2 \quad \text{for } \lambda_{ph} \ll d. \quad (36)$$

Here  $n^{3d}$  is the volume Cooper pair density. In formula (16) it is necessary only to perform the replacement

$$e^2 n^{3d}/m = L^{-1} = c^2 d / 4\pi \lambda_L^2, \quad (37)$$

where the kinetic inductance  $L$  is an experimental measurable quantity [18]. In spite of much lower deformation potential for the YBaCuO (because of weak binding between different CuO layers), the absence of ohmic dissipation make this layers an ideal system for coherent phonon generation.

In conclusion we hope that a new coherent phonon spectroscopy of 2D plasmons will be developed in the near future.

#### Acknowledgments

The authors would like to thank ICTP, Trieste, where this work was begun, for the hospitality. The authors are also indebted to Prof. G. Mukhopadhyay for the critical analysis and suggestions to bring the manuscript in the present form.

#### References

- [ 1 ] Ando T., Fowler A.B., and Stern F., Rev. Mod. Phys. 54 (1982) 437.
- [ 2 ] Pinczuk A. and Abstreiter G., in *Light Scattering in Solids V* (Topics of Applied Physics 66), edited by Cardona M. and Güntherodt, (Springer-Verlag, 1989, Berlin, Heidelberg, New York), p. 437.
- [ 3 ] Abstreiter G., in *Molecular Beam Epitaxy and Heterostructures*, edited by Chang L.L. and Ploog K., (Martinus Nijhoff, 1985, Dordrecht, Boston, Lancaster), p. 348.
- [ 4 ] Chemla D.S. and Pinczuk A., IEEE J. Quantum Electronics QE-22 (1986) 1609-1921.
- [ 5 ] Olego D., Pinczuk A., Gossard A.C., and Wiegmann W., Phys. Rev. B 25 (1982) 7867.
- [ 6 ] Challis L.J., Toombs G.A. and Sheard F.W., in *Physics of Phonons*, edited by Paszkiewicz, Lecture Notes in Physics, vol. 285 (Springer, 1987, Berlin) p. 348.
- [ 7 ] Krasheninnikov M.V., Sultanov M.B. and Chaplik A.V., Sov. Phys. JETP 50 (1979) 828.
- [ 8 ] Mishonov T.M., J. Phys.: Condens. Matter 1 (1989) 585.
- [ 9 ] Rothenfusser M., Kosler L. and Dietche W., Phys. Rev. 34 (1986) 5518.
- [10] Bron W.E., Rep. Prog. Phys. 43 (1980) 30.
- [11] Allen S.J. Jr., Tsui D.C. and Logan R.A., Phys. Rev. Lett. 38 (1977) 980.
- [12] Batke E., Heitman D., Kotthaus J.P., and Ploog K., Phys. Rev. Lett. 55 (1985) 236.
- [13] Chen W.P., Chen Y.J. and Burstein E.A., Surf. Sci. 58 (1976) 263.
- [14] Price P.J., Ann of Phys. 133 (1981) 217.
- [15] Shayegan M., Goldman V.J., Jiang C., Saioto T., and Santos M., Appl. Phys. Lett. 52 (1988) 1086.
- [16] English J.H., Gossard A.C., Störmer H.L., and Baldwin K.W., Appl. Phys. Lett. 50 (1987) 1826.
- [17] Grill W. and Weis O., Phys. Rev. Lett. 35 (1975) 588.
- [18] Fiory A.T., Hebard A.F., Mankiewich, and Howard R.E., Phys. Rev. Lett. 61 (1988) 1419.
- [19] Mishonov T.M. and Groshev A.V., JINR-Dubna-Commun. E17-89-752 (1989).

Received by Publishing Department  
on December 7, 1989.