


И-304



ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

E17-91-304

R. Tanaś¹, Ts. Gantsog²

QUANTUM FLUCTUATIONS IN THE STOKES
PARAMETERS OF LIGHT PROPAGATING
IN A KERR MEDIUM WITH DISSIPATION

Submitted to "Journal of Modern Optics"

¹Permanent address: Nonlinear Optics Division,
Institute of Physics, Adam Mickiewicz University,
60-780 Poznan, Poland

²On leave from the Department of Theoretical
Physics, Mongolian State University, Ulan Bator
210646, Mongolia


1991

1 Introduction

Optically induced birefringence of an isotropic medium subjected to a strong optical field is a well known fact [1, 2]. Such phenomena like the optical Kerr effect and self-induced ellipse rotation can be explained without recourse to field quantization. On the other hand, if a strong optical field propagating through a nonlinear Kerr medium is treated as a quantum field some new phenomena, like photon antibunching [3]– [5] and squeezing [6], can appear. Since Kerr media are also considered as suitable candidates for performing quantum non-demolition measurements [7, 8], there is growing interest in revealing those aspects of nonlinear propagation that are directly related to the quantum properties of the field.

The polarization state of light propagating through a nonlinear Kerr medium can be effectively described in terms of the Stokes parameters. The Stokes parameters, which are real numbers in the classical description of the field, become Hermitian operators in the quantum description. On having defined the Stokes operators, which are quantum mechanical observables, one is naturally led to address the problem of quantum fluctuations in these quantities as well as quantum field effects on the polarization state of the field propagating in a Kerr medium. Quantum fluctuations in the Stokes parameters of strong light propagating in an isotropic nonlinear medium have recently been discussed by Tanaś and Kielich [9], who treated the medium as ideally transparent, i.e., without losses. Quantum evolution of the field propagating in a Kerr medium has been also considered by Agarwal and Puri [10]. Quite recently, Chaturvedi and Srinivasan [11] using the thermofield dynamics notation have found the exact solution of the master equation for coupled nonlinear oscillators. This solution can be adopted to describe propagation of elliptically polarized light in a Kerr medium with dissipation. An approximate approach based on the Heisenberg-Langevin equations of motion for the operators of the two coupled nonlinear oscillators was given earlier by Horák and Peřina [12].

In this paper we apply the solution of Chaturvedi and Srinivasan [11] to study



quantum fluctuations of the Stokes parameters and the polarization state of strong light propagating in a nonlinear Kerr medium with dissipation. Even including (linear) losses, the exact analytical formulas describing the expectation values and variances of the Stokes operators are derived. The results obtained in this paper are generalizations of earlier results by Tanaš and Kielich [9] into the case of a medium with dissipation.

2 The master equation and its solution

Quantum properties of elliptically polarized light propagating in an isotropic nonlinear Kerr medium can be described by the following effective interaction Hamiltonian [4, 9, 10]

$$H_I = \frac{1}{2}\hbar\kappa \left\{ a_+^{\dagger 2} a_+^2 + a_-^{\dagger 2} a_-^2 + 4da_+^{\dagger} a_-^{\dagger} a_- a_+ \right\}, \quad (1)$$

where a_{\pm} are the annihilation operators of the circularly right (+) and left (-) polarized modes both of frequency ω , the nonlinear coupling constant κ is real and is given by [4, 9]

$$\kappa = \frac{V}{\hbar} \left(\frac{2\pi\hbar\omega}{n^2(\omega)V} \right)^2 2\chi_{xyxy}(\omega) \quad (2)$$

with V denoting the quantization volume, $n(\omega)$ — the linear refractive index of the medium, $\chi_{xyxy}(\omega)$ — the third-order nonlinear susceptibility tensor of the medium. The parameter d in equation (1) is defined by [4]

$$2d = 1 + \frac{\chi_{xxyy}(\omega)}{\chi_{xyxy}(\omega)} \quad (3)$$

and describes the asymmetry of the nonlinear properties of the medium. Ritzke [5] has calculated this asymmetry parameter for atoms with a degenerate one-photon transition obtaining the results

$$d = \begin{cases} (2J-1)(2J+3)/[2(2J^2+2J+1)] & \text{for } J \leftrightarrow J \text{ transitions} \\ (2J^2+3)/[2(6J^2-1)] & \text{for } J \leftrightarrow J-1 \text{ transitions} \end{cases} \quad (4)$$

The coupling between the two modes depends crucially on this asymmetry parameter.

If there is no damping in the system, the interaction Hamiltonian (1) can be directly applied to derive the Heisenberg equations of motion for the field operators which, after replacing the time t by $-n(\omega)z/c$ to deal with the field propagation instead of the field in a cavity, have the simple exponential solutions of the form

$$a_{\pm}(\tau) = \exp \left\{ i\tau \left[a_{\pm}^{\dagger}(0)a_{\pm}(0) + 2da_{\mp}^{\dagger}(0)a_{\mp}(0) \right] \right\} a_{\pm}(0), \quad (5)$$

where

$$\tau = \frac{n(\omega)\kappa z}{c}. \quad (6)$$

To describe the field state evolution the evolution operator $U(\tau)$ can be used, which is given by

$$\begin{aligned} U(\tau) &= \exp \left\{ i\frac{\tau}{2} \left[a_+^{\dagger 2} a_+^2 + a_-^{\dagger 2} a_-^2 + 4da_+^{\dagger} a_-^{\dagger} a_- a_+ \right] \right\} \\ &= \exp \left\{ i\frac{\tau}{2} \left[\hat{n}_+(\hat{n}_+ - 1) + \hat{n}_-(\hat{n}_- - 1) + 4d\hat{n}_+\hat{n}_- \right] \right\} \end{aligned} \quad (7)$$

where we have introduced the number operators $\hat{n}_{\pm} = a_{\pm}^{\dagger} a_{\pm}$ for the two circularly polarized modes. If the initial state of the field is a coherent state of elliptically polarized light one obtains [10]

$$\begin{aligned} |\psi(\tau)\rangle &= U(\tau)|\alpha_+, \alpha_-\rangle \\ &= \sum_{n_+, n_-} b_{n_+}^{(+)} b_{n_-}^{(-)} \exp \left\{ i(n_+\varphi_+ + n_-\varphi_- \right. \\ &\quad \left. + i\frac{\tau}{2} [\hat{n}_+(\hat{n}_+ - 1) + \hat{n}_-(\hat{n}_- - 1) + 4d\hat{n}_+\hat{n}_-] \right\} |\hat{n}_+, \hat{n}_-\rangle, \end{aligned} \quad (8)$$

where

$$b_{n_{\pm}}^{(\pm)} = \exp(-|\alpha_{\pm}|^2/2) \frac{|\alpha_{\pm}|^{n_{\pm}}}{\sqrt{n_{\pm}!}}, \quad \alpha_{\pm} = |\alpha_{\pm}| \exp(i\varphi_{\pm}), \quad (9)$$

α_{\pm} are the amplitudes of the initial coherent states of the two modes, $|\alpha_{\pm}|^2$ are the mean numbers of photons, and φ_{\pm} are the phases of α_{\pm} . Properties of such states have been discussed by Agarwal and Puri [10].

When dissipation is introduced into the system, the state of the field can no longer be described by a pure state, like equation (8), and the density operator description is

necessary. A standard way of introducing the dissipation into the system is to couple it to a reservoir of oscillators and after well known steps, to write down a master equation describing the evolution of the system with dissipation. For the system of two coupled nonlinear oscillators obtained by the Hamiltonian (1), the corresponding master equation in the interaction picture has the following form [11]

$$\frac{\partial \rho}{\partial t} = \frac{1}{i\hbar} [H_I, \rho] + \sum_{i=+,-} \left\{ \frac{\gamma_i}{2} \left([a_i \rho, a_i^\dagger] + [a_i, \rho a_i^\dagger] \right) + \gamma_i \bar{n}_i [a_i, \rho], a_i^\dagger \right\}, \quad (10)$$

where γ_i are the damping constants and \bar{n}_i are the mean numbers of thermal photons.

The exact solution to the master equation (10) has been found by Chaturvedi and Srinivasan [11]. In this paper we adopt their solution to describe propagation of light in a Kerr medium with dissipation. We assume that the reservoir is at zero temperature ($\bar{n}_i = 0$) and that the field is initially in a coherent state. Moreover, we replace the time evolution of the density matrix by the coordinate dependence (assuming that the field propagates along the z coordinate). This means that the damping constants γ_\pm can be interpreted as the absorption coefficients (per unit length) related to the linear absorption of the medium. With these assumptions the solution to the master equation (10), in the number state basis, has the following form

$$\begin{aligned} \rho_{m_+, m_-; n_+, n_-}(\tau) &= \langle m_+, m_- | \rho(\tau) | n_+, n_- \rangle \\ &= b_{m_+}^{(+)} b_{n_+}^{(+)} b_{m_-}^{(-)} b_{n_-}^{(-)} \exp \left[-\frac{\lambda \tau}{2} (\sigma_+ + \sigma_-) + \Gamma_{\delta_+, \delta_-}(\tau) \right] \\ &\times \exp \left\{ i \left[\delta_+ \varphi_+ + \delta_- \varphi_- + \frac{\tau}{2} [\delta_+ (\sigma_+ + 2d\sigma_- - 1) \right. \right. \\ &\left. \left. + \delta_- (\sigma_- + 2d\sigma_+ - 1)] + \Lambda_{\delta_+, \delta_-}(\tau) \right] \right\}, \end{aligned} \quad (11)$$

where the $b_{n_\pm}^{(\pm)}$ are given by (9), and we have introduced the following notation

$$\left. \begin{aligned} \sigma_\pm &= m_\pm + n_\pm \\ \delta_\pm &= m_\pm - n_\pm \end{aligned} \right\}, \quad (12)$$

$$\Gamma_{m,n}(\tau) = \lambda [A_{m,n}^{(+)}(\tau) + A_{n,m}^{(-)}(\tau)] + \eta_{m,n} B_{m,n}^{(+)}(\tau) + \eta_{n,m} B_{n,m}^{(-)}(\tau), \quad (13)$$

$$\Lambda_{m,n}(\tau) = \eta_{m,n} A_{m,n}^{(+)}(\tau) + \eta_{n,m} A_{n,m}^{(-)}(\tau) - \lambda [B_{m,n}^{(+)}(\tau) + B_{n,m}^{(-)}(\tau)], \quad (14)$$

$$\eta_{m,n} = m + 2dn, \quad (15)$$

$$A_{m,n}^{(\pm)}(\tau) = \frac{|\alpha_\pm|^2 \lambda}{\lambda^2 + \eta_{m,n}^2} [1 - \exp(-\lambda \tau) \cos(\eta_{m,n} \tau)], \quad (16)$$

$$B_{m,n}^{(\pm)}(\tau) = \frac{|\alpha_\pm|^2 \lambda}{\lambda^2 + \eta_{m,n}^2} \exp(-\lambda \tau) \sin(\eta_{m,n} \tau). \quad (17)$$

In equations (11)-(17) τ is given by (6), and

$$\lambda = \gamma_+ / \kappa = \gamma_- / \kappa, \quad (18)$$

describes the relative (with respect to the nonlinear coupling κ) damping constant assumed to be the same for both modes of the field.

The solution (11) is exact and, despite the complexity of $\Gamma_{\delta_+, \delta_-}(\tau)$ and $\Lambda_{\delta_+, \delta_-}(\tau)$, its structure is quite transparent. If there is no absorption in the medium, $\lambda = 0$, both $\Gamma_{\delta_+, \delta_-}(\tau)$ and $\Lambda_{\delta_+, \delta_-}(\tau)$ are zero, and the density matrix factorizes into components of the field state (8). The solution (11) will be used to calculate the expectation values and variances of the Stokes operators.

3 Quantum fluctuations in the Stokes parameters

The changes of the polarization of initially elliptically polarized light when it propagates through a nonlinear Kerr medium can be easily accounted for with the use of the Stokes parameters. In quantum treatment of the two-mode field considered in this paper, the following Hermitian Stokes operators can be defined [13]

$$\left. \begin{aligned} S_0 &= a_+^\dagger a_+ + a_-^\dagger a_- \\ S_1 &= a_+^\dagger a_- + a_-^\dagger a_+ \\ S_2 &= -i(a_+^\dagger a_- - a_-^\dagger a_+) \\ S_3 &= a_+^\dagger a_+ - a_-^\dagger a_- \end{aligned} \right\}, \quad (19)$$

where a_\pm (a_\pm^\dagger) are the annihilation (creation) operators of the two circularly polarized modes. If the boson commutation relations are applied for a_\pm and a_\pm^\dagger , it is easy to

check that the Stokes operators themselves satisfy the commutation relations

$$\begin{aligned} [S_1, S_2] &= 2iS_3 \quad \text{and cyclic interchange of indices} \\ [S_i, S_0] &= 0, \quad i = 1, 2, 3. \end{aligned} \quad (20)$$

Moreover, we have

$$S_1^2 + S_2^2 + S_3^2 = S_0(S_0 + 2). \quad (21)$$

The quantum mechanical expectation values of the Stokes operators (19) are the Stokes parameters describing the polarization of the light beam. For elliptically polarized light the parameters of the polarization ellipse are given by [9]

$$\left. \begin{aligned} \tan 2\theta &= \langle S_2 \rangle / \langle S_1 \rangle, \\ \tan 2\eta &= \langle S_3 \rangle / (\langle S_1 \rangle^2 + \langle S_2 \rangle^2)^{1/2}, \end{aligned} \right\} \quad (22)$$

where θ is the azimuth of the polarization ellipse denoting the angle between the major axis of the polarization ellipse and the x axis, and η is the ellipticity parameter, $-\pi/4 \leq \eta \leq \pi/4$; $\tan \eta$ describes the ratio of the semi-minor axis and the semi-major axis of the polarization ellipse and the sign defines its handedness (plus indicates right-handed polarization on the helicity convention).

The degree of polarization of the field can be defined as

$$P = \frac{(\langle S_1 \rangle^2 + \langle S_2 \rangle^2 + \langle S_3 \rangle^2)^{1/2}}{\langle S_0 \rangle}. \quad (23)$$

For the initial coherent state of elliptically polarized light the Stokes parameters have the values

$$\left. \begin{aligned} \langle \alpha_+, \alpha_- | S_0 | \alpha_+, \alpha_- \rangle &= |\alpha_+|^2 + |\alpha_-|^2 = |\alpha|^2 \\ \langle \alpha_+, \alpha_- | S_1 | \alpha_+, \alpha_- \rangle &= 2\text{Re}(\alpha_+^* \alpha_-) = |\alpha|^2 \cos 2\eta \cos 2\theta \\ \langle \alpha_+, \alpha_- | S_2 | \alpha_+, \alpha_- \rangle &= 2\text{Im}(\alpha_+^* \alpha_-) = |\alpha|^2 \cos 2\eta \sin 2\theta \\ \langle \alpha_+, \alpha_- | S_3 | \alpha_+, \alpha_- \rangle &= |\alpha_+|^2 - |\alpha_-|^2 = |\alpha|^2 \sin 2\eta \end{aligned} \right\}, \quad (24)$$

where $|\alpha|^2$ is the total mean number of photons in the field while $\theta = (\varphi_- - \varphi_+)/2$ and η define the polarization ellipse. The degree of polarization P is in this case equal to unity. This means that the coherent state of the field corresponds to a classical,

fully polarized field. However, the non-commutability of the Stokes operators puts well known limits on measurements of the physical quantities represented by these operators. For example, according to the commutation relations (20), we have the following Heisenberg uncertainty relation

$$((\Delta S_1)^2)(\Delta S_2)^2)^{1/2} \geq |\langle S_3 \rangle|. \quad (25)$$

Quantum fluctuations in the Stokes parameters of light propagating in a Kerr medium without losses have been discussed by Tanaš and Kielich [9]. In this paper we wish to generalize those results including dissipation into the system. The exact solution (11) to the master equation enables us to derive exact analytical formulas for the expectation values and variances of the Stokes operators as well as the characteristics of the field polarization for the medium with dissipation.

The expectation values of the Stokes operators are given by

$$\begin{aligned} \langle S_0 \rangle &= \sum_{n,m=0}^{\infty} (n+m) \rho_{n,m;n,m}(\tau) \\ &= \sum_{n,m=0}^{\infty} (n+m) b_n^{(+)} b_m^{(-)} \exp[-\lambda\tau(n+m)] \\ &\quad \times \exp\{(|\alpha_+|^2 + |\alpha_-|^2)[1 - \exp(\lambda\tau)]\} \\ &= |\alpha|^2 \exp(-\lambda\tau), \end{aligned} \quad (26)$$

$$\begin{aligned} \langle S_1 \rangle &= 2\text{Re} \sum_{n,m=0}^{\infty} \sqrt{(n+1)(m+1)} \rho_{n,m+1;n+1,m}(\tau) \\ &= 2|\alpha_+||\alpha_-| \exp\left\{-\lambda\tau + \Gamma_{-1,1}(\tau)\right. \\ &\quad \left. + (|\alpha_+|^2 + |\alpha_-|^2) \left\{ \exp(-\lambda\tau) \cos[(1-2d)\tau] - 1 \right\}\right\} \\ &\quad \times \cos\{\varphi_+ - \varphi_- + (|\alpha_+|^2 - |\alpha_-|^2) \exp(-\lambda\tau) \sin[(1-2d)\tau] - \Lambda_{-1,1}(\tau)\} \\ &= |\alpha|^2 \cos 2\eta \exp\{-\lambda\tau + \Gamma_{-1,1}(\tau) + |\alpha|^2 \left\{ \exp(-\lambda\tau) \cos[(1-2d)\tau] - 1 \right\}\right. \\ &\quad \left. \times \cos\{2\theta - |\alpha|^2 \sin 2\eta \exp(-\lambda\tau) \sin[(1-2d)\tau] + \Lambda_{-1,1}(\tau)\}\right\}, \end{aligned} \quad (27)$$

$$\begin{aligned} \langle S_2 \rangle &= 2\text{Im} \sum_{n,m=0}^{\infty} \sqrt{(n+1)(m+1)} \rho_{n,m+1;n+1,m}(\tau) \\ &= |\alpha|^2 \cos 2\eta \exp\{-\lambda\tau + \Gamma_{-1,1}(\tau) + |\alpha|^2 \left\{ \exp(-\lambda\tau) \cos[(1-2d)\tau] - 1 \right\}\right\} \end{aligned}$$

$$\times \sin \{2\theta - |\alpha|^2 \sin 2\eta \exp(-\lambda\tau) \sin[(1-2d)\tau] + \Lambda_{-1,1}(\tau)\}, \quad (28)$$

$$\langle S_3 \rangle = \sum_{n,m=0}^{\infty} (n-m) \rho_{n,m;n,m}(\tau) = |\alpha|^2 \sin 2\eta \exp(-\lambda\tau), \quad (29)$$

where $|\alpha|^2$, $2\theta = \varphi_- - \varphi_+$, and η are the parameters defining initial state of the field; λ describes the dissipation of the medium, and $\Gamma_{-1,1}(\tau)$ and $\Lambda_{-1,1}(\tau)$ are given by (13) and (14) or (A.3) and (A.4). It is to be kept in mind that the latter two quantities depend on τ , and that they vanish for $\lambda = 0$. For $\lambda = 0$ the results (26)–(29) go over into the earlier results of Tanaš and Kielich [9].

According to (22) and (23), from (26)–(29) we obtain

$$\tan 2\theta(\tau) = \tan \{2\theta - |\alpha|^2 \sin 2\eta \exp(-\lambda\tau) \sin[(1-2d)\tau] + \Lambda_{-1,1}(\tau)\}, \quad (30)$$

$$\tan 2\eta(\tau) = \exp \{-|\alpha|^2 \{\exp(-\lambda\tau) \cos[(1-2d)\tau] - 1\}\} \tan 2\eta, \quad (31)$$

$$P^2(\tau) = \sin^2 2\eta + \cos^2 2\eta \exp \left\{ 2|\alpha|^2 \{\exp(-\lambda\tau) \cos[(1-2d)\tau] - 1\} + 2\Gamma_{-1,1}(\tau) \right\}. \quad (32)$$

Formulas (30)–(32) are exact quantum expressions describing the evolution of the polarization state of elliptically polarized field propagating in a Kerr medium with dissipation. It is evident from (30)–(32) (see also the Appendix) that all polarization parameters depend on $1-2d$, that is, they crucially depend on the asymmetry parameter d of the nonlinear medium. For $d = 1/2$, i. e. for $1/2 \leftrightarrow 1/2$ transitions, the polarization state of the field does not change. For the fully symmetric susceptibility tensor $\chi(\omega)$, $d = 1$, and there are changes in the polarization of the field. From (30) and (A.2) we obtain the following expression describing the rotation of the polarization ellipse

$$\begin{aligned} \theta(\tau) &= \theta - \frac{1}{2} \sin 2\eta \frac{1-2d}{\lambda^2 + (1-2d)^2} \left\{ (1-2d) |\alpha|^2 \exp(-\lambda\tau) \sin[(1-2d)\tau] \right. \\ &\quad \left. + \lambda [|\alpha|^2 - |\alpha|^2 \exp(-\lambda\tau) \cos[(1-2d)\tau]] \right\} \\ &= \theta - \frac{1}{2} \sin 2\eta \frac{1-2d}{\lambda^2 + (1-2d)^2} \left\{ (1-2d) S_0(\tau) \sin[(1-2d)\tau] \right. \\ &\quad \left. + \lambda [S_0(0) - S_0(\tau) \cos[(1-2d)\tau]] \right\}, \quad (33) \end{aligned}$$

where

$$S_0(\tau) = \langle S_0 \rangle = |\alpha|^2 \exp(-\lambda\tau), \quad (34)$$

according to (26).

Formula (33) is the exact quantum formula, which evolves in two different “timescales” (in fact lengthscales): one associated with the nonlinearity of the medium (τ), and another one related to the dissipation ($\lambda\tau$). For $\lambda = 0$, formula (33) reproduces the earlier obtained result [9], which is still the quantum result and is periodic in $(1-2d)\tau$. Only for $\tau \ll 1$, when $\sin[(1-2d)\tau] \approx (1-2d)\tau$, the classical result is obtained. Another limit is $\lambda \gg 1$, in which only the term proportional to λ in the large braces contributes if the terms of order λ^{-1} are retained. Again, the transition to the classical field is obtained by taking $\tau \ll 1$, i. e., by replacing $\cos[(1-2d)\tau]$ with unity. This gives us the quantity $[S_0(0) - S_0(\tau)]/\lambda$ that appears in the solution of the problem for the classical field and the medium with dissipation. It is interesting to note that, despite the fact that this classical result has been obtained from the quantum formula under assumption $\lambda \gg 1$, putting $\lambda = 0$ in the resulting formula leads to the same classical result which is obtained from the first term in the large braces of eq. (33) for $\lambda = 0$ and $\tau \ll 1$. Thus, the classical result without dissipation can be obtained from eq. (33) in two different ways. For $\lambda \approx 1$, the effects of nonlinearity and dissipation are equally important, and in this case formula (33) must be used in its full form. In Fig. 1 the evolution of $\theta(\tau)$ is shown for different values of λ , $\eta = \pi/8$, $d = 1$, and $|\alpha|^2 = 0.25, 1, 4, 16$.

The ellipticity $\eta(\tau)$ given by eq. (31) exhibits the τ -dependence that is a purely quantum effect (we have assumed the same absorption for both circular components of the field). The exponential does not appear for classical fields. For $\lambda = 0$, eq. (31) goes over into the earlier result [9]. The evolution of $\eta(\tau)$ is drawn in Fig. 2 for different values of λ , $\eta = \pi/8$, $d = 1$, and $|\alpha|^2 = 0.25, 1, 4, 16$. As it is seen from Fig. 2, the ellipticity $\eta(\tau)$ approaches $\pi/4$ when $|\alpha|^2 \gg 1$. This means the circular polarization of the field. Thus the quantum fluctuations of the field cause that the field which remains polarized during the propagation can only be circularly polarized. For $\lambda = 0$

there is a quantum periodicity of the evolution which is removed by the dissipation. For $\eta = 0$, according to (31), there is no change in the ellipticity of the field which suggests that initially linear polarization remains linear during the evolution. This is true when speaking about the polarized part of the field. However, one can easily check, using eq. (32), that in this case the degree of polarization rapidly deeps to zero, and there is no, practically, the polarized part of the field. The degree of polarization $P(\tau)$ is plotted in Fig. 3 for $\eta = 0; d = 1$, and different values of $|\alpha|^2$ and λ . The reduction of the degree of polarization is quite evident, and the dissipation prevents the recurrence of the initial degree of polarization. From eq. (32) it is seen that there is a lower bound for the degree of polarization equal to $|\sin 2\eta|$. This means, for example, that for $\eta = \pi/8$ taken in Figs. 1 and 2 the degree of polarization $P(\tau)$ cannot fall below the value $1/\sqrt{2}$, i. e., the field retains quite a bit of its polarization. For the circular polarization the degree of polarization does not change at all. So, the effect of quantum field fluctuations is most dramatic when linearly polarized light propagates through an isotropic nonlinear medium. The changes in the degree of polarization for media without dissipation were discussed earlier [9, 10, 14]. We have also discussed the relation of the Stokes parameters with the phase properties of the field [15].

To study the quantum fluctuations in the Stokes parameters, we calculate the variances of the Stokes operators and look at their behaviour during the evolution. For the expectation values of the squares of the Stokes operator we get the relations

$$\begin{aligned}
\langle S_{1,2}^2 \rangle &= \pm 2 \operatorname{Re} \sum_{n,m} \sqrt{(n+1)(n+2)(m+1)(m+2)} \rho_{n,m+2;n+2,m}(\tau) \\
&\quad + \sum_{n,m} (n+m+2nm) \rho_{n,m;n,m} \\
&= \pm \frac{1}{2} |\alpha|^4 \cos^2 2\eta \exp \left\{ |\alpha|^2 [\exp(-\lambda\tau) \cos[2(1-2d)\tau] - 1] \right. \\
&\quad \left. - 2\lambda\tau + \Gamma_{-2,2}(\tau) \right\} \cos \left\{ 4\theta - |\alpha|^2 \sin 2\eta \exp(-\lambda\tau) \right. \\
&\quad \left. \times \sin[2(1-2d)\tau] + \Lambda_{-2,2}(\tau) \right\} \\
&\quad + \frac{1}{2} |\alpha|^4 \cos^2 2\eta \exp(-2\lambda\tau) + |\alpha|^2 \exp(-\lambda\tau), \quad (35)
\end{aligned}$$

$$\begin{aligned}
\langle S_3^2 \rangle &= \sum_{n,m} (n+m)^2 \rho_{n,m;n,m}(\tau) \\
&= |\alpha|^4 \sin^2 2\eta \exp(-2\lambda\tau) + |\alpha|^2 \exp(-\lambda\tau). \quad (36)
\end{aligned}$$

Formulas (35) and (36) together with (26)–(29) and (A.1)–(A.6) allow calculations of the variances of the Stokes parameters. The variances are intensity dependent, i. e., the quantum noise related with the measurement of the Stokes parameters is also intensity dependent. However, it is rather the relative noise, or the signal-to-noise ratio, that are interesting from the experimental point of view. One can ask the question: How will the quantum fluctuations of the field affect the signal-to-noise ratio? Our exact analytical formulas will give immediately the answer to this question. Let us define the signal-to-noise ratio for the measurements of the Stokes parameters as

$$R_i(\tau) = \frac{|\langle S_i \rangle|}{\sqrt{\langle S_i^2 \rangle - \langle S_i \rangle^2}}, \quad (i = 1, 2, 3). \quad (37)$$

The evolution of the signal-to-noise ratio (37) is plotted in Figs. 4-7 for various sets of the parameters. In Fig. 4 there are plots of $R_1(\tau)$ for $\eta = 0, d = 1, |\alpha|^2 = 0.25, 1, 4, 16$, and (a) $\lambda = 0$, (b) $\lambda = 0.1$. For $\lambda = 0$, i. e. the absence of damping in the system, the evolution of the signal-to-noise ratio is periodic. The initial value of the ratio is $|\alpha|$, as it should be for the coherent state $|\alpha\rangle$. However, owing to the quantum fluctuations of the field the ratio rapidly falls down if the field is strong. That is, even without damping in the system, the signal-to-noise ratio deteriorates drastically making the measurement of $\langle S_1 \rangle$ less certain. The presence of damping deteriorates the signal-to-noise ratio still further, as it is seen from Fig. 4.(b), and it removes the periodicity of the evolution. For the linear polarization of the initial field, $\eta = 0$ and $\sin \eta = 0$, and both $\langle S_2 \rangle$ and $\langle S_3 \rangle$ are zero all the time. That is, the only signal we can measure in the case of linear polarization is $\langle S_1 \rangle$.

For elliptical polarization with $\eta = \pi/8$, we have the situation illustrated in Figs. 5-7, where $R_1(\tau), R_2(\tau)$, and $R_3(\tau)$ are plotted for $\eta = \pi/8$ and all other parameters being the same as in Fig. 4. The signal-to-noise ratio $R_1(\tau)$ is lower initially than for linear polarization, it is now $|\alpha| |\cos 2\eta|$ instead of $|\alpha|$, there are some oscillations at




FIG. 1. Evolution of the azimuth $\theta(\tau)$ for $\eta = \pi/8$. The other parameters are taken the same for all the figures, and they are: $d = 1$, $|\alpha|^2 = 0.25$ (dotted line), 1 (dashed line), 4 (dashed-dotted line), 16 (solid line). Figure (a) is for $\lambda = 0$, and (b) for $\lambda = 0.1$.




FIG. 2. Evolution of the ellipticity $\eta(\tau)$, for $\eta = \pi/8$.




FIG. 3. Evolution of the degree of polarization $P(\tau)$, for $\eta = 0$.




FIG. 4. Evolution of the signal-to-noise ratio $R_1(\tau)$, for $\eta = 0$.




FIG. 5. Same as Fig. 4, but for $\eta = \pi/8$.




FIG. 6. Evolution of $R_2(\tau)$, for $\eta = \pi/8$.




FIG. 7. Evolution of $R_3(\tau)$, for $\eta = \pi/8$.

the initial stages of the evolution if the field is strong, but deterioration of the ratio owing to quantum fluctuations and dissipation is evident. The ratio $R_2(\tau)$, shown in Fig. 6, grows up initially indicating the conjugate character of S_2 with respect to S_1 , as one could expect from the commutation relation (20) and the uncertainty relation (25). However, at later times quantum fluctuations of the field deteriorate this ratio as well. Of course, the dissipation still worsens the situation.

The behaviour of $R_3(\tau)$ is different: if there is no damping $R_3(\tau)$ is constant and equal to $|\alpha| |\sin 2\eta|$, and it is damped when $\lambda \neq 0$. This is shown in Fig. 7. So, the quantum fluctuations of the field do not deteriorate the precision of measurement of the Stokes parameter $\langle S_3 \rangle$. This can be explained by the fact that this Stokes parameter describes the circularly polarized part of the field which is not affected during the propagation in the isotropic nonlinear medium. The only evolution is the result of the linear damping of the field. The quantum character of the field propagating in an isotropic Kerr medium manifests itself most strongly when the field is linearly polarized.

4 Conclusion

We have considered the quantum field of elliptical polarization propagating in an isotropic, nonlinear Kerr medium with dissipation finding the exact analytical formulas describing the evolution of the Stokes parameters, which define the polarization of the field, and the evolution of their variances, which describe quantum fluctuations in the Stokes parameters. We have applied the exact solution of the master equation obtained recently by Chaturvedi and Srinivasan [11] to find the evolution of the Stokes parameters and their variances. The quantum character of the field affects essentially the polarization of light propagating in the medium. In case of linear polarization there is a rapid decrease of the degree of polarization. For the elliptical polarization there is a lower bound for the degree of polarization equal to $|\sin 2\eta|$, which means that the circular polarization remains unchanged during the evolution. The results for the azimuth of the polarization ellipse, its ellipticity, and the degree of polarization are

illustrated graphically for various sets of parameters. There are quantum effects that can be found in the evolution of these quantities despite the fact that they are defined by the expectation values of the Stokes operators that are linear in intensity.

The quantum noise of the Stokes parameters of light propagating in a Kerr medium is discussed in detail. The evolution of the signal-to-noise ratio for the measurements of the Stokes parameters is studied. It is shown that quantum fluctuations of the strong field drastically deteriorate the signal-to-noise ratio for the measurements of the Stokes parameters $\langle S_1 \rangle$ and $\langle S_2 \rangle$ related with the linear polarization of the field. The parameter $\langle S_3 \rangle$, related to the circular polarization of the field, is not affected by the quantum fluctuations. As one could expect, the dissipation in the system lowers the signal-to-noise ratio in any case. Our exact formulas that include linear damping allow for the quantitative assessment of its destructive role in the quantum evolution of the field.

Appendix

For convenience, in this Appendix we write down the explicit expressions for $\Gamma_{m,n}(\tau)$ and $\Lambda_{m,n}(\tau)$ needed for our calculations in this paper. Let us define the quantities

$$A_k(\tau) = \lambda \frac{|\alpha|^2 \lambda}{\lambda^2 + [k(1-2d)]^2} (1 - \exp(-\lambda\tau) \cos[k(1-2d)\tau]) \quad (\text{A.1})$$

$$B_k(\tau) = \frac{|\alpha|^2 \lambda}{\lambda^2 + [k(1-2d)]^2} \exp(-\lambda\tau) \sin[k(1-2d)\tau]. \quad (\text{A.2})$$

Using (A.1) and (A.2), from the general expressions (13)–(17), we can derive the following relations:

$$\Gamma_{-1,1}(\tau) = \lambda A_1(\tau) + (1-2d)B_1(\tau), \quad (\text{A.3})$$

$$\Lambda_{-1,1}(\tau) = \sin 2\eta [-(1-2d)A_1(\tau) + \lambda B_1(\tau)], \quad (\text{A.4})$$

$$\Gamma_{-2,2}(\tau) = \lambda A_2(\tau) + 2(1-2d)B_2(\tau), \quad (\text{A.5})$$

$$\Lambda_{-2,2}(\tau) = \sin 2\eta [-2(1-2d)A_2(\tau) + \lambda B_2(\tau)]. \quad (\text{A.6})$$

References

- [1] Kielich, S., 1981, *Nonlinear Molecular Optics* (Moscow: Nauka)
- [2] Shen, Y. R., 1985, *The Principles of Nonlinear Optics* (New York: Wiley)
- [3] Ritze, H. H., and Bandilla, A., 1979, *Optics Commun.* **29**, 126
- [4] Tanaś, R., and Kielich, S., 1979, *Optics Commun.* **30**, 443
- [5] Ritze, H. H., 1980, *Z. Phys.* B **39**, 353
- [6] Tanaś, R., and Kielich, S., 1983, *Optics Commun.* **45**, 351;
1984, *Optica Acta*, **31**, 81
- [7] Levenson, M. D., Shelby, R. M., Reid, M., and Walls, D. F., 1986, *Phys. Rev. Lett.*, **57**, 2473
- [8] Bachor, H. A., Levenson, M. D., Walls, D. F., Perlmutter, S. H., and Shelby, R. M., 1988, *Phys. Rev. A*, **38**, 180
- [9] Tanaś, R., and Kielich, S., 1990, *J. Mod. Opt.*, **37**, 1935
- [10] Agarwal, G. S., and Puri, R. R., 1989, *Phys. Rev. A*, **40**, 5179
- [11] Chaturvedi, S., and Srinivasan, V., 1991, *Phys. Rev. A*, **43**, ***
- [12] Horák, R., and Peřina, J., 1989, *J. Opt. Soc. Am. B*, **6**, 1239
- [13] Jauch, J. M., and Rohrlich, F., 1959, *The Theory of Photons and Electrons* (London: Addison-Wesley)
- [14] Horák, R., 1989, *Optics Commun.*, **72**, 239
- [15] Tanaś, R., and Gantsog, Ts., Quantum effects in the polarization of light propagating in a Kerr medium, to be published

Received by Publishing Department
on July 2, 1991.

Танась Р., Ганцог Ц. E17-91-304
Квантовые флуктуации параметров Стокса
для света, распространяющегося в ячейке
Керра с диссипацией

Для света, распространяющегося в ячейке Керра с диссипацией, получены точные аналитические выражения, описывающие эволюцию средних значений и дисперсией операторов Стокса. Показано, что квантовые флуктуации поля существенно возмущают поляризацию поля. Получены и проиллюстрированы графически явные квантовые соотношения, описывающие как наклон и эллиптичность поляризационного эллипса, так и степень поляризации. Обсуждаются квантовые флуктуации параметров Стокса, показано, что в процессе эволюции отношение сигнала к шуму должно сокращаться квантовыми флуктуациями. Явно показана роль диссипации на основе точного аналитического решения.

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1991

Tanaś R., Gantsog Ts. E17-91-304
Quantum Fluctuations in the Stokes
Parameters of Light Propagating
in a Kerr Medium with Dissipation

Exact analytical expressions describing the evolution of the expectation values and the variances of the Stokes operators are derived for the elliptically polarized light propagating in a Kerr medium with dissipation. It is shown that quantum fluctuations of the field essentially affect the polarization of the field. The explicit quantum formulas describing the azimuth and the ellipticity of the polarization ellipse as well as the degree of polarization are derived and illustrated graphically. Quantum fluctuations in the Stokes parameters are discussed, and the evolution of the signal-to-noise ratio is shown to be reduced by the quantum field fluctuations. Role of the dissipation is shown explicitly in a fully quantitative way from the exact analytical solutions.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research, Dubna 1991