

5329

ЭКЗ. ЧИТ. ЗАЛА

СООБЩЕНИЯ
ОБЪЕДИНЕННОГО
ИНСТИТУТА
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

Дубна

E2 - 5329



**В.М. Barbashov, S.P. Kuleshov,
V.A. Matveev, V.N. Pervushin, A.N. Sissakian**

ЛАБОРАТОРИЯ ТЕОРЕТИЧЕСКОЙ ФИЗИКИ

**THE POISSON DISTRIBUTION
OF SECONDARY PARTICLES
IN STRAIGHT-LINE PATHS
APPROXIMATION**

1970

E2 - 5329

**B.M. Barbashov, S.P. Kuleshov,
V.A. Matveev, V.N. Pervushin, A.N. Sissakian**

**THE POISSON DISTRIBUTION
OF SECONDARY PARTICLES
IN STRAIGHT-LINE PATHS
APPROXIMATION**

I. Introduction

In recent papers^{/1-5/} a straight-line paths approximation (SLPA) has been formulated which then has been applied to considering problems of high-energy particle scattering in the framework of standard models of quantum field theory. This approximation is closely associated with the methods which are extensively being developed by many authors^{/6-9/} for the so-called eikonal summation of the Feynmann graphs. The eikonal approximation which is used for studying the asymptotic behaviour of the sum of a definite class of perturbation diagrams is based on the modification of the nucleon propagator according to which the bilinear terms on the mesons momenta are rejected. This modification has been studied well and proved in the infrared region^{/10-13/}. By the present time we have already some grounds of its validity for a certain class of s -channel diagrams^{/14,15/} in the asymptotic domain of high energies and fixed momentum transfers.

In formulating SLPA it is convenient to start from the Feynman interpretation of the scattering amplitude as a sum over the paths. In so doing, the method of averaging over the functional variable used in refs.^{/1-5/} and in the present paper is equivalent to the account of paths which approach most closely the straight-line ones. In the case of high-energy and fixed momentum transfer scattering the particle trajectories are approximately intercepts of straight lines having the direction of the particle momenta before and after collisions,

respectively. Such a physical picture permits us to give to the method used the name of the straight-line paths approximation.

The present paper is devoted to the investigation of the behaviour of the distribution of secondary "mesons" produced in high-energy "nucleon" collisions on the SLPA basis. It is useful to study aspects of this problem in order to clarify the hadron interaction mechanism in the asymptotic domain in question.

For the sake of simplicity, we shall consider the field theory models in which scalar nucleons^{x/} exchange scalar and vector mesons. The inelastic amplitude and the n particle production cross section are factorized in these models under the condition that the components of the emitted meson momenta be restricted by the "softness" conditions.

It is interesting to study the differential cross section of inelastic processes, which in SLPA has the shape of the Poisson distribution over the secondary particle number, as a function of the choice of the integral cutoff parameters in various asymptotic regions.

It should be noted that the Poisson distribution was considered earlier in various phenomenological models/16/, electrodynamics/17-20/, as well as in other approaches. Recently it has been shown/21,1/ that such a distribution corresponds to the physical picture where the nucleons interact via their meson "clouds". It is interesting to note that the total differential cross section summed up over all the emitted mesons may have no pronounced diffraction peak in a certain domain of momentum transfers. In this connection an analogy should be indicated with the automodel behaviour of the cross sections of high-energy deep inelastic interactions of hadrons with leptons/22,23/.

The problems formulated above are studied first on a simple example of potential scattering. Further the consideration is made in the framework of ordinary quantum field theory models.

^{x/}In the framework of this method a generalization may be made to the case of spinor particles, too.

2. Secondary Particle Distribution on Nucleon-Potential Scattering

We consider a potential scattering with multiple production of secondary particles. The field with which the scattered nucleon interacts is represented as the sum of the external (classic) V and quantized ϕ fields. The amplitude of the scattering of a nucleon in the external field with n meson production is found according to the well-known formula

$$f_n(p, q; k_1, \dots, k_n) = \langle 0 | T_\phi F(p, q | g\phi + V) | k_1, \dots, k_n \rangle, \quad (2.1)$$

where T_ϕ is the sign of the T -product of the operators ϕ .

The generating functional $F(p, q | g\phi + V)$ is expressed through the Fourier transform of the nucleon Green function $G(p, q | g\phi + V)$ as follows

$$F(p, q | g\phi + V) = \lim_{(p^2, q^2) \rightarrow m^2} (p^2 - m^2)(q^2 - m^2) G(p, q | g\phi + V). \quad (2.2)$$

The transition to the mass shell is easily performed using the method of ref.^{24/}. Thus, after extracting the poles, eq. (2.2) takes the form^{x/}

$$F(p, q | g\phi + V) = \int d^4 x e^{i x(p-q)} \langle \Gamma(x; \nu) e^{i g \int d^4 y \phi(y) J(x, y; \nu)} \rangle, \quad (2.3)$$

^{x/} In the transition to the mass shell the terms corresponding to the diagrams without interaction of the nucleon with the field V are eliminated.

where

$$J(x, y; \nu) = \int d\xi \delta^4 [x + a(\xi) + 2 \int_0^\xi \nu(\eta) d\eta - y]; \quad a(\xi) = 2p\theta(\xi) + 2q\theta(-\xi), \quad (2.4)$$

$$\Gamma(x; \nu) = V(x) \int_0^1 d\lambda \exp \{ i\lambda \int d^4 z V(z) J(x, z; \nu) \} \quad (2.5)$$

and the averaging over ν means

$$\langle \Phi(\nu) \rangle_\nu = \frac{\int d^4 \nu \exp \{ -i \int \nu^2(\eta) d\eta \} \Phi(\nu)}{\int d^4 \nu \exp \{ -i \int \nu^2(\eta) d\eta \}}. \quad (2.6)$$

Let us assume that all the particles produced (both real and virtual) are "soft" mesons/1/, namely that the region of change of their momenta D is limited so that they do not lead to a strong change of the momentum transfer. It is easily seen that the x -dependence in eq. (2.4) can be neglected.

We neglect the variable ν which describes the deflection of particles from the straight-line paths.

Under these assumption the functional F takes the form

$$F(p, q | g\phi + V) = f_{el}(p, q) \exp \{ ig \int_D d^4 y \phi(y) J(y) \}, \quad (2.7)$$

where the elastic scattering amplitude^{/25/} is expressed as follows

$$f_{el}(p, q) = \int d^4x e^{ix(p-q)} \langle \Gamma(x; \nu) \rangle_\nu. \quad (2.8)$$

Hence, it is not difficult to get^{/17/} that the cross section of scattering on a potential with n "soft" meson production has the shape of the Poisson distribution. Indeed,

$$f_n = f_{el} \langle 0 | T_\phi \exp \left\{ ig \int_D d^4x \phi(x) J(x) \right\} | n \rangle = \quad (2.9)$$

$$= f_{el} \frac{1}{\sqrt{n!}} \left(\prod_{i=1}^n \frac{J(k_i)}{(2\pi)^{3/2} \sqrt{2k_{0i}}} \right) \exp \left\{ \frac{ig^2}{2} \int_D d^4k J^2(k) D(k) \right\}$$

where $D(k)$ is the casual function. Consequently,

$$d\sigma_n = d\sigma_{el} \frac{1}{n!} \left[\frac{1}{(2\pi)^3} \int_D \frac{d^3k J^2(k)}{2k_0} \right]^n. \quad (2.10)$$

Now taking into account that

$$D(k) - D^*(k) = \frac{i}{(2\pi)^3} \delta(k^2 - m^2) \quad (2.11)$$

we finally get

$$d\sigma_n = d\sigma_0 \frac{a^n}{n!} e^{-a}, \quad (2.12)$$

where a is defined by the following formula

$$a = \frac{g^2}{(2\pi)^3} \int_D \frac{d^3 k J^2(k)}{2k_0} \quad (2.13)$$

and the current $J(k)$ is of the form

$$J(k) = \frac{1}{2pk + i\epsilon} - \frac{1}{2qk - i\epsilon} \quad (2.14)$$

If the momentum transfer is in the region $s \gg t \gg m^2$, $\mu^2 \neq 0$ then

$$a \sim g^2 \frac{\ln^2 \alpha t}{t} \quad (2.15)$$

where α depends only on masses.

Note that in electrodynamics, in eq. (2.15), there appears an infrared singularity and the factor $\frac{1}{t}$ vanishes. In this case the asymptotics has a so-called doubly-logarithmic character $\sqrt{20}$.

3. Poisson Distribution of Secondary Particles in Quantum Field Theory Models

Now we proceed to the case of multiple production of "soft" particles in the model $L_{int} = g:\psi^2 \phi:$, where, for simplicity, the

fields ψ and ϕ are assumed to be scalar. The vector exchange model was considered earlier^{1/}. Here we present only some results of this model.

The amplitude of scattering of two nucleons (field ψ) with meson production (field ϕ) is determined by the formula^{25/}

$$(2\pi)^4 \delta(p_1 + p_2 - q_1 - q_2 - \sum_{i=1}^n k_i) f(p_1, p_2; q_1, q_2; k_1, \dots, k_n) = \quad (3.1)$$

$$= \langle 0 | T_{\phi_1} T_{\phi_2} F(p_1, p_2; q_1, q_2; \phi_1, \phi_2) | n \rangle,$$

where the functional F , after the transition to the mass shell, is of the form^{5/}

$$F(p_1, p_2; q_1, q_2; \phi_1, \phi_2) = \int dx_1 dx_2 \ll A(x_1, x_2; \nu_1, \nu_2) \cdot \exp \{ i g \int d^4 x \sum_{i=1}^2 \phi_i(x) J_i(x; x_1, x_2; \nu_1, \nu_2) \} \gg_{\nu_1, \nu_2} \quad (3.2)$$

The following notations are introduced:

$$A(x_1, x_2; \nu_1, \nu_2) = \exp \{ i(p_1 - q_1)x_1 + i(p_2 - q_2)x_2 \} D(x_1 - x_2) \cdot \quad (3.3)$$

$$\cdot \int_0^1 d\lambda \exp \{ i\lambda g^2 \int d\xi_1 d\xi_2 D[x_1 - x_2 + a_1(\xi_1) - a_2(\xi_2) + \int_0^{\xi_1} \nu_1(\eta) d\eta - \int_0^{\xi_2} \nu_2(\eta) d\eta] \}$$

$$J_1(x; x_1, x_2; \nu_1, \nu_2) = \int d\xi \delta^4[x_1 - x + a_1(\xi) + \int_0^\xi \nu_1(\eta) d\eta] \quad (3.4)$$

and the symbol $\langle\langle \dots \rangle\rangle_{\nu_1, \nu_2}$ means the averaging over the both functional variables ν_1 and ν_2 (eq. 2.6).

We consider the case when the produced meson momenta are negligibly small compared with the nucleon momenta. This makes it possible to neglect the x_1 and x_2 dependence in eq. (3.4) for quantum current J_1 . In other words, we consider the production of the so-called "soft" mesons which do not nearly affect the motion of scattered rapid particles.

According to the method developed and employed in refs./10, 12, 1-5/ for the estimation of the functional integrals over ν we use the following approximation

$$\begin{aligned} \langle\langle A(x_1, x_2; \nu_1, \nu_2) \exp\{ig \int d^4x \phi(x) J(x; \nu_1, \nu_2)\} \rangle\rangle_{\nu_1, \nu_2} = \\ = \langle\langle A(x_1, x_2; \nu_1, \nu_2) \rangle\rangle_{\nu_1, \nu_2} \exp\{ig \int d^4x \phi(x) J(x)\}, \end{aligned} \quad (3.5)$$

where

$$J(x) = \langle\langle J(x; \nu_1, \nu_2) \rangle\rangle_{\nu_1, \nu_2}.$$

As a first simple example we consider the case when both the real and virtual mesons are "soft". In order to take into account the "softness" of mesons we cut off the region of integration over the meson momenta. Owing to the fact that now the integrals are no longer divergent in the upper limit, the averaging over ν , according to eq. (3.5), may be replaced by other more rough approxima-

tion when the F dependence in quantum currents is eliminated. As was already noted in the case of potential scattering, this approximation means an entire neglect of the deflexion nucleon from straight-line paths.

Thus, in this approximation the functional F takes the form

$$F(p_1, p_2; q_1, q_2; \phi_1, \phi_2) = f_{el}(p_1, p_2; q_1, q_2) \exp \left\{ i \int_D d^4k \sum_{i=1}^2 \phi_i(k) J_i(k) \right\}, \quad (3.6)$$

where $J_i(k) = \frac{1}{2p_1 k} - \frac{1}{2q_1 k}$; D is the region of integration over the meson momenta limited by the "softness" conditions; f_{el} is the elastic scattering amplitude without taking into account radiation corrections.

We note that, in principle, for f_{el} it is possible to use the eikonal representation obtained in refs. 1-3, 6-9/.

Using (3.6) for the cross section for inelastic process with "soft" meson production it is not difficult to get the expression 1/

$$d\sigma_n = d\sigma_0 \frac{a_1^n}{n!} e^{-a_2}, \quad (3.7)$$

where

$$a_1 = \frac{g^2}{(2\pi)^3} \int \frac{d^3k}{2k_0} |J_1 + J_2|^2 \quad (3.8)$$

and

$$a_2 = \frac{g^2}{(2\pi)^3} \int \frac{d^3 k}{2k_0} [J_1^2 + J_2^2]. \quad (3.9)$$

are the contributions of the real and virtual meson, respectively, and

$$d\sigma_0 = |f_{el}|^2 d\Omega.$$

Since the interference term $J_1 J_2$ in the domain $\frac{t}{s} \ll 1$ gives smaller contribution than the quadrupole terms/1/ we have $a_1 = a_2 = a$ and the distribution over the secondary particles number takes the simple form

$$d\sigma_n = d\sigma_0 \frac{a^n}{n!} e^{-a}. \quad (3.10)$$

We notice that eq. (3.7) holds in a more general case. If we restrict ourselves to the assumption about the softness of only real mesons (this condition is necessary for the amplitude to be factorized), and perform the averaging over ν , according to eq. (3.5), then eq. (3.7) is valid. But in this case it is necessary to make the following substitutions

$$a_1 \rightarrow \bar{n}_{m.e.} = \frac{g^2}{(2\pi)^3} \int \frac{d^3 k}{2k_0} [J_1^2 + J_2^2], \quad (3.11)$$

where

$$J_1 = \frac{1}{\mu^2 + 2k p_1} - \frac{1}{-\mu^2 + 2k q_1}, \quad (3.12)$$

$$a_2 \rightarrow \bar{n}_{r.c.} = 2 \operatorname{Re} \left\{ \frac{g^2}{(2\pi)^4} \frac{i}{2} \int \frac{d^4 k}{k^2 - \mu^2 + i\epsilon} \left| \tilde{J}_1 + \tilde{J}_2 \right|^2 \right\}, \quad (3.13)$$

where

$$\tilde{J}_1 = \frac{1}{k^2 + 2kp_1} - \frac{1}{k^2 + 2kq_1}. \quad (3.14)$$

When the mesons are vector particles the quantities \tilde{J} and \tilde{J}^v read

$$\tilde{J}_{1\mu}^v = \frac{k_\mu + 2p_{1\mu}}{2p_1 k + k^2} + \frac{k^2 - 2q_{1\mu}}{2q_1 k + k^2}, \quad (3.15)$$

$$\tilde{J}_{1\mu}^v = \frac{k_\mu + 2p_{1\mu}}{2p_1 k + \mu^2} + \frac{k_\mu + 2q_{1\mu}}{2q_1 k - \mu^2}. \quad (3.16)$$

Thus, see that in quantum field models the n "soft" meson production cross section has the behaviour of the Poisson distribution.

4. Investigation of the Radiative Correction of Real Mesons Contributions to the Cross Section

We first consider the integral (3.13) with currents (3.14) and (3.15) which corresponds to the radiative correction contribution.

In the models in question the integration over the functional variables ν_1 and ν_2 , according to eq. (3.5), leads to a factorization of the radiative corrections in the scattering amplitude and the cross section. Then in the nucleon propagators the k^2 -dependence is kept and the integrals converge in the upper limit.

It is not difficult to see that the integral (3.13) in this approximation does not contain the s -dependence^{4/}.

In the asymptotic domain $|t| \ll m^2$ it is

$$\bar{n}_{r.c.} = t \frac{g^2}{24 (2\pi)^2 m^4} \ln\left(\frac{m^2}{\mu^2}\right) + O\left(\frac{t^2}{m^4}\right) \quad (4.1)$$

if the scalar current (3.14) is inserted in (3.13) and

$$\bar{n}_{r.c.}^v = t \frac{2g^2}{3(2\pi)^2 m^2} \left[\ln\left(\frac{m^2}{\mu^2}\right) + \frac{1}{2} \right] + O\left(\frac{t^2}{m^4}\right) \quad (4.2)$$

if the vector current (3.15) is used.

As was already noted in ref.^{1/}, the quantity $\frac{\bar{n}_{r.c.}^v}{t}$ is the width of the diffraction peak.

The softness conditions imposed on the meson momenta are as follows^{1,5/}

$$\frac{1}{\sqrt{s}} \sum_{i=1}^n k_{0i} \ll 1, \quad \left| \sum_{i=1}^n \vec{k}_{i\perp} \right| \ll |\vec{p}_{1\perp} - \vec{q}_{1\perp}| = |\vec{p}_{2\perp} - \vec{q}_{2\perp}|, \quad (4.3)$$

where the particle momentum components are given in the c.m.s., the initial nucleon momenta being chosen along the z -axis.

In order to cut off the integrals (3.11) with currents (3.12) and (3.16) in the upper limit it is natural to single out in the whole momentum space a cylindric domain oriented along the z -axis

$$|k_z| \leq \alpha p_0 = \epsilon_z, \quad (4.4)$$

$$|\vec{k}_\perp| \leq \epsilon_\perp \quad (4.5)$$

since in the majority of cases the experimentally observed secondary mesons are emitted forward.

As was shown in ref./1/ in the infrared asymptotic limit $\mu \rightarrow 0$ as well as in a broader region defined by the condition

$$\epsilon_\perp \approx m^2, \quad 1 \gg \alpha^2 \gg \frac{\mu^2}{m^2}, \quad \text{where} \quad \alpha \equiv \frac{\epsilon_z}{p_0} \quad (4.6)$$

$$\ln \left(\frac{\mu^2}{m^2} \right) \gg \ln \left(\frac{1}{\alpha^2} \right),$$

the absolute contributions from the radiative corrections and emitted mesons coincide^{x/}. In this case, in summing in the expression for $d\sigma_n$ over the number of all the emitted mesons the t -dependence vanishes. This leads to the peak being also vanishing in the diffraction cross section. A similar feature has been noted in ref./21/ and is analogous to the automodel behaviour of deep inelastic interactions of hadrons with leptons at high energies/22,23/.

^{x/}The results of estimation of the integrals for different domains are given in the Appendix.

Note that the quantity $\bar{n}_{m.e.}$ in the Poisson distribution has a simple physical meaning of the average number of emitted particles, i.e. multiplicity. Thus, in the region indicated, under the condition that the cutoff boundaries are energy independent, the multiplicity of "soft" particles turns out to be independent of the incident nucleon energy. As a result, the n soft particle production cross section at high energies is also independent of s .

As was noted in Introduction, in the framework of the present method there is a certain ambiguity in choosing the cutoff parameters and the result is sensitive to this choice. For example, if in the infrared domain the cutoff parameter ϵ_{\perp} is chosen to be increasing with energy $\epsilon_{\perp} = \beta s$ ($\beta \ll 1$ and fixed) then the multiplicity increases logarithmically with energy.

Let us consider the case when the μ scalar meson mass is fixed and is not small as compared with the nucleon mass m and $s \rightarrow \infty$. The average number of emitted particles in this domain is expressed as

$$\bar{n}_{m.e.} \sim a^4 \int d^2 \vec{k} \frac{(k_{\perp} \Delta_{\perp})}{(\mu^2 + k_{\perp}^2)^4} \quad (4.7)$$

It is seen that at a fixed cutoff parameter ϵ_z the multiplicity decreases with increasing energy $s \rightarrow \infty$. This means that with increasing the number of "soft" mesons, the momenta of which lie in a given interval defined by fixed parameter ϵ_z , decreases.

In addition, the condition may be imposed that the maximum longitudinal component of the meson momenta increases linearly with energy. In so doing, the average particles multiplicity tends with increasing energy to a finite limit. A similar result is obtained for the vector mesons.

Taking the opportunity the authors express their deep gratitude to N.N. Bogolubov, D.I. Blokhintsev, A.A. Logunov, A.N. Tavkhelidze

for stimulating discussions and valuable remarks. We thank also O.A. Khrustalev, V.I. Savrin, N. Ye. Tyirin and M.A. Smondyrev for interesting discussions.

R e f e r e n c e s

1. B.M. Barbashov, S.P. Kuleshov, V.A. Matveev, V.N. Pervushin, A.N. Sissakian, A.N. Tavkheldidze. *JINR Preprint*, E2-5217, Dubna (1970).
2. B.M. Barbashov, S.P. Kuleshov, V.A. Matveev, A.N. Sissakian. *JINR Preprint*, E2-4692, Dubna (1969).
3. Б.М. Барбашов, С.П. Кулешов, В.А. Матвеев, А.Н. Сисакян. *ТМФ*, 3, 342 (1970).
4. B.M. Barbashov, S.P. Kuleshov, V.A. Matveev, A.N. Sissakian. *JINR Preprint*, E2-4983, Dubna (1970).
5. B.M. Barbashov, S.P. Kuleshov, V.N. Pervushin, A.N. Sissakian. *JINR Preprint*, E2-4955, Dubna (1970).
6. H.D.I. Abarbanel, C. Itzykson. *Phys.Rev.Lett.*, 23, 53 (1969).
7. И.В. Андреев. *ЖЭТФ*, 58, 257 (1970).
8. M. Levy, J. Sucher. *Phys.Rev.*, 186, 1656 (1969).
9. H. Cheng, T.T. Wu. Report submitted to the XV-th International Conference on High Energy Physics., Kiev-1970.
10. Б.М. Барбашов. *ЖЭТФ*, 48, 607 (1965).
11. Г.А. Милехин, Е.С. Фрадкин. *ЖЭТФ*, 45, 1926 (1963).
12. Б.М. Барбашов, М.К. Волков. *ЖЭТФ*, 50, 660 (1966).
13. E.S. Fradkin. *Nucl.Phys.*, 76, 588 (1966).
14. Б.М. Барбашов, В.В. Нестеренко. Препринт ОИЯИ, P2-4900, Дубна, на (1970).
15. G. Tiktopoulos, S.B. Treiman. *Preprint Princeton University* (1970).
16. В.И. Саврин, Н.Е. Тюрин, О.А. Хрусталеv. Доклад, представленный на ХУМеждународную конференцию по физике высоких энергий. Киев-1970.

17. R.J. Glauber. *Phys.Rev.*, 84, 395 (1951).
18. G. Mack. *Phys.Rev.*, 154, 1617 (1967).
19. H.A. Kastrup. *Phys.Rev.*, 147, 1130 (1966).
20. В.Г. Горшков. *ЖЭТФ*, 56, 597 (1969).
21. V.A. Matveev, A.N. Tavkhelidze. *JINR preprint*, E2-5141, Dubna (1970).
22. В.А. Матвеев, Р.М. Мурадян, А.Н. Тавхелидзе. *Препринт ОИЯИ*, P2-4578, Дубна (1969).
23. A.N. Tavkhelidze. *Talk given at the Coral Gables Conference*, Miami (1970).
24. В.Н. Первушин. *ТМФ*, 4, 22 (1970).
25. Н.Н. Боголюбов, Д.В. Ширков. "Введение в теорию квантованных полей", ГИТТЛ, М (1957).

Received by Publishing Department
on August 27, 1970.

Appendix

Estimations of integrals, corresponding to real and virtual

"mesons"

Scalar "mesons"

$\bar{n}_{r.c.}$	$2 \operatorname{Re} \left\{ \frac{g^2}{(2\pi)^4} \frac{1}{2} \int \frac{d^4 k}{k^2 - \mu^2 + i\epsilon} \left[\frac{1}{k^2 + 2kp_1} - \frac{1}{k^2 + 2kq_1} \right] \right\}$	$t \frac{g^2}{24(2\pi)^4 m^4} \ln \left(\frac{m^2}{\mu^2} \right) + O \left(\frac{t^2}{m^4} \right)$	$t \frac{g^2}{24(2\pi)^4 m^4} \ln \left(\frac{m^2}{\mu^2} \right)$	$\bar{n}_{r.c.} = \bar{n}_{r.c.}(t)$	$\bar{n}_{r.c.}$
$\bar{n}_{m.e.}$	$\frac{g^2}{(2\pi)^3} \int_{ k_1 \leq p_1} \frac{dk_1}{2k_0} \int_{ k_2 \leq \epsilon_1} d^2 \vec{k}_2 \left[\frac{1}{2kp_1} - \frac{1}{2kq_1} \right]^2$	$-t \frac{g^2}{24(2\pi)^4 m^4} (A+B)$	$-t \frac{g^2}{24(2\pi)^4 m^4} \ln \left(\frac{\epsilon_1^2}{\mu^2} \right)$	$\frac{g^2}{(2\pi)^3} \int_{ k_2 \leq \epsilon_1} d^2 \vec{k}_2 \frac{(\vec{k}_2 \vec{\Delta}_1)^2}{(\mu^2 + k_2^2)^4}$	$\bar{n}_{m.e.}$
	General form of the integral taking into account only J_1	Asymptotic at $ t \ll m^2, (t < 0); \alpha \gg \frac{\epsilon_1}{p_0}$	Asymptotic at $ t \ll m^2, \alpha \gg \frac{\epsilon_1}{p_0}; \frac{\mu^2}{m^2} \ll \frac{m^2}{s}, \mu \rightarrow 0$	Asymptotic at $ t \ll m^2, \alpha \gg \frac{\epsilon_1}{p_0}; \frac{\mu^2}{m^2} \geq \frac{m^2}{s}; s \rightarrow \infty$	
$\bar{n}_{r.e.}$	$2 \operatorname{Re} \left\{ \frac{g^2}{(2\pi)^4} \frac{1}{2} \int \frac{d^4 k}{k^2 - \mu^2 + i\epsilon} \left[\frac{k_\mu + 2p_{1\mu}}{2p_1 k + k^2} - \frac{k_\mu + 2q_{1\mu}}{2q_1 k + k^2} \right] \right\}$	$t \frac{2g^2}{3(2\pi)^4 m^2} \left[\ln \left(\frac{m^2}{\mu^2} \right) + \frac{1}{2} \right] O \left(\frac{t^2}{m^2} \right)$	$t \frac{2g^2}{3(2\pi)^4 m^2} \ln \left(\frac{m^2}{\mu^2} \right)$	$\bar{n}_{r.e.} = \bar{n}_{r.e.}(t)$	$\bar{n}_{r.e.}$
$\bar{n}_{m.r.}$	$-\frac{g^2}{(2\pi)^3} \int_{ k_1 \leq p_0} \frac{dk_1}{2k_0} \int_{ k_2 \leq \epsilon_1} d^2 \vec{k}_2 \left[\frac{p_{1\mu}}{p_1 k} - \frac{q_{1\mu}}{q_1 k} \right]^2$	$-t \frac{g^2}{6(2\pi)^4 m^2} (4A+B)$	$t \frac{2g^2}{3m^2(2\pi)^2} \ln \left(\frac{\epsilon_1^2}{\mu^2} \right)$	$t \frac{2g^2}{(2\pi)^4} \int_{ k_2 \leq \epsilon_1} d^2 \vec{k}_2 \frac{1}{(\mu^2 + k_2^2)^2} + O(\alpha^2)$	$\bar{n}_{m.e.}$
Vector "mesons"					

$$A = \ln \left| 1 + \frac{c}{\mu^2} \right| - \ln \left| 1 + \frac{c}{\mu^2 + \epsilon_1^2} \right|; \quad B = -\frac{\epsilon_1^2 c [(\epsilon_1^2 + \mu^2 + c)(\mu^2 + c) + \epsilon_1^2(\epsilon_1^2 + \mu^2)]}{(\epsilon_1^2 + \mu^2)(\mu^2 + c)(\epsilon_1^2 + \mu^2 + c)^2}; \quad C = \frac{4p_0^2 d^2 (p_0 - p_1)}{p_0 + p_2}$$

Пуассоновский характер распределения вторичных частиц
в приближении прямолинейных путей

В рамках стандартных теоретико-полевых моделей рассмотрена проблема множественного рождения "мягких" мезонов при столкновении двух нуклонов высоких энергий. С помощью приближения прямолинейных путей получено пуассоновское распределение числа вторичных частиц, а также исследована зависимость среднего числа частиц от способов обрезания интегралов.

Сообщения Объединенного института ядерных исследований
Дубна, 1970

Barbashov B.M., Kuleshov S.P., Matveev V.A. E2-5329
Pervushin V.N., Sissakian A.N.

The Poisson Distribution of Secondary Particles
in Straight-Line Paths Approximation

The problem of multiple production of "soft" mesons in high-energy two-nucleon collision is considered in the framework of standard quantum field theory models. The straight-line paths approximation is used to obtain the Poisson distribution of the secondary particle number and to investigate the average particles as a function of the choice of the integral cutoff parameters in various asymptotic regions.

Communications of the Joint Institute for Nuclear Research.
Dubna, 1970