

ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

ДУБНА



G-42

E2 - 7711

8/IV-74

1332/2-74

A.Gheorghe , D.B.Ion , El.Mihul

ON RELATIVISTIC MULTIPARTICLE
KINEMATICS IN INVARIANT VARIABLES

1974


ЛАБОРАТОРИЯ
ТЕОРЕТИЧЕСКОЙ ФИЗИКИ

E2 - 7711

A.Gheorghe*, D.B.Ion*, El.Mihul*

**ON RELATIVISTIC MULTIPARTICLE
KINEMATICS IN INVARIANT VARIABLES**

Submitted to Ann. de l'Inst. H.Poincare,
sec. A.



* On leave of absence from the Institute
of Atomic Physics, Bucharest, Romania.

Георге А., Ион Д.Б., Михул Э.

E2 - 7711

О реалистической многочастичной кинематике в инвариантных переменных

Проведены полный анализ алгебраических ограничений на матрицы Грама, соответствующие любому конечному числу 4-импульсов, и аналитическое восстановление любой компоненты 4-импульсов через матрицу Грама.

Препринт Объединенного института ядерных исследований.
Дубна, 1974

Gheorghe A., Ion D.B., Mihul E.I.

E2 - 7711

On Relativistic Multiparticle Kinematics
in Invariant Variables

A complete system of algebraic constraints on the Gram matrices of four-momenta and an analytic reconstruction of any finite set of four-momenta from their Gram matrix are established. The physical regions and phase-space volume elements in general invariant variables of both exclusive and inclusive multiparticle processes are also given.

Preprint. Joint Institute for Nuclear Research.
Dubna, 1974

1. INTRODUCTION

The purpose of this work is to extend some previous analyses /1-6/ in relativistic multiparticle kinematics in invariant variables. The present approach is essentially inspired by certain old algebraic results of Jacobi, Hildenfänger, Frobenius, Gauss and Weyl.

The paper is organized as follows. In Section 2, we present two kinematical rules for the Gram matrices of finite sets of four-momenta. The first rule gives a complete system of Gram determinantal equalities and inequalities which guarantees that a real symmetric $n \times n$ matrix is a Gram matrix of n four-momenta. The second rule represents a concrete analytic reconstruction of any finite set of four-momenta from their Gram matrix.

In Section 3, we apply the above kinematical rules to a description of the physical regions of both exclusive and inclusive multiparticle processes in general invariant variables. As applied to this description, some algebraic and geometric properties of the considered physical regions are discussed. On the other hand, a convenient unified scheme for finding the ranges of several useful sets of invariant variables is established and the phase-space volume elements are determined with respect to this scheme.

2. KINEMATICAL RULES FOR GRAM MATRICES

We first establish a complete system of algebraic constraints on the Gram matrices of n four-momenta.

It is convenient to start with some notation. The four-momentum q is written as $q = (q^0, \vec{q}) = (q^0, q^1, q^2, q^3)$. The Minkowski scalar product of two four-momenta q and q' is given by $qq' = q^0q'^0 - \vec{q}\vec{q}'$ (with the notation $(q)^2 = qq$ for the Lorentz square of q).

The Gram matrix of the four-momenta q_1, \dots, q_n is defined by

$$U = \begin{pmatrix} (q_1)^2 & q_1 q_2 & \dots & q_1 q_n \\ q_1 q_2 & (q_2)^2 & \dots & q_2 q_n \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ q_1 q_n & q_2 q_n & \dots & (q_n)^2 \end{pmatrix}. \quad (1)$$

If $U = (u_{ij})$ is a real symmetric $n \times n$ matrix, the minor of U with all rows and columns deleted except for the i_1, \dots, i_h -th lines and j_1, \dots, j_h -th columns is denoted by

$$G_{i_1 \dots i_h}^{j_1 \dots j_h}(U) = \det(u_{ij})_{i=i_1, \dots, i_h; j=j_1, \dots, j_h} \quad (2)$$

We also use the following notations:

$$G_{i_1 \dots i_h}^{i_1 \dots i_h}(U) = G_{i_1 \dots i_h}^{i_1 \dots i_h}(U), \quad (3)$$

$$\sigma_{i_1 \dots i_h}(U) = \text{sgn}(-1)^{h-1} G_{i_1 \dots i_h}^{i_1 \dots i_h}(U), \quad (4)$$

where, for any real number α , $\text{sgn } \alpha = 1$ if $\alpha > 0$, $\text{sgn } \alpha = 0$ if $\alpha = 0$, and $\text{sgn } \alpha = -1$ if $\alpha < 0$.

Let us now state the following rule:

RULE 1. Let U be a real symmetric $n \times n$ matrix. Then the following statements are equivalent:

a) there exist n four-momenta such that U is their Gram matrix;

b) U has at most one positive eigenvalue and at most three negative eigenvalues;

c) $r = \text{rank } U \leq 4$ and in the case $r > 0$ there exist some indices $i_1, \dots, i_r \in \{1, \dots, n\}$ such that $\sigma_{i_1 \dots i_r}(U) \neq 0$ and

$$\sigma_{i_1 \dots i_h}(U) \leq \sigma_{i_1 \dots i_{h+1}}(U), \quad 1 \leq h < r, \quad (5)$$

$$\sigma_{i_1 i_2 i_3 i_4}(U) = 1 \quad \text{if } r = 4, \quad (6)$$

$$\sigma_{i_2}(U) = -1 \quad \text{if } r = 4, \quad \sigma_{i_1 \dots i_h}(U) = 0, \quad h = 1, 2, 3. \quad (7)$$

Proof. Several parts of the above rule are well-known /5/. Suppose that c) holds. Let ν denote the number of all indices h such that $\sigma_{i_1 \dots i_h}(U) = 0$; $1 \leq h < r$. According to a remark of Poon/5/, the implication c) \Rightarrow b) represents Jacobi's rule if $\nu = 0$ and Hildenfänger's rule if $\nu = 1$. Moreover, we recall that the implication c) \Rightarrow b) for $\nu = 2$ represents a rule of Frobenius (for Jacobi's, Hildenfänger's and Frobenius' rules see the full section 3 from /7/ Ch. X). The only remaining case is $\nu = 3$. Using (6) and the conditions of (7), we obtain $\sigma_{i_2}(U) = \sigma_{i_2 i_3}(U) = -1$ and $\sigma_{i_1 i_2 i_3 i_4}(U) = 1$. Then (5) holds if the indices i_1, i_2, i_3, i_4 are replaced by i_2, i_3, i_4, i_1 , respectively, and the case $\nu = 3$ is reduced to the cases $\nu = 0$ and $\nu = 1$. Hence c) implies b).

Suppose that b) holds. Then it follows from the canonical diagonalization of a real symmetric matrix that

there exist a real orthogonal $n \times n$ matrix $\Omega = (\omega_{ij})$ and a real diagonal $n \times n$ matrix $D = (d_{ij})$ such that $U = \Omega D \Omega^t$, $d_{11} \geq 0$, $d_{ii} \leq 0$, $d_{jj} = 0$, $2 \leq i \leq 4$, $j > 4$ (see, for example, /7/ Ch. X, § 5). The symbol t denotes the matrix transpose. Let us denote $q_i^{j-1} = \omega_{ij} |d_{jj}|^{1/2}$, $i = 1, \dots, n$; $j = 1, 2, 3, 4$. Then U is the Gram matrix of the four-momenta q_1, \dots, q_n . Hence b) implies a).

In order to prove the implication a) \Rightarrow c), we introduce certain orthogonal bases for the linear subspaces of the Minkowski space. The basis $\{e_1, \dots, e_d\}$ is called a standard basis if the four-momenta e_1, \dots, e_d satisfy the following relations:

$$e_h e_{h'} = \begin{cases} \epsilon & \text{if } h = h' = 1 \\ -1 & \text{if } h = h' > 1, \\ 0 & \text{if } h \neq h' \end{cases} \quad (8)$$

where $h, h' = 1, \dots, d$ and $\epsilon \in \{-1, 0, 1\}$. There exists a standard basis for any non-zero linear subspace N of the Minkowski space. Indeed, if N is d -dimensional, then there exists an orthogonal basis $\{f_1, \dots, f_d\}$ for N such that $(f_h)^2 \geq (f_{h+1})^2$, $h = 1, \dots, d-1$ /8/. It is easy to see that if two four-momenta are linearly independent and their scalar product is equal to zero, then at least one of these four-momenta has a strictly negative Lorentz square. Then we can define $e_1 = f_1$ if $(f_1)^2 = 0$, and $e_h = |(f_h)^2|^{-1/2} f_h$ if either $h > 1$ or $h = 1$ and $(f_1)^2 \neq 0$. Therefore $\{e_1, \dots, e_d\}$ is a standard basis for N . Notice that if $\epsilon \leq 0$, then $(q)^2 \leq 0$ for any $q \in N$. Hence $\epsilon = 1$ when $d = 4$ (in this case N is the whole Minkowski space of four-momenta). It is clear that $d \leq 4$.

Suppose that U is the Gram matrix of the four-momenta q_1, \dots, q_n . Let N be the linear space spanned by the four-momenta q_{i_1}, \dots, q_{i_h} , where $i_1, \dots, i_h \in \{1, \dots, n\}$. If $q_1 = \dots = q_n = 0$, then $U = 0$ and the implication a) \Rightarrow c) is trivial. Therefore we can

suppose that $d > 0$. Let $\{e_1, \dots, e_d\}$ be a standard basis for N . We introduce the $d \times d$ matrix $V = (e_j e_{j'})$, $1 \leq j, j' \leq d$, and the $h \times d$ matrix $T = (t_{kj})$, $1 \leq k \leq h$, $1 \leq j \leq d$, defined by

$$q_i = \sum_{j=1}^d t_{ij} e_j, \quad i = i_1, \dots, i_h. \quad (9)$$

The Gram matrix of the four-momenta q_{i_1}, \dots, q_{i_h} is $U' = TVT^t$. If q_{i_1}, \dots, q_{i_h} are linearly dependent, then it is obvious that $\sigma_{i_1 \dots i_h}(U) = (-1)^{h-1} \det U' = 0$. Moreover, we have $\text{rank } U' \leq 4$ and, in particular, $r = \text{rank } U \leq d \leq 4$.

If q_{i_1}, \dots, q_{i_h} are linearly independent, then $h = d$, $\det T \neq 0$, and (8) implies $\sigma_{i_1 \dots i_h}(U) = (e_1)^2$. In the case $r = h = 4$ one obtains $\sigma_{i_1 i_2 i_3 i_4}(U) = \det V = 1$ and (6) holds. If $r > 1$, we choose $i_1, \dots, i_{r'} \in \{1, \dots, n\}$, $1 < r' \leq r$, such that $\sigma_{i_1 \dots i_{r'}}(U) \neq 0$ and consider that $h \in \{1, \dots, r'-1\}$. Then the four-momenta $q_{i_1}, \dots, q_{i_{r'}}$ are linearly independent and the linear space N' spanned by $q_{i_1}, \dots, q_{i_{h+1}}$ admits a standard basis $\{e'_1, \dots, e'_{h+1}\}$ with $\sigma_{i_1 \dots i_{h+1}}(U) = (e'_1)^2$. Since N is a subspace of N' , e_1 is a linear combination of e'_1, \dots, e'_{h+1} and (8) implies $(e'_1)^2 \geq (e_1)^2$. Hence (5) holds for the choice $r = r'$.

Suppose next that $r = r' > 2$ and $\sigma_{i_1}(U) = \sigma_{i_1 i_2}(U) = 0$. It is clear that $(q_{i_1})^2 = q_{i_1} q_{i_2} = 0$. Since q_{i_1} and q_{i_2} are linearly independent, we have $(q_{i_2})^2 < 0$ and (7) also holds. Hence a) implies c) and RULE 1 is proved.

Let I_n denote the space of all Gram matrices of n four-momenta. Any Gram matrix U satisfies both assertions b) and c) of Rule 1. Moreover, it follows from the proof of the implication a) \Rightarrow c) that

$$\sigma_{i_1 \dots i_h}(U) \leq \sigma_{i_1 \dots i_{h+1}}(U), \quad h = 1, \dots, r'-1, \quad (10)$$

for any $i_1, \dots, i_{r'} \in \{1, \dots, n\}$ such that $\sigma_{i_1 \dots i_{r'}}(U) \neq 0$.

The space I_n is completely defined by the polynomial equalities and inequalities given by (5) - (7) and the condition $\text{rank } U \leq 4$. If $n > 4$, the last condition is equivalent to the following kinematical constraints of Asribekov /2/:

$$G_{i_1 i_2 i_3 i_4 i_5}^{j_1 j_2 j_3 j_4 j_5}(U) = 0, \quad (11)$$

where $1 \leq i_1 < \dots < i_5 \leq n$ and $1 \leq j_1 < \dots < j_5 \leq n$.

According to a result of Weyl, any identity with respect to the coefficients of U is a consequence of (11) (see, for example, the full section 17 from /9/, Ch. II). Moreover, according to a result of Kronecker /10/, the condition $\text{rank } U = r$ is equivalent to the following relations:

$$\sum_{1 \leq j_1 < \dots < j_h \leq n} G_{j_1 \dots j_h}^{i_1 \dots i_h}(U) = 0, \quad h = r+1, \dots, n, \quad (12)$$

$$G_{i_1 \dots i_r}(U) \neq 0$$

for some $i_1, \dots, i_r \in \{1, \dots, n\}$.

Hence a real symmetric $n \times n$ matrix U which satisfies (12) for $r \leq 4$ and the relation $G_{i_1 \dots i_r}(U) \neq 0$ for a choice of the indices $i_1, \dots, i_r \in \{1, \dots, n\}$ is a Gram matrix of n four-momenta if and only if (5), (6), and (7) hold.

We recall that Rule 1 for any Gram matrix with positive diagonal coefficients has been obtained by Omnes /1/ and Byers and Yang /3/.

The reconstruction of a set of four-momenta from their Gram matrix is given by the following rule:

RULE 2. Consider a Gram matrix $U \in I_n$ with $\text{rank } U = r > 0$. Then there exists a choice of the indices $i_1, \dots, i_r \in \{1, \dots, n\}$ such that either $\sigma_{i_1 \dots i_h}(U) \neq 0$, $h = 1, \dots, r$, or there exists $k \in \{1, \dots, r-1\}$ satisfying the following conditions:

1) $\sigma_{i_1 \dots i_h}(U) = -1$ if $1 \leq h < k$; 2) $\sigma_{i_1 \dots i_h}(U) = 1$ if

$k < h \leq r$; 3) $\sigma_{i_1 \dots i_{k-1} i_h}(U) = 0$ if $k \leq h \leq r$;

4) $G_{i_1 \dots i_{k-1} i_{k+1}}^{i_1 \dots i_{k-1} i_{k+1}}(U) \neq 0$; 5) $i_k < i_{k+1}$ if and only if

$$(-1)^k G_{i_1 \dots i_{k-1} i_k}^{i_1 \dots i_{k-1} i_{k+1}}(U) > 0.$$

Consider an orthogonal $n \times n$ matrix $\hat{O} = (o_{jj'})$ such that either $\hat{O} = 1$ for $\sigma_{i_1 \dots i_h}(U) \neq 0$, $h = 1, \dots, r$, or

$$o_{jj'} = \begin{cases} 1 & \text{if } j = j' \neq i_k, i_{k+1} \\ 0 & \text{if } j \neq j' \text{ and either } j \neq i_k, i_{k+1} \\ & \text{or } j' \neq i_k, i_{k+1} \\ -1/\sqrt{2} & \text{if } j < j' \text{ and } j, j' \in \{i_k, i_{k+1}\} \\ 1/\sqrt{2} & \text{if } j \geq j' \text{ and } j, j' \in \{i_k, i_{k+1}\} \end{cases} \quad (13)$$

for $\sigma_{i_1 \dots i_k}(U) = 0$.

Then the matrix $U' = \hat{O} U \hat{O}^t$ satisfies the relations $\sigma_{i_1 \dots i_h}(U') \neq 0$, $1 \leq h \leq r$. Moreover, U' is the Gram matrix of the four-momenta q_1, \dots, q_n defined by

$$q_i = \sum_{j=1}^n o_{ji} q'_j, \quad i = 1, \dots, n, \quad (14)$$

$$q'_j = \begin{cases} \frac{\sigma_{i_1 \dots i_h}(U') G_{i_1 \dots i_{h-1} j}^{i_1 \dots i_{h-1} j}(U')}{|G_{i_1 \dots i_{h-1}}(U') G_{i_1 \dots i_h}(U')|^{1/2}} & \text{if } h \leq r \\ 0 & \text{if } h > r \end{cases} \quad (15)$$

where $j = 1, \dots, n$; $h = 1, \dots, 4$, and the following convention is used:

$-\sigma_{i_1 \dots i_{h-1}}(U') = G_{i_1 \dots i_{h-1}}(U') = 1$ if $h = 1$. Here

$(\mu_0, \mu_1, \mu_2, \mu_3)$ is a permutation of $(0, 1, 2, 3)$ such that $\mu_h = 0$ if $\sigma_{i_1 \dots i_{h-1}}(U') \cdot \sigma_{i_1 \dots i_h}(U') = -1$ and $\mu_h < \mu_{h'}$ if $h < h'$; $\mu_h, \mu_{h'} \neq 0$.

Proof. Since $r > 0$, we can choose the indices $j_1, \dots, j_r \in$

$\in \{1, \dots, n\}$ such that $G_{j_1 \dots j_r}(U) \neq 0$ and $\sigma_{j_1 \dots j_h}(U) \geq \sigma_{j_1 \dots j_h}(U)$ for $h \geq h'$. We now show that there exists a permutation (i_1, \dots, i_r) of (j_1, \dots, j_r) which satisfies the conditions of Rule 2. According to Rule 1c), the argument divides into two cases:

Case I. Suppose that $\sigma_{j_1 \dots j_h}(U) \neq 0$, $h = 1, \dots, r$. Then choose $i_h = j_h$, $h = 1, \dots, r$.

Case II. Suppose that there exists $k \in \{1, \dots, r-1\}$ such that $\sigma_{j_1 \dots j_h}(U) = -1$ if $1 \leq h < k$ and $\sigma_{j_1 \dots j_{k-1} j_h}(U) = 0$ if $k \leq h \leq r$. Since $G_{j_1 \dots j_r}(U) \neq 0$ it follows that

there exist two indices $h', h'' \in \{k, \dots, r\}$ such that

$G_{j_1 \dots j_{k-1} j_{h'}}(U) \neq 0$ and $j_{h'} < j_{h''}$. We choose $i_h = j_h$

if $h \in \{1, \dots, r\}$; $h', h'' \neq h$, and $i_k = j_{h'}, i_{k+1} = j_{h''}$

(resp. $i_k = j_{h''}, i_{k+1} = j_{h'}$) if $(-1)^k G_{j_1 \dots j_{k-1} j_{h'}}(U)$

is negative (resp. positive). Note that from the identity

$$\begin{aligned} G_{i_1 \dots i_{k-1}}(U) G_{i_1 \dots i_{k+1}}(U) &= \\ &= G_{i_1 \dots i_k}(U) G_{i_1 \dots i_{k-1} i_{k+1}}(U) - [G_{i_1 \dots i_{k-1} i_{k+1}}(U)]^2 \end{aligned} \quad (16)$$

it follows that $\sigma_{i_1 \dots i_{k+1}}(U) = 1$. Then (5) implies $\sigma_{i_1 \dots i_h}(U) = 1$ if $k < h \leq r$.

If k is as in Case II, using (13), we obtain

$$(-1)^{k-1} G_{i_1 \dots j_k}(U') = |G_{i_1 \dots i_{k-1} i_{k+1}}(U)|, \quad (17)$$

$$G_{i_1 \dots i_h}(U') = G_{i_1 \dots i_h}(U), \quad 1 \leq h \leq r, \quad h \neq k. \quad (18)$$

Therefore $\sigma_{i_1 \dots i_h}(U') \neq 0$, $1 \leq h \leq r$. Straightforward computations starting from (15) show that $U' = (q'_i q'_{j'})$; $j, j' = 1, \dots, n$ (see, for example, the Gauss algorithm from [7], Ch. II, § 4). Then by (14), we obtain $U = (q_i q_{j'})$; $i, i' = 1, \dots, n$, and Rule 2 is proved.

We remark that the implication c) \Rightarrow a) of Rule 1 follows from the proof of Rule 2. Then from the proof of Rule 1 excepting the implication c) \Rightarrow b), we obtain a direct proof of Jacobi's, Hildenfänger's and Frobenius' rules.

Let I_{nr} denote the set of all Gram matrices of rank r belonging to I_n . Consider the set of all matrices $U \in I_{nr}$ such that $G_{i_1 \dots i_r}(U) \neq 0$ for a choice of the indices $i_1, \dots, i_r \in \{1, \dots, n\}$ (in the case $r > 0$). Then Rule 2 shows that $\{u_{i_h j}\}$, $h = 1, \dots, r$; $j \in \{1, \dots, n\}$; $j \neq i_{h'}$, for $h' < h$, is a set of $nr - r(r-1)/2$ independent kinematical variables with respect to I_{nr} . According to (15) and the identities

$$[G_{i_1 \dots i_{r-1} j}(U')]^2 = G_{i_1 \dots i_r}(U') G_{i_1 \dots i_{r-1} j}(U'), \quad (19)$$

we can choose the independent variables u_{jj} , $u_{i_h j}$ with $h = 1, \dots, r-1$; $j \in \{1, \dots, n\}$; $j \neq i_{h'}$, if $h' \leq h$. But the Gram matrix U can be reconstructed from these

variables only if the signs of $G_{i_1 \dots i_{r-1} j}(U')$, $1 \leq j \leq n$, $j \neq i_1, \dots, i_r$, are known (see also Röhrlich's dichotomy from [4]).

Finally, any representative of a Gram matrix is given by the following completion of Rule 2:

RULE 2'. For any matrix $U \in I_n$ and four-momenta q'_1, \dots, q'_n such that U is their Gram matrix, there exists a standard basis $\{e_1, \dots, e_d\}$ such that one of the following decompositions holds:

$$q'_i = \sum_{h=1}^r \alpha_{ih} e_h, \quad i = 1, \dots, n, \quad r = d, \quad (20)$$

$$q'_i = \sum_{h=2}^{r+1} \alpha_{ih} e_h + \beta_i e_1, \quad (e_1)^2 = 0, \quad (21)$$

$$\sum_{i=1}^n \beta_i^2 = 1, \quad r = d-1 \leq 2,$$

where d ($0 \leq d \leq 4$) is the dimension of the linear space spanned by q'_1, \dots, q'_n ; r is the rank of U , each β_i is a real number, $\alpha_{ih} = q_i^{\mu_{h-1}}$ for $r = d$ and $\alpha_{ih} = q_i^{\mu_{h-2}}$ for $r = d-1$ with q_1, \dots, q_n given by Rule 2. The sums from the r.h.s. of (19) and (20) are dropped if $r = 0$.

Proof. We recall a result of Hall and Wightman^{/10/} if two sets of n four-momenta $\{q_1, \dots, q_n\}$ and $\{q'_1, \dots, q'_n\}$ have the same Gram matrix U , then there exist a Lorentz transformation Λ , some real numbers $\beta'_1, \dots, \beta'_n$ and a four-momentum ω such that

$$q'_i = \Lambda q_i + \beta'_i \omega, \quad (\omega)^2 = \omega(\Lambda q_i) = 0, \quad i = 1, \dots, n. \quad (22)$$

Here Λ is identified to the matrix (Λ^μ_ν) , $0 \leq \mu, \nu \leq 3$, such that the four-vectors $f_h = (\Lambda^0_{\mu_{h-1}}, \Lambda^1_{\mu_{h-1}}, \Lambda^2_{\mu_{h-1}}, \Lambda^3_{\mu_{h-1}})$

$h = 1, 2, 3, 4$, satisfy the relations $(f_1)^2 = -(f_2)^2 = -(f_3)^2 = -(f_4)^2 = 1$, $f_h f_{h'} = 0$, $1 \leq h' < h \leq 4$, with the permutation $(\mu_0, \mu_1, \mu_2, \mu_3)$ of $(0, 1, 2, 3)$ given by Rule 2 with respect to U . Note that the four-vectors f_1, f_2, f_3 and f_4 are linearly independent. Λq is

defined by $(\Lambda q)^\mu = \sum_{\nu=0}^3 \Lambda^\mu_\nu q^\nu$, $0 \leq \mu \leq 3$, for any

four-momentum q .

Suppose that the four-momenta q_1, \dots, q_n are determined by U as in Rule 2 (in the case $r > 0$). Then by (22), we have

$$q'_i = \sum_{h=1}^r q_i^{\mu_{h-1}} f_h + \beta'_i \omega, \quad i = 1, \dots, n, \quad (23)$$

where the sum over h is dropped if $r = 0$. If $\omega = 0$ or $\beta'_i = 0$ for any $i = 1, \dots, n$, we choose $e_h = f_h$, $h = 1, \dots, r$, and (20) holds. In the opposite case, we set $e_1 = (\sum_{i=1}^n \beta_i^2)^{1/2} \omega$

and $e_h = f_{h+1}$, $h = 1, \dots, r$. According to the proof of Rule 1 a standard basis $\{e_1, \dots, e_d\}$ with $(e_1)^2 = 0$ has $d \leq 3$. Then (21) holds and Rule 2' is proved.

Rules 2 and 2' give an analytical parametrization of any n four-momenta with respect to their scalar products, the parameters of the Lorentz group, and the coordinates of an n -sphere (not all independent).

3. PHYSICAL REGIONS AND PHASE SPACES

Consider a set of n particles c_1, \dots, c_n . By the spectral condition, we attach to each particle c_i a four-momentum p_i such that

$$(p_i)^2 = m_i^2, \quad p_i^0 > 0, \quad i = 1, \dots, n, \quad (24)$$

where m_i ($m_i \geq 0$) is the mass of c_i .

In order to give a general treatment for the usual choices of invariant kinematical variables (like the multiperipheral momentum transfers squared and multiparticle invariant masses) we introduce the following four-momenta:

$$q_i = \sum_{j=1}^n t_{ij} p_j, \quad i = 1, \dots, n, \quad (25)$$


where the coefficients t_{ij} are appropriate real numbers or functions of the scalar products $q_i q_j$, $1 \leq i, j \leq n$.

By (24), the four-momenta q_i satisfy (25) for some t_{ij} only if either all scalar products of q_i vanish or the linear space spanned by q_1, \dots, q_n has a standard basis $\{e_1, \dots, e_r\}$ with $(e_1)^2 = 1$. Then it follows from (20) that r is the rank of the Gram matrix U of q_1, \dots, q_n and


$$\sigma_{i_1 \dots i_r}(U) = 1 \quad (26)$$

if $G_{i_1 \dots i_r}(U) \neq 0$ and $i_1, \dots, i_r \in \{1, \dots, n\}$. Moreover, according to (20) and to the Sylvester law of inertia, (26) holds if and only if U has one positive eigenvalue (more details can be found in ^{5/}). We denote by I_{n+} the space of all Gram matrices $U \in I_n$ such that either $U=0$ or U satisfies (26).

Consider now the exclusive reaction



and the inclusive reaction



We suppose that the above reactions are not forbidden. The condition of energy-momentum conservation gives $p = p_n$ for (27) and $p^0 \geq 0$, $\sqrt{(p)^2} \in S$ for (28), where $p = p_1 + p_2 - p_3 - \dots - p_{n-1}$ and S is the spectrum of the invariant mass of the unobserved system from (28).

Suppose next that the $(n-1) \times (n-1)$ matrix $T = (t_{ij})$, $1 \leq i, j \leq n-1$, given by (25), is invertible. According to (24) and (25) and using the Gram matrix U of the four-momenta q_1, \dots, q_{n-1} , we obtain the following image of the physical region for reaction (27) (resp. (28)) in the space I_{n-1+} :

$$D_n = \{U | U = TU'T^t \in I_{n-1+}, U'_{ii} = m_i^2, \sum_{i,j=1}^{n-1} \epsilon_i \epsilon_j U'_{ij} = m_n^2, \sum_{j=1}^{n-1} \epsilon_j U'_{ij} \leq 0, \quad (29)$$

$$U'_{ij} \geq 0, \epsilon_1 = \epsilon_2 = -1, \epsilon_3 = \dots = \epsilon_{n-1} = 1; i, j = 1, \dots, n-1\},$$

respectively

$$D'_n = \{U | U = TU'T^t \in I_{n-1+}, U'_{ii} = m_i^2, [\sum_{i,j=1}^{n-1} \epsilon_i \epsilon_j U'_{ij}]^{1/2} \in S, \quad (30)$$

$$U'_{ij} \geq 0, \sum_{j=1}^{n-1} \epsilon_j U'_{ij} \leq 0, \epsilon_1 = \epsilon_2 = -1, \epsilon_3 = \dots = \epsilon_{n-1} = 1; i, j = 1, \dots, n-1\}.$$

The region D_n can be decomposed into the subregions $D_{nr} = D_n \cap I_{n-1r}$ (each D_{nr} consists of all matrices $U \in D_n$ with $\text{rank} U = r$). By Rule 1, we have $r \leq r_0 = \min(4, n-1)$. The case $r=0$ holds only when all masses vanish and all four-momenta are parallel. In the case $T=1$, the results of Jacobson ^{6/} show that each nonvoid subregion D_{nr} is a connected real analytic manifold of dimension $n(r-1) - r(r+1)/2$, $1 < r \leq r_0$. D_{n1} is not empty if and only if $U_{ij} = m_i m_j, 1 \leq i, j \leq n$, and $m_1 + m_2 = m_3 + \dots + m_n \neq 0$. Moreover, the inverse image of D_n in the space of all n -tuples of four-momenta is an analytic manifold if and only if D_{nr} is empty for $r \leq 1$. These results persist if the coefficients of T are analytic functions of the coefficients of U . Note that the closure \bar{D}_{nr} of D_{nr} , $1 < r \leq r_0$, is not an analytic manifold, but it is a semi-algebraic variety if the coefficients of T are polynomials of the coefficients of U (i.e., the set \bar{D}_{nr} consisting of all matrices $U \in D_n$ with $\text{rank} U \leq r$ is completely defined by the polynomial inequalities and equalities given by Rule 1 c) and (26)). Notice also that the boundary of the physical region D_n ($n \geq 4$) is \bar{D}_{nr-1} and the boundary of D_{nr} is \bar{D}_{nr-1} ($r > 2$). Finally, we remark that the physical regions of all channels crossed to (27) can be obtained by replacing $\epsilon_1, \dots, \epsilon_{n-1}$ in (29) by $\pm \epsilon_1, \dots, \pm \epsilon_{n-1}$ except for the cases $\epsilon_1 = \dots = \epsilon_{n-1}$ and $\epsilon_i = -\epsilon_j$, $j \in \{1, \dots, n-1\}$, $j \neq i$, for $i \in \{1, \dots, n-1\}$ fixed with the mass m_j smaller than the sum of the masses m_j , $j \neq i$ (i.e., the decay channel of particle c_i).

The region D'_{n-1} can be decomposed into the regions $D'_{n-1}(M)$, $M \in S$, where each $D'_{n-1}(M)$ is defined by the r.h.s. of (29) with m_n replaced by M . Hence any region $D'_{n-1}(M)$ has the same algebraic and geometric properties as D_n . Moreover, a similar behaviour persists for the

union D''_{n-1} of the regions $D'_{n-1}(M)$ with $M > M_c$, where M_c is the threshold mass of the continuum. Thus if $r > 1$, the region $D''_{n-1r} = D''_{n-1} \cap I_{n-1r}$ (resp. its closure \bar{D}_{n-1r}) is a connected real analytic manifold of dimension $n(r-1) - r(r+1)/2 + 1$ (resp. a semialgebraic variety) provided the coefficients of T are analytic functions (resp. polynomials) of the coefficients of U . The closure \bar{D}''_{n-1} of D''_{n-1} is the union of $D_{n-1}(M_c)$ and D''_{n-1} . \bar{D}''_{n-1r} consists of all matrices $U \in \bar{D}''_{n-1}$ with rank $U \leq r$. Notice that the boundaries of D''_{n-1} and \bar{D}''_{n-1} ($n \geq 4$) consist of all matrices $U \in \bar{D}''_{n-1}$ such that either $U \in D'_{n-1}(M_c)$ or rank $U < \min(4, n-1)$.

We now digress a little on phase-space analysis. The phase-space volume element of reaction (27) is defined by

$$dW(P, m_3, \dots, m_n) = \delta^4(P - \sum_{i=3}^n p_i) \prod_{j=3}^n \theta(p_j^0) \delta((p_j^2 - m_j^2)) d^4 p_j, \quad (31)$$

where $P = p_1 + p_2$ and the symbols δ and θ denote the usual Dirac and Heaviside distributions.

Let us consider the following transformation of variables:

$$p_i \rightarrow \{p_i q_i, p_i q'_i, p_i q''_i, (p_i)^2\}, \quad i = 3, \dots, n-1, \quad (32)$$

where the four-momenta p_i, q_i, q'_i and q''_i are linearly independent and the four-momenta q_i, q'_i and q''_i are differentiable functions of $p_j, 1 \leq j \leq i$. Then the Jacobian of (32) is

$$J = 2^{3-n} \prod_{i=3}^{n-1} [-D(p_i, q_i, q'_i, q''_i)]^{-1/2}, \quad (33)$$

Here and in the remainder of this paper we shall use the notation $D(Q_1, \dots, Q_h)$ for the determinant of the Gram matrix of Q_1, \dots, Q_h .

Notice that if

$$\gamma_i = \text{sgn det} \begin{vmatrix} p_i^0 & p_i^1 & p_i^2 & p_i^3 \\ q_i^0 & q_i^1 & q_i^2 & q_i^3 \\ q'_i{}^0 & q'_i{}^1 & q'_i{}^2 & q'_i{}^3 \\ q''_i{}^0 & q''_i{}^1 & q''_i{}^2 & q''_i{}^3 \end{vmatrix} \quad (34)$$

is fixed at a nonzero value ($\gamma_i = \pm 1$), then (32) is a one-to-one transformation.

Suppose that

$$D(-q_3, q'_3) < 0, \quad q_{n-1} = P - \sum_{i=3}^{n-2} p_i, \quad n > 4. \quad (35)$$

Using (32) - (35), the relation

$$\int d(p_3 q_3) [-D(p_3, q_3, q'_3, q''_3)]^{-1/2} = \pi [-D(q_3, q'_3)]^{-1/2}, \quad (36)$$

and integrating (31) over $p_n, (p_i)^2, p_3 q_3$, and $p_{n-1} \cdot q_{n-1}$ ($i=3, \dots, n$), the phase-space volume element can be written as

$$d\tilde{W}(P, m_3, \dots, m_n) = \chi \pi [-D(q_3, q'_3)]^{-1/2} \prod_{i=4}^{n-1} [-D(p_i, q_i, q'_i, q''_i)]^{-1/2} \times \prod_{j=3}^{n-2} d(p_j q_j) \prod_{j'=3}^{n-1} d(p_{j'}, q_{j'}) \prod_{j''=4}^{n-1} d(p_{j''}, q_{j''}), \quad (37)$$

where $(p_i)^2 = m_i^2, 3 \leq i \leq n-1$, and

$$p_{n-1} q_{n-1} = \frac{1}{2} [(q_{n-1})^2 + m_{n-1}^2 - m_n^2]. \quad (38)$$

The function χ gives the ranges of the invariant variables. Thus, if $q_i^0 > 0, (q_i)^2 \geq 0$, Rule 1 c) implies

$$\chi = \theta((q_{n-1})^2 - (m_{n-1} + m_n)^2) \prod_{i=3}^{n-2} \theta(p_i q_i - m_i \sqrt{(q_i)^2}) \times$$

$$\times \prod_{j=3}^{n-1} \theta(D(p_j, q_j, q'_j)) \prod_{k=4}^{n-1} \theta(-D(p_k, q_k, q'_k, q''_k)). \quad (39)$$

Note that if a function invariant to the connected Lorentz group is integrated with respect to the phase-space volume element (31), then the factor $2^{3-n} [\theta(\gamma_3) + \theta(-\gamma_3)] \dots [\theta(\gamma_{n-1}) + \theta(-\gamma_{n-1})]$ must be included in χ . However, this factor is not necessary for the estimation of cross-sections with parity conservation.

If $(P)^2$ is fixed, we have $3n-11$ essential invariant variables $p_j, q_j, p_{j'}, q_{j'}, p_{j''}, q_{j''}, 3 \leq j \leq n-2, 3 \leq j' \leq n-1, 4 \leq j'' \leq n-1$. By Rule 2, the scalar products of $q_3, q'_3, q_i, q'_i, q''_i (4 \leq i \leq n-1)$ are functions of the above variables. It is easy to estimate these functions in the scheme of Byers and Yang ^{/3/} ($q_3 = q'_3 = P; q_j = p_3 + \dots + p_{j-1}; q'_3 = p_1; q''_k = p_{k-1}; 4 \leq i, k \leq n-1; 4 \leq j \leq n-2$) or in the scheme of Poon ^{/5/} (i.e. the following multiperipheral scheme: $q'_j = p_{j-1}; q_3 = P; q_i = P - p_3 - \dots - p_{i-1}; q''_k = p_2 - p_3 - \dots - p_{k-1}, 4 \leq i \leq n-2; 3 \leq j \leq n-1; 4 \leq k \leq n-1$). Notice that (37) is a simple generalization of the phase-space volume element of Byers and Yang ^{/3/} (see ^{/11-13/} for some applications to the results of Byers and Yang). Note also that (37) unifies many usual schemes ^{/14/}.

Finally, we remark that the phase-space volume element of reaction (28) excepting the discrete part of the spectrum is given by $d\tilde{W}(P, m_3, \dots, m_{n-1}, M) dM^2; M \geq M_c$.

ACKNOWLEDGEMENTS

The authors are pleased to thank Prof. A.L.M. Mihul for many useful discussions and Mr. S. Berceanu for information on several aspects of inclusive kinematics.

REFERENCES

1. R. Omnes. Dispersion Relations and Elementary Particles, Les Houches, 1960; New York, Wiley, 1960, p. 341..
2. V.E. Asribekov. Phys. Lett., 2, 284 (1962).
3. N. Byers and C.N. Yang. Rev. Mod. Phys., 36, 595 (1964).
4. F. Röhrlich. Nuovo Cimento, 38, 673 (1965).
5. C.H. Poon. Nucl. Phys., B20, 509 (1970).
6. D.A. Jacobson. Nuovo Cimento, 45A, 905 (1966).
7. F.R. Gantmacher. The Theory of Matrices, Nauka, Moscow, 1966.
8. S. Lang. Algebra, Reading, Mass., Addison-Wesley Publ. Co., 1965, p. 365.
9. H. Weyl. The Classical Groups, Their Invariants and Representations, Princeton N.J., Princeton University Press, 1939.
10. D. Hall and A.S. Wightman. Dan. Vid. Selsk. Mat. Fys. Medd., 31, No. 5 (1959).
11. A.R. Morrow. Ann. Phys., 57, 115 (1970).
12. P. Nyborg and O. Skeggestad. Notes on Phase Space in Kinematics and Multiparticle Systems, London, Gordon and Breach, 1968.
13. E. Byckling and K. Kajantie. Particle Kinematics, New York, Wiley, 1973.
14. El. Mihul. Multiparticle Reactions and Lorentz Invariants (to be published); presented at the CERN/JINR School of Physics, Loma-Koli, 1970.

Received by Publishing Department
on February 1, 1974.