

ОБЪЕДИНЕННЫЙ  
ИНСТИТУТ  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ  
ДУБНА

2690/82

7/6-82

E2-82-159

A.V.Radyushkin

OPTIMIZED LAMBDA-PARAMETRIZATION  
FOR THE QCD RUNNING COUPLING  
CONSTANT  
IN SPACELIKE AND TIMELIKE REGIONS

Submitted to "Physics Letters B".

1982

## 1. INTRODUCTION


Perturbative QCD is intensively applied now to various processes involving large momentum transfers, both in spacelike ( $q^2 = -Q^2 < 0$ ) and timelike ( $q^2 > 0$ ) regions (for a review see <sup>1,3/</sup>). However, the coupling constant  $g(\mu)$  (i.e., the expansion parameter) is defined usually with the reference to some Euclidean (spacelike) configuration of momenta of scale  $\mu$ . For spacelike  $q$  this produces no special complications. One simply uses the renormalization group to sum up the logarithmic corrections  $(g^2(\mu) \ln(Q^2/\mu^2))^N$  that appear in higher orders and arrives at the expansion in the effective coupling constant  $\alpha_s(Q^2)$  which in the lowest approximation is given by the famous asymptotic freedom formula <sup>1/</sup>.

$$\alpha_s(Q^2) = \frac{4\pi}{(11-2N_f/3) \ln(Q^2/\Lambda^2)}, \quad (1)$$

where  $\Lambda$  is the "fundamental" scale of QCD. In general, the  $\Lambda$ -parametrization of  $\alpha_s(Q^2)$  is a series expansion in  $1/L$  (where  $L = \ln(Q^2/\Lambda^2)$ ), and the definition of  $\Lambda$  is fixed only if the  $O(1/L^2)$ -term is added to eq. (1) <sup>4/</sup>.

For timelike  $q$  there appear, however,  $i\pi$ -factors ( $\ln(Q^2/\mu^2) \rightarrow \ln(q^2/\mu^2) \pm i\pi$ ), and it is not clear a priori what is the effective expansion parameter in this region. This problem has been discussed recently in a very suggestive paper by Pennington and Ross <sup>5/</sup>. These authors analysed the ratio  $R(q^2) = \sigma(e^+e^- \rightarrow \text{hadrons}) / \sigma(e^+e^- \rightarrow \mu^+\mu^-)$  for which the analytic continuation from the spacelike to timelike region is well-defined and investigated which of the three ansätze ( $\alpha_s(q^2)$ ,  $|\alpha_s(-q^2)|$  and  $\text{Re}\alpha_s(-q^2)$ ) better absorbs the  $(\pi^2/L^2)^N$ -corrections \* in the timelike region  $q^2 > 0$ . Their conclusion was that  $|\alpha_s(-q^2)|$  is better than  $\alpha_s(q^2)$  and  $\text{Re}\alpha_s(-q^2)$ . Nevertheless, it is easy to demonstrate by a straightforward calculation that  $|\alpha_s(-q^2)|$  cannot absorb all the  $(\pi^2/L^2)^N$ -terms associated with the analytic continuation of the  $\ln(Q^2/\mu^2)$ -factors. Our main goal in the present letter is to show that by using the  $\Lambda$ -parametrization for

\* Odd powers of  $(i\pi/L)$  cancel because  $R$  is real.



$\alpha_s(Q^2)$  in the spacelike region it is possible to construct for  $R(Q^2)$  in the timelike region the expansion in which all the  $(\pi^2/L^2)^N$  -terms are summed up explicitly.

## 2. $\Lambda$ -PARAMETRIZATION IN SPACELIKE REGION

The starting point for the  $\Lambda$ -parametrization is the Gell-Mann-Low equation taken as a series expansion in  $G = \alpha_s/4\pi$ :

$$L \equiv \ln(Q^2/\Lambda^2) = \frac{1}{b_0 G} + \frac{b_1}{b_0^2} \ln G + \Delta + \frac{b_2 b_0 - b_1^2}{b_0^3} G + O(G^2), \quad (2)$$

where  $b_k$  are  $\beta$ -function coefficients:  $b_0 = 11 - 2N_f/3$ ,  $b_1 = 102 - 38N_f/3$ ,  $b_2^{MS} = 2857/2 - 5033N_f/18 + 325N_f^2/54$ . The parameter  $\Delta$  in eq. (2) is due to the lower boundary of the GML integral<sup>8,9/</sup>. By a particular choice of  $\Delta$  one fixes the definition of  $\Lambda$ :  $\Lambda = \Lambda(\Delta)^*$ . Eq. (2) is solved by iterations and the result is reexpanded in  $1/L$ :

$$\alpha_s(Q^2) = \frac{4\pi}{b_0 L} \left\{ 1 - \frac{L_1}{L} + \frac{1}{L^2} \left[ L_2 - \frac{b_1}{b_0^2} L_1 + \frac{b_2 b_0 - b_1^2}{b_0^4} \right] + O(1/L^3) \right\}, \quad (3)$$

where

$$L_1 = \frac{b_1}{b_0^2} \ln(b_0 L) - \Delta. \quad (4)$$

The expansion (3) is useful, of course, only if it converges rapidly enough. In fact, the convergence of the  $1/L$  series depends (i) on the value of  $L$  we are interested in and (ii) on the choice of  $\Delta$ .

We emphasize that the most important for perturbative QCD is the region  $L > 3$ , since  $L=3$  corresponds to  $\alpha_s \sim 0.5$ , and the reliability of perturbation theory for larger  $\alpha_s$  is questionable. Hence, in a realistic situation the naive expansion parameter  $1/L$  is smaller than (but usually close to) one third. Of course,  $1/3$  is not very small, so one must check the coefficients of the  $1/L$  expansion more carefully. First, there is a  $\Delta$ -convention-independent term  $(b_2 b_0 - b_1^2)/(b_0^4 L^2)$  which reduces for  $N_f = 3$  to roughly  $0.25/L^2$  and gives, therefore, less than 3%-correction to the simplest formula (1). There are also  $\Delta$ -convention-dependent terms like  $L_1/L$ ,  $L_1/L^2$  and one should choose  $\Delta$  so as to minimize the upper value of the ratio  $L_1/L$  in the  $L$ -region of interest.

\* Of course,  $\Lambda$  depends also on the renormalization scheme chosen.

If one takes, e.g.,  $\Delta = \Delta_{opt} = (b_1/b_0^2) \ln(4b_0)$ , then  $L_1 = (b_1/b_0^2) \ln(L/4)$  and the ratio  $L_1/L$  is smaller than 7% in the whole region  $L > 3$ . Another choice<sup>10/</sup> is to take  $\Delta = \Delta(Q_0^2) = (b_1/b_0^2) \ln(b_0 L_0)$ , where  $L_0 = \ln(Q_0^2/\Lambda^2)$  and  $Q_0^2$  lies somewhere in the middle of the  $Q^2$ -region analysed. In this case  $L_1 = (b_1/b_0^2) \ln(L/L_0)$ , i.e.,  $L_1/L$  is zero for  $Q^2 = Q_0^2$  and smaller than 7% for all  $Q^2$  in the region where  $L > 3$ . An important observation is that both the choices minimize the corrections not only in eq. (3) but also in the GML equation (2).

Really, for small  $G$  the only dangerous term in eq. (2) is  $\ln G$ , hence, the best thing to do is to compensate it by taking  $\Delta = -(b_1/b_0^2) \ln \bar{G}$ , where  $\bar{G}$  is  $\alpha_s(Q^2)/4\pi$  averaged (in some sense) over the relevant  $Q^2$ -region. After this has been done, one may safely solve eq. (2) by iterations and perform the  $1/L$ -expansion. For a proper choice of  $\Delta$  eq. (3) has 1% accuracy for  $L > 3$ , and, moreover, the total correction to the simplest formula (1) is less than 10%. However, accepting the most popular prescription  $\Delta_{pop} = (b_1/b_0^2) \ln b_0 = \Delta(Q^2 = e\Lambda^2)$  (the only motivation for  $\Delta_{pop}$  being the "aesthetic" criterion that  $L_1$  should have the shortest form  $L_1 = (b_1/b_0^2) \ln L$ ) one minimizes  $L_1/L$  in the region  $Q^2 \sim 3\Lambda^2$  nobody is really interested in. Moreover, in the important region  $L \sim 3$  one has  $L_1^{pop}/L \sim 1/3$  and the convergence of the  $1/L$ -series is very poor in this case.

Thus, the  $\Lambda$ -parametrization (eq. (3)) gives a rather compact and sufficiently precise expression for the effective coupling constant in the spacelike region provided a proper choice of the  $\Delta$ -parameter has been made.

## 3. $\Lambda$ -PARAMETRIZATION AND $R(e^+e^- \rightarrow \text{HADRONS}; s)$

The standard procedure (see, e.g.,<sup>11/</sup> and references therein) is to calculate the derivative  $D(Q^2) = Q^2 dt/dQ^2$  of the vacuum polarization  $t(Q^2)$  related to  $R$  by

$$R(s) = \frac{1}{2\pi i} (t(-s+i\epsilon) - t(-s-i\epsilon)). \quad (5)$$

In perturbative QCD  $D(Q^2)$  is given by the  $\alpha_s(Q^2)$ -expansion:

$$D(Q^2) = \sum_q e_q^2 \left\{ 1 + \frac{\alpha_s(Q^2)}{\pi} + d_2 \left( \frac{\alpha_s(Q^2)}{\pi} \right)^2 + d_3 \left( \frac{\alpha_s(Q^2)}{\pi} \right)^3 + \dots \right\}. \quad (6)$$

Only  $d_2$  is known now<sup>11,12/</sup>, its value depending on the renormalization scheme chosen. Using eq. (5) and the definition of  $D$ , one can relate  $R(s)$  (or, more precisely, its perturbative QCD version  $R^{QCD}(s)$ ) directly to  $D(Q^2)$

$$R^{\text{QCD}}(s) = \frac{1}{2\pi i} \int_{-s-i\epsilon}^{-s+i\epsilon} D(\sigma) \frac{d\sigma}{\sigma}. \quad (7)$$

Integration in eq. (7) goes below the real axis from  $-s-i\epsilon$  to zero and then above the real axis to  $-s+i\epsilon$ .

In a shorthand notation  $D \Rightarrow R \equiv \Phi[D]$ . In some important cases the integral (7) can be calculated explicitly: (8)

$$1 \Rightarrow 1,$$

$$\frac{1}{L_\sigma} \Rightarrow \frac{1}{\pi} \text{arctg}(\pi/L_s) = \frac{1}{L_s} \left\{ 1 - \frac{1}{3} \frac{\pi^2}{L_s^2} + \dots \right\}, \quad (9)$$

$$\frac{\ln(L_\sigma/L_0)}{L_\sigma^2} \Rightarrow \frac{\ln(\sqrt{L_s^2 + \pi^2}/L_0) - (L_s/\pi) \text{arctg}(\pi/L_s) + 1}{L_s^2 + \pi^2} = \quad (10)$$

$$= \frac{\ln(L_s/L_0)}{L_s^2} \left\{ 1 - \frac{\pi^2}{L_s^2} + \dots \right\} + \frac{5}{6} \frac{\pi^2}{L_s^4} + \dots \quad (11)$$

$$\frac{1}{L_\sigma^2} \Rightarrow \frac{1}{L_s^2 + \pi^2} = \frac{1}{L_s^2} \left\{ 1 - \frac{\pi^2}{L_s^2} + \dots \right\},$$

$$\frac{1}{L_\sigma^n} \Rightarrow (-1)^n \frac{1}{(n-1)!} \left( \frac{d}{dL_s} \right)^{n-2} \frac{1}{L_s^2 + \pi^2} = \frac{1}{L_s^n} \left\{ 1 - \frac{\pi^2}{L_s^2} \frac{n(n+1)}{6} + \dots \right\}, \quad (12)$$

where  $L_s = \ln(s/\Lambda^2)$ ,  $L_\sigma = \ln(\sigma/\Lambda^2)$  and  $L_0$  is the constant depending on the  $\Lambda$ -choice.

Using the  $\Lambda$ -parametrization for  $a_s(\sigma)$  and incorporating eqs. (8)-(12) (as well as their generalizations for  $\ln^2 L/L^3$ ,  $\ln L/L^3$  etc.) produces the expansion for  $R^{\text{QCD}}(s)$

$$R^{\text{QCD}}(s) = \left( \sum_q e_q^2 \right) \left\{ 1 + \sum_{k=1} d_k \Phi[(a_s/\pi)^k] \right\} \quad (13)$$

in which all the  $(\pi^2/L^2)^N$ -terms are summed up explicitly.

#### 4. QUEST FOR THE BEST EXPANSION PARAMETER

Note that the expansion (13) is not an expansion in powers of some particular parameter since the application of the  $\Phi$ -operation normally violates nonlinear relations:  $\Phi[1/L^2] \neq (\Phi[1/L])^2$ , etc. A priori, there are no grounds to believe that a power expansion is better than any other (say, Fourier). In fact, the expansion (13) converges better than the genera-

ting expansion (6) for  $D(\sigma)$  because, as it follows from eqs. (9)-(12),  $\Phi[a_s^N]$  is always smaller than  $a_s^N$ . Moreover,

$(\Phi[a_s^{N+1}])^{1/N+1} < (\Phi[a_s^N])^{1/N}$  i.e., the effective expansion parameter decreases in higher orders. Thus, if one succeeded in obtaining a good  $a_s^N$  expansion for  $D(\sigma)$  (with all  $d_N$  being small numbers), then the resulting  $\Phi[a_s^N]$ -expansion for  $R^{\text{QCD}}(s)$  is even better, and the best thing to do is to leave it as it is.

However, if one insists that the result for  $R^{\text{QCD}}(s)$  should have a form of a power expansion, then the best expansion parameter is evidently  $\Phi[a_s/\pi]$  because the largest nontrivial (i.e.,  $O(a_s/\pi)$ ) term of the expansion is reproduced in the exact form and only higher terms are spoiled. The analogue of the simplest  $\Lambda$ -parametrization for  $a_s(Q^2)$  (eq. (1)) is then

$$\tilde{a}_s(Q^2) = \frac{4}{b_0} \text{arctg}\left(\frac{\pi}{\ln(Q^2/\Lambda^2)}\right). \quad (14)$$

Using eqs. (8)-(13) it is easy to realize that  $a_s(Q^2)$  is really a bad expansion parameter, because if one reexpands  $\tilde{a}_s(Q^2)$  in  $a_s(Q^2)$ , then there appear terms with large coefficients

$$\tilde{a}_s(Q^2) = a_s(Q^2) \left\{ 1 - \frac{1}{3} \left( \frac{\pi b_0}{4} \right)^2 \left( \frac{a_s(Q^2)}{\pi} \right)^2 + \dots \right\} \approx a_s \left\{ 1 - 17 \left( \frac{a_s}{\pi} \right)^2 + \dots \right\}. \quad (15)$$

If one reexpands  $\tilde{a}_s(Q^2)$  in  $\text{Re} a_s(-Q^2)$  then the corresponding coefficient is even 2 times larger, whereas if  $\tilde{a}_s(Q^2)$  is reexpanded in  $|a_s(-Q^2)|$ , the coefficient is 2 times smaller. This observation is in full agreement with the result of ref.<sup>/5/</sup> quoted in the introduction.

#### 5. CONCLUDING REMARKS

It should be noted that the change of the expansion parameter as given by eq. (15) affects only the  $(a_s/\pi)^3$  coefficient of the  $R^{\text{QCD}}$ -expansion which has not been calculated yet. So, within the present-day accuracy, all expansions for  $R^{\text{QCD}}$  have the same coefficients. It is worth emphasizing, nevertheless, that the  $\pi^2/L^2$  terms produce for  $a_s \geq 0.3$  more than 20%-correction to  $a_s$ , i.e., they are more important (for an optimal choice of the  $\Lambda$ -parameter) than the 2-loop corrections in eq. (3)).

To conclude, we have described the construction of an optimized (i.e., rapidly convergent)  $\Lambda$ -parametrization for the effective QCD coupling constant in the spacelike region, and then we used it to obtain the fastest convergent expansion for the time-like quantity  $R^{\text{QCD}}(s)$ . The technique outlined in the present paper can be applied also to other  $R^{\text{QCD}}$ -like

quantities. Such quantities do appear, e.g., in the QCD sum rule approach<sup>/13/</sup> in which the analysis of hadronic properties is based on the study of vacuum correlators of various currents. They appear also in an alternative approach<sup>/14/</sup> based on the finite-energy sum rules<sup>/15/</sup>. It should be stressed that in the latter approach the  $R^{QCD}$ -like quantities enter into the basic integral relation, and the analysis is most conveniently performed if one has a simple analytic expression similar to that described above.

#### ACKNOWLEDGEMENTS

I am grateful to A.V.Efremov and D.V.Shirkov for their interest in this work and support. I thank D.I.Kazakov, O.V.Tarasov and A.A.Vladimirov for useful discussions.

#### REFERENCES

1. Politzer H.D. Physics Reports, 1974, 14C, p. 129.
2. Mueller A.H. Physics Reports, 1981, 73, p. 237. Reya E. Physics Reports, 1981, 69, p. 195.
3. Efremov A.V., Radyushkin A.V. Rivista del Nuovo Cimento, 1980, 3, No. 2.
4. Bace M. Physics Letters B, 1978, 78, p. 132.
5. Pennigton M.R., Ross G.G. Physics Letters B, 1981, 102, p. 167.
6. Caswell W. Phys.Rev.Letters, 1974, 33, p. 244. Jones D.R.T. Nucl.Physics B, 1974, 75, p. 531. Egoryan E.Sh., Tarasov O.V. Teor.Mat.Fizika, 1979, 41, p. 26.
7. Tarasov O.V., Vladimirov A.A., Zharkov A.Yu. Phys.Letters B, 1980, 93, p. 429.
8. Bardeen W.A. et al. Phys.Rev. D, 1978, 18, p. 3998.
9. Vladimirov A.A. Yad.Fizika, 1980, 31, p. 1083.
10. Abbott L.F. Phys.Rev.Letters, 1980, 44, p. 1569.
11. Chetyrkin K.G., Kataev A.L., Tkachov F.V. Nucl.Phys.B, 1980, 174, p. 345.
12. Dine M., Sapirstein J. J.Phys.Rev.Lett., 1979, 43, p. 668. Celmaster W., Gonsalvez R. Phys.Rev.D, 1980, 21, p.3112.
13. Shifman M.A., Vainshtein A.I., Zakharov V.I. Nucl.Phys.B, 1979, 147, pp. 385, 448.
14. Krasnikov N.V., Pivovarov A.A. Preprint NBI-HE-81-38, Niels Bohr Institute, Copenhagen (October 1981).
15. Chetyrkin K.G., Krasnikov N.V., Tavkhelidze A.N.. Physics Letters, B, 1978, 76, p. 83.

Received by Publishing Department  
on February 26 1982.

#### WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices - in US \$,  
including the packing and registered postage

D9-10500	Proceedings of the Second Symposium on Collective Methods of Acceleration. Dubna, 1976.	11.00
D2-10533	Proceedings of the X International School on High Energy Physics for Young Scientists. Baku, 1976.	11.00
D13-11182	Proceedings of the IX International Symposium on Nuclear Electronics. Varna, 1977.	10.00
D17-11490	Proceedings of the International Symposium on Selected Problems of Statistical Mechanics. Dubna, 1977.	18.00
D6-11574	Proceedings of the XV Symposium on Nuclear Spectroscopy and Nuclear Theory. Dubna, 1978.	4.70
D3-11787	Proceedings of the III International School on Neutron Physics. Alushta, 1978.	12.00
D13-11807	Proceedings of the III International Meeting on Proportional and Drift Chambers. Dubna, 1978.	14.00
	Proceedings of the VI All-Union Conference on Charged Particle Accelerators. Dubna, 1978. 2 volumes.	25.00
D1,2-12450	Proceedings of the XII International School on High Energy Physics for Young Scientists. Bulgaria, Primorsko, 1978.	18.00
D-12965	The Proceedings of the International School on the Problems of Charged Particle Accelerators for Young Scientists. Minsk, 1979.	8.00
D11-80-13	The Proceedings of the International Conference on Systems and Techniques of Analytical Computing and Their Applications in Theoretical Physics. Dubna, 1979.	8.00
D4-80-271	The Proceedings of the International Symposium on Few Particle Problems in Nuclear Physics. Dubna, 1979.	8.50
D4-80-385	The Proceedings of the International School on Nuclear Structure. Alushta, 1980.	10.00
	Proceedings of the VII All-Union Conference on Charged Particle Accelerators. Dubna, 1980. 2 volumes.	25.00
D4-80-572	N.N.Kolesnikov et al. "The Energies and Half-Lives for the $\alpha$ - and $\beta$ -Decays of Transfermium Elements"	10.00
D2-81-543	Proceedings of the VI International Conference on the Problems of Quantum Field Theory. Alushta, 1981	9.50
D10,11-81-622	Proceedings of the International Meeting on Problems of Mathematical Simulation in Nuclear Physics Researches. Dubna, 1980	9.00

Orders for the above-mentioned books can be sent at the address:  
Publishing Department, JINR  
Head Post Office, P.O.Box 79 101000 Moscow, USSR

## SUBJECT CATEGORIES OF THE JINR PUBLICATIONS

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches
19.	Biophysics

Радюшкин А.В. Оптимальная лямбда-параметризация эффективной константы связи в КХД для пространственно-и времениподобной областей E2-82-159

Сформулирован алгоритм, позволяющий в явном виде просуммировать  $(\pi^2/\ln^2(Q^2/\Lambda^2))^N$  -поправки к  $a_s(Q^2)$ , обусловленные аналитическим продолжением из пространственно-подобной во времениподобную область передач импульса. Показано, что во времениподобной области наилучшим параметром разложения является  $(4/b_0) \operatorname{arctg}(\pi/\ln(q^2/\Lambda^2))$ .

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1982

Radyushkin A.V. Optimized Lambda-Parametrization for the QCD Running Coupling Constant in Spacelike and Timelike Regions E2-82-159

The algorithm is described that enables one to perform an explicit summation of all the  $(\pi^2/\ln^2(Q^2/\Lambda^2))^N$  -corrections to  $a_s(Q^2)$  that appear owing to the analytic continuation from spacelike to timelike region of momentum transfer.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1982