


84-193

C 323.5



СООБЩЕНИЯ
ОБЪЕДИНЕННОГО
ИНСТИТУТА
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

2857/84

2530

E2-84-193

St. Mr̄wezyński

ON MULTIPLICITY DISTRIBUTIONS
AND A PRESSURE ENSEMBLE

1984

High multiplicities of particles produced in high energy collisions seem to be a good reason for applying statistical methods to study production processes. It is well-known^{/1/} that fluctuations of the number, N , of particles in a grand canonical ensemble are (outside a phase-transition region) proportional to the root square of an average number of particles, $\langle N \rangle$. This leads to a disagreement with multiplicity data on hadron-hadron collisions which provides a linear dependence of dispersion,

$$D = \sqrt{\langle N^2 \rangle - \langle N \rangle^2}, \text{ on average value - the Wróblewski formula}^{/2/}.$$


The volume of all systems, in particular those with different number of particles, belonging to the grand canonical ensemble is the same. However, the hot hadron gas which occurs in high energy collisions is not closed in any container with a fixed volume*. Thus, in our opinion, a more appropriate basis for studying the problem is the so-called pressure ensemble^{/3/}. In this ensemble one considers systems under fixed external pressure. Up to our knowledge, Gorenstein was the first who applied the pressure ensemble to hadron physics^{/4/}. In our considerations we use some results of Gorenstein's paper^{/4/}, where the distribution of the number of quarks in the MIT bag and possible implications for multiplicity distributions were discussed.

The grand canonical pressure partition function is defined as

$$\Pi(p, T, \mu) = \int_0^{\infty} dV \exp\left(-\frac{pV}{T}\right) \Xi(V, T, \mu),$$

where p is the pressure, T is the temperature, V is the volume and μ is the chemical potential. Ξ is the grand canonical partition function.

* It should be underlined that even in the grand canonical ensemble the volume of a system is not fixed during the time evolution of the system if summation over the ensemble is performed at a definite moment of time, for example the moment of decaying of the system. Thus, the volume is fixed in such a sense that the volumes of all systems belonging to the ensemble are equal at this definite moment of time.



$$\bar{N}(V, T, \mu) = \sum_{N=0}^{\infty} Z^N Q_N(V, T),$$

where $Z \equiv \exp \beta \mu$, $\beta = 1/T$, and Q_N is the canonical partition function of N particles.

The probability of finding N particles, \mathcal{P}_N , in the grand canonical pressure ensemble is expressed as

$$\mathcal{P}_N = \frac{Z^N}{\Pi} \int_0^{\infty} dV \exp\left(-\frac{pV}{T}\right) Q_N(V, T). \quad (1)$$

Let us consider the model of hadron gas extensively discussed in the literature^{/5/}, where attractive forces are represented by the mass spectrum of particles, $\rho(m)$, and repulsive forces by a Van der Waals correction to the volume. The canonical partition function of such a gas looks like

$$Q_N(V, T) = \frac{1}{N!} \int \prod_{i=1}^N [d^3 p_i dm_i \rho(m_i) (V - \sum_{i=1}^N v_i)] \cdot \exp\left(-\sum_{i=1}^N \beta \sqrt{p_i^2 + m_i^2}\right) \theta\left(V - \sum_{i=1}^N v_i\right). \quad (2)$$

The integration is performed over the momenta and masses of particles. v_i is the volume of an i -th hadron. There are so many different types of particles that, except for high densities at low temperature, quantum effects are negligible and Boltzmann statistics is sufficient.

Substituting (2) in (1), one finds the geometrical distribution

$$\mathcal{P}_N = (1-q) q^N, \quad (3)$$

where

$$q = \frac{z'T}{p} \int d^3 p dm \exp\left(-\beta \sqrt{p^2 + m^2} - \beta p v\right) \rho(m).$$

It is seen that the form of (3) is independent of details of the hadron gas model. In particular, the distribution (3) is valid for ideal gas.

The geometrical distribution fulfils approximately KNO scaling^{/6/} for $N \gg 1$. Thus, the Wróblewski formula is also approximately satisfied. Although, the shape of the geometrical distribution is far from the experimental one.

As pointed out^{/7/} many years ago, experimental data cannot be described within the model where the existence of one fireball is assumed. Only a few-fireball model can be realistic^{/8/}. In the case of e^+e^- annihilation into hadrons, jets can be interpreted as a result of the existence of two or more fireballs^{/9/}.

Let us consider a model of k -fireballs. For this purpose we transform the discrete distribution (3) into the continuous one (4)

$$\mathcal{P}(n) = a \exp(-an), \quad (4)$$

where n is a continuous variable. When $\langle N \rangle \gg 1$,

$$\mathcal{P}_N \approx \mathcal{P}(N), \quad q = \exp(-a) \approx 1 - a.$$

Because the statistical methods are applicable to $\langle N \rangle \gg 1$, it is sufficient for our discussion to use the continuous distribution (4).

We restrict ourselves to the simplest situation where the thermodynamic characteristics of all fireballs are the same. Assuming that the fireballs independently contribute to the resulting multiplicity, the distribution of particles emitted by these fireballs is the convolution of k distributions (4):

$$\begin{aligned} \mathcal{P}^k(n) &= \int_0^{\infty} \prod_{i=1}^k [dn_i \mathcal{P}(n_i)] \delta\left(n - \sum_{i=1}^k n_i\right) = \\ &= \frac{a^k}{(k-1)!} \exp(-an) n^{k-1}. \end{aligned} \quad (5)$$

The average value of n is expressed by the formula

$$\langle n \rangle = \frac{k}{a} \quad (6)$$

and the second moment of (5) is the following

$$D = \sqrt{\langle n^2 \rangle - \langle n \rangle^2} = \frac{\sqrt{k}}{a} = \frac{\langle n \rangle}{\sqrt{k}}. \quad (7)$$

The ratio $D/\langle n \rangle$ is simply related to a number of the sources of particles.

Multiplying $\mathcal{P}^k(n)$ by $\langle n \rangle$, we get

$$\langle n \rangle \mathcal{P}^k(n) = \frac{k^k}{(k-1)!} \left[\frac{n}{\langle n \rangle}\right]^{k-1} \exp\left(-k \frac{n}{\langle n \rangle}\right).$$




Fig.1. The multiplicity distribution of charged particles produced in antiproton-proton collisions at $\sqrt{s} = 540$ GeV. The data are taken from Ref.^{/10/}. The solid line is found from formula (5) with $k = 4$.

Thus, the distribution (5) exactly fulfils KNO scaling. An analogous discrete distribution satisfies KNO approximately*.

In the case of hadron-hadron or lepton-lepton collisions we expect that an increase of multiplicity with incident energy is due to decreasing the thermodynamical factor "a" present in (6). The assumption of constancy of the number of fireballs leads to KNO scaling and, in particular, to the linear function $D(\langle n \rangle)$. In order to compare the distribution (5) with experimental data on hadron-hadron collisions, we assume that the number of fireballs is the same in each event. This assumption is quite realistic in the case of e^+e^- or $\bar{p}p$ annihilation. Because in hadron-hadron interactions there are collisions with different inelasticity coefficient, the above assumption is only approximately valid. Comparing (7) with the experimental value of the $D/\langle n \rangle$ ratio which is about $0.5^{/2/}$, we find that $k \approx 4$.

In Fig.1 we compare the experimental multiplicity distribution in $\bar{p}p$ interactions at collider energies^{/10/} with the predictions of (5) for $k = 4$. The agreement seems to be quite satisfactory.

In the case of collisions with nuclei the number of fireballs can strongly depend on impact parameter. Thus, the multiplicity distribution in inelastic nucleus-nucleus collisions is the sum of distributions (5) with different k . Such summation gives the distribution which is "wider" than that in $\bar{p}p$ interactions. If we consider central nucleus-nucleus collisions where an impact parameter is restricted to small values, we expect that the number k does not change significantly from event to event. Because the number of fireballs in central collisions with nuclei

*The distribution (5) has been also found by other authors, whose argumentation, however, has been different than ours. See: N. Suzuki, Prog. Theor. Phys., 1974, 51, p. 1629. A. Giovannini, Nuovo Cim., 1976, 34A, p. 647. Ref.^{/4/}; P. Carruthers, C.C. Shih, Phys. Lett., 1983, 127B, p. 242.




Fig.2. The dispersion versus the average number of negative pions produced in nucleus-nucleus collisions at a momentum 4.5 GeV/c per nucleon. Crosses - inelastic collisions, circles - central collisions. The full and open circles correspond to different central triggers, for details see Ref.^{/11/}. The straight line corresponding to multiplicity distributions for proton-proton inelastic collisions is found according to Wróblewski formula^{/2/}.

right line corresponding to multiplicity distributions for proton-proton inelastic collisions is found according to Wróblewski formula^{/2/}.

is greater than in $\bar{p}p$ interactions, the multiplicity distribution should be "narrower" than in $\bar{p}p$. In Fig.2 taken from ref.^{/11/} it is shown the dispersion versus an average value for π^- produced in nucleus-nucleus, inelastic and central, collisions for fixed incident energy projectile and different targets. Both features of the multiplicity distributions discussed above are seen. If we assume that at fixed projectile energy the thermodynamical factor "a" is independent of target mass and multiplicity increases with increasing the number of fireballs, we find, comparing (6) and (7), $D = \sqrt{a \langle n \rangle}$. As shown in Fig.2, such a Poisson type relation has been found in the experiment^{/11/}.

If it were possible to choose such a phase-space region of secondaries where only one source contributes, the geometrical distribution (3) would be found in this region. A good candidate for such a region is the backward hemisphere in LAB for nucleus-nucleus collisions. Because the emission of backward particles is kinematically unfavourable, we expect that only fireball with the smallest value of rapidity significantly contributes to this region. Consequently, the multiplicity distribution of backward particles should be geometrical. It occurs that such a distribution has been experimentally found^{/12-14/}, see Fig.3.

We summarize our considerations as follows. In the pressure ensemble one finds the geometrical distribution of the number of particles emitted from one source. Assuming the existence of a few fireballs in high energy collisions, we are led to the multiplicity distribution being the convolution of a few geometrical ones. The resulting distribution satisfies KNO scaling and fulfils other features of experimental multiplicity distributions in hadron-hadron and nucleus-nucleus interactions.

I am grateful to M. Gaździcki, A.I. Golokhvastov, S.A. Khorozov and M. Šumbera for fruitful discussions.




Fig.3. The multiplicity distribution of protons with momentum greater than 240 MeV/c emitted in the backward (in LAB) hemisphere for $^{12}\text{C}-^{181}\text{Ta}$ collisions at a momentum of 4.2 GeV/c per nucleon. The data are taken from Ref. /12/. The solid line corresponds to the geometrical distribution (3).

REFERENCES

1. See, e.g., K.Huang. "Statistical Mechanics", John Wiley and Sons, Inc., New York, 1963.
2. Wroblewski A. Acta Phys.Pol., 1973, B4, p. 857.
3. Ter Haar D., Wergeland H. Phys.Rep., 1971, C1, p. 31. Münster A. "Statistical Physics", Springer, Berlin, Heidelberg, New York, 1969.
4. Gorenstein M.I. Yad.Fiz., 1980, 31, p. 1630.
5. Baacke J. Acta Phys.Pol., 1977, B8, p. 625. Hagedorn R. J.Refelski. Phys.Lett., 1980, B97, p. 136. Hagedorn R. Z.f.Phys., 1983, C17, p. 265. Gorenstein M.I., Petrov V.K., Zinoviev G.M. Phys.Lett., 1981, B106, p. 327. Gorenstein M.I. et al. Teor.Mat.Fiz., 1982, 52, p. 346. Kapusta J.I. Phys.Rev., 1981, D23, p. 2453.
6. Koba Z., Nielsen H.B., Olesen P. Nucl.Phys., 1972, B40, p. 317.
7. Ciok P. et al. Nuovo Cim., 1958, 8, p. 166.
8. For review of thermodynamical fireball models see E.L.Feinberg, Uspekhi Fiz. Nauk, 1983, 139, p. 1.
9. Hoang T.F. Z.f.Phys., 1983, C18, p. 19.
10. Alpgard K. et al. Phys.Lett., 1983, B121, p. 209.
11. Anikina M.Kh. et al. Z. f. Phys., 1981, C9, p. 105.
12. Angelov N. et al. Yad.Fiz., 1980, 32, p. 1582.
13. Angelov N. et al. Yad.Fiz., 1981, 33, p. 190.
14. Vokal S. et al. JINR, 1-83-652, Dubna, 1983. (submitted to Z. f. Phys. A).

Received by Publishing Department
on March 27, 1984.

WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices - in US \$,
including the packing and registered postage

D4-80-385	The Proceedings of the International School on Nuclear Structure. Alushta, 1980.	10.00
	Proceedings of the VII All-Union Conference on Charged Particle Accelerators. Dubna, 1980. 2 volumes.	25.00
D4-80-572	N.N.Kolesnikov et al. "The Energies and Half-Lives for the α - and β -Decays of Transfermium Elements"	10.00
D2-81-543	Proceedings of the VI International Conference on the Problems of Quantum Field Theory. Alushta, 1981	9.50
D10,11-81-622	Proceedings of the International Meeting on Problems of Mathematical Simulation in Nuclear Physics Researches. Dubna, 1980	9.00
D1,2-81-728	Proceedings of the VI International Seminar on High Energy Physics Problems. Dubna, 1981.	9.50
D17-81-758	Proceedings of the II International Symposium on Selected Problems in Statistical Mechanics. Dubna, 1981.	15.50
D1,2-82-27	Proceedings of the International Symposium on Polarization Phenomena in High Energy Physics. Dubna, 1981.	9.00
D2-82-568	Proceedings of the Meeting on Investigations in the Field of Relativistic Nuclear Physics. Dubna, 1982	7.50
D9-82-664	Proceedings of the Symposium on the Problems of Collective Methods of Acceleration. Dubna, 1982	9.20
D3,4-82-704	Proceedings of the IV International School on Neutron Physics. Dubna, 1982	12.00
D2,4-83-179	Proceedings of the XV International School on High-Energy Physics for Young Scientists. Dubna, 1982	10.00
	Proceedings of the VIII All-Union Conference on Charged Particle Accelerators. Protvino, 1982. 2 volumes.	25.00
D11-83-511	Proceedings of the Conference on Systems and Techniques of Analytical Computing and Their Applications in Theoretical Physics. Dubna, 1982.	9.50
D7-83-644	Proceedings of the International School-Seminar on Heavy Ion Physics. Alushta, 1983.	11.30
D2,13-83-689	Proceedings of the Workshop on Radiation Problems and Gravitational Wave Detection. Dubna, 1983.	6.00

Orders for the above-mentioned books can be sent at the address:
Publishing Department, JINR
Head Post Office, P.O.Box 79 101000 Moscow, USSR

SUBJECT CATEGORIES OF THE JINR PUBLICATIONS

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches
19.	Biophysics

Мрувчински С.

E2-84-193

О распределениях по множественности и ансамбле с давлением

Пользуясь так называемым ансамблем с давлением, получаем в рамках многофайрбольной модели распределение по множественности частиц, образованных во взаимодействиях при высоких энергиях. Показано, что полученное распределение удовлетворяет KNO скейлингу и может описать довольно широкий класс экспериментальных данных по множественности, от $p\bar{p}$ взаимодействия при энергии коллайдера до ядро-ядро соударения при нескольких ГэВ на нуклон.

Работа выполнена в Лаборатории высоких энергий ОИЯИ.

Сообщение Объединенного института ядерных исследований. Дубна 1984

Mrówczyński St.

E2-84-193

On Multiplicity Distributions and a Pressure Ensemble

Using a pressure ensemble, the multiplicity distribution in the multi-fireball model is found. The distribution satisfies KNO scaling and describes a wide class of multiplicity data from $p\bar{p}$ interactions at collider energy to nucleus-nucleus collisions at a few GeV per nucleon.

The investigation has been performed at the Laboratory of High Energies, JINR.

Communication of the Joint Institute for Nuclear Research. Dubna 1984