

**СООБЩЕНИЯ
ОБЪЕДИНЕННОГО
ИНСТИТУТА
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА**

E2-86-218

K.G.Denisenko*, St.Mrowczyński

DELTA'S IN HADRON GAS

* Radium Institute, Leningrad, USSR

I. INTRODUCTION

The classical (nonquantum) kinetic theory approach to the systems including unstable particles has been developed in our recent paper ^{1/}. The hadron gas is an example of the system where the presence of unstable particles, hadron resonances, essentially influences the properties of the gas. The decay width of hadron resonances is often greater than the temperature of the hadron systems, what suggests the importance of resonance instability. On the other hand, the lifetime of the resonances can be longer than the average time interval between successive collisions in the gas. So, the resonances should be treated in a similar manner as stable particles are. The starting point of our approach has been the introduction of the profile function which is the generalization of the delta function that "keeps" a stable particle on the mass-shell. Then the phase space element of the resonance with four-momentum p is chosen in the form $\Delta(p^2)d^4p$, where $\Delta(p^2)$ describes the mass smearing of the resonance. Using the profile function we have defined the resonance distribution function and macroscopical quantities. Kinetic equations have been formulated, where, besides binary collisions, resonance formation processes and resonance decays have been taken into account. It has been shown that the profile function can be uniquely determined through experimentally measurable quantities if the transition rates of the processes with the resonance involved, satisfy the detailed balance condition or the bilateral normalization condition. For the resonance formation cross section in the Breit-Wigner form the profile function looks like

$$\Delta(M^2) = \frac{\Gamma}{2\pi M[(M - \bar{M})^2 + \Gamma^2/4]}, \quad (1)$$

where \bar{M} and Γ are the average resonance mass and the resonance decay width. Finally it has been demonstrated that the equilibrium characteristics of the resonances can be expressed in the form

$$\mathcal{Q}_R = \int dM M \Delta(M^2) \mathcal{Q}_{st}(M), \quad (2)$$

where $\mathcal{Q}_{st}(M)$ is the respective characteristic for stable particles with mass M .

The aim of this paper is twofold. In Sec. II we show that our model is, in the case of equilibrium systems, equivalent under certain assumptions to the S-matrix formulation of statistical mechanics by Dashen, Ma and Bernstein^{/2/}. In their approach unstable particles (resonances) occur through the S-matrix elements of the resonance scattering of stable particles.

In Sec. III the equilibrium characteristics of classical gas of nucleons, deltas and pions are studied. We show that if one treats the delta isobars as stable particles their number can be significantly underestimated.

II. S-MATRIX EXPANSION OF THE GRAND POTENTIAL

To relate our kinetic theory approach to the Gibbs statistical mechanics, let us write down the grand canonical potential of the system of stable and unstable particles. Only the interaction, which leads to the resonance formation and resonance decay, is included, i.e., the resonance scattering of stable particles is assumed to dominate two-body interaction. So the grand potential reads

$$\Omega = \sum_i \Omega^i(m_i, \mu_i) + \sum_j \int dM M \Delta^i(M^2) \Omega^i(M, \mu_j), \quad (3)$$

where $\Omega(m, \mu)$ is the grand canonical potential of the one-component ideal gas of particles with mass m and chemical potential μ . In the first term summation is performed over the sorts of stable particles while in the second one over the unstable particle sorts. Formula (3), of course, follows from (2).

The S-matrix expansion of the grand canonical potential for the system of relativistic particles is the following


$$\Omega = \Omega_0 - TV \sum_{\nu} a_{\nu} e^{\beta \mu \mathcal{N}}, \quad (4)$$

where Ω_0 is the grand potential of noninteracting particles, V is the volume of the system, $T = \beta^{-1}$ is the temperature, \mathcal{N} is the set of conserved charges and μ is the set of chemical potentials, $\nu = (\mathcal{N}, a)$ where a describes all quantum numbers required in fixing the system with \mathcal{N} charges:

$$a_{\nu} = \int \frac{d^3 \bar{p}}{(2\pi)^3} dM e^{-\beta \sqrt{\bar{p}^2 + M^2}} \frac{1}{4\pi i} \text{Tr}_{\nu} (S^{-1} \frac{\overleftrightarrow{\partial}}{\partial M} S)_c$$

where S is the S-operator and the subscript c indicates that only the so-called connected diagrams are taken into account. Let us now discuss, as in paper^{/2/}, the hadron gas consisting of nucleons and pions. Suppose the scattering processes are dominated by πN resonance scattering which lead to the delta

formation. We take into account only the collisions



and ignore all other interaction processes. For such a system the first term in Eq. (4) is the grand canonical potential of the ideal gas of nucleons and pions, and the second term relates to the πN resonant interaction. For simplicity we neglect the role of particle spins, which are taken into account through the particle degeneration factors only. Taking the matrix element of the process (5) in the Breit-Wigner form one finds^{/2/*}

$$\frac{1}{4\pi i} \text{Tr} (S^{-1} \frac{\overleftrightarrow{\partial}}{\partial M} S)_c = - \text{Re} \frac{\partial}{\partial M} \frac{\Gamma}{2\pi} (M - \bar{M} + i\Gamma/2)^{-1} + \text{Im} \frac{\Gamma^2}{2\pi} (M - \bar{M} - i\Gamma/2)^{-1} \frac{\partial}{\partial M} (M - \bar{M} + i\Gamma/2)^{-1}. \quad (6)$$

Substituting Eq. (6) in the formula (4) we get the following grand canonical potential valid in the lowest order S-matrix expansion (4)

$$\Omega = \Omega_0 + g_{\Delta} z \int dM M \Delta(M^2) \cdot \{-TV \int \frac{d^3 \bar{p}}{(2\pi)^3} e^{-\beta \sqrt{\bar{p}^2 + M^2}} \}. \quad (7)$$

where g_{Δ} is the delta degeneration factor and $z = e^{\beta \mu}$ is the delta fugacity. We have assumed that the gas is symmetric, i.e., the total electric charge of the system equals (in natural units) half of the total baryon charge. In such a case the chemical potentials of all sorts of baryons, in particular, of the deltas with different electric charge, are equal to each other and the chemical potentials of pions vanish. The function $\Delta(M^2)$ from Eq. (7) is exactly equal to that from formula (1). In the parentheses one recognizes the grand canonical potential of the ideal gas of spinless particles with mass M . So, the form of the potential (7) coincides with this one of (3). In that way we have shown that the idea of the profile function follows from the S-matrix formulation of statistical mechanics.

* There is a minor misprint in Eq. (7.11) of Ref. 2. The second term of the r.h.s. of this equation should be additionally divided by π . It should be also remembered that in Ref. 2 Γ denotes the decay half-width while in this paper Γ is the decay width.

III. GAS CHARACTERISTICS

Keeping in mind the formula (2) we discuss in this section the properties of classical gas consisting of nucleons, deltas and pions.

The Breit-Wigner formula (1) is correct for sufficiently narrow resonances what is not the case for the isobars Δ . So, to describe the mass distribution of deltas we modify formula (1) as follows

$$\Delta(M^2) = \xi \frac{\Gamma}{M[(M - \bar{M})^2 + \Gamma^2/4]} \theta(M - m_N - m_\pi),$$

where θ is the step function and m_N , m_π are the masses of nucleons and pions respectively, $\bar{M} = 1232$ MeV, $\Gamma = 115$ MeV. The coefficient ξ is found from the normalization condition $\int dM \Delta(M^2) = 1$.

The baryon density of the system reads

$$\rho = z g_\Delta \int dM \Delta(M^2) n(T, M) + z g_N n(T, m_N), \quad (8)$$

with

$$n(T, m) = \frac{1}{2\pi^2} T m K_2(\beta m)$$

and $g_N = 4$ and $g_\Delta = 16$; K_ν is the McDonald function. As quoted previously, the fugacities of N and Δ are equal to each other because the gas is assumed symmetric. The first term in Eq. (8) comes from the deltas; while the second one, from nucleons. Solving Eq. (8) with respect to z one finds the densities of deltas, ρ_Δ , and nucleons. In Fig. 1 we present the ratio of ρ_Δ to ρ as a function of temperature. In the case of symmetric gas this ratio is density independent. The dashed line is found under assumption that the deltas are stable particles, i.e., $\Gamma = 0$. It is seen that particularly at low temperature one strongly underestimates the density of the isobars if these particles are treated as stable ones. Our classical calculations cannot be extrapolated to the temperature $T < 30$ MeV since in such a case the quantum effects are essential.

The energy density of the system is

$$\epsilon = z g_\Delta \int dM \Delta(M^2) u(T, M) + z g_N u(T, m_N) + g_\pi u(T, m_\pi), \quad (9)$$

where $g_\pi = 3$ and

$$u(T, m) = \frac{1}{2\pi^2} T^2 m^2 [\beta m K_1(\beta m) + 3 K_2(\beta m)].$$




Fig. 1. The ratio of the delta density to the baryon density of the system versus temperature.

The first, second and the third term of Eq. (9) comes from the isobars, nucleons and pions, respectively. The numerical calculations show that the energy density is completely insensitive to the value of Γ of the resonance. The reason is the following. At low temperature, where the number of deltas strongly depends on the decay width, the absolute number of deltas is small as compared to the number of nucleons, see Fig. 1. Consequently the delta contribution to the energy density of the system is small. At high temperature, where the number of deltas is comparable to that of nucleons, the description of resonances is, in practice, the same for $\Gamma = 0$ and $\Gamma = 115$ MeV, see Fig. 1.

In the framework of the thermodynamical model of the pion production in relativistic nucleus-nucleus collisions the number of secondary pions (of all sorts) equals the number of pions plus the number of deltas present in the fireball at the moment of time of the fireball decay. The deltas are added since these isobars decay into pions which are finally registered. To discuss how the number of secondary pions changes, if one takes into account the finiteness of the isobar decay width, we have calculated the ratio

$$R = \frac{\rho_\pi + \rho_\Delta(\Gamma = 115 \text{ MeV})}{\rho_\pi + \rho_\Delta(\Gamma = 0)},$$

where ρ_π is the pion density. In Fig. 2 we present this ratio as a function of temperature for three values of the baryon density measured in the units of normal nuclear density $\rho_0 = 0.17 \text{ fm}^{-3}$. It is seen that at low temperatures R significantly exceeds unity. As is known the thermodynamical model overestimates, at least by a factor of 2, the multiplicity of secondary pions in nucleus-nucleus collisions^{3,4}. The proper description of deltas makes this difficulty even more serious.

It is known that in the nucleon-nucleon collisions at few GeV/c incident momentum, about 80% of produced pions come from




Fig.2. The ratio R as a function of temperature for three values of the baryon density.




Fig.3. The ratio of the delta density to the density of pions versus temperature for three values of the baryon density.

the delta decays ^{1/5}. So, it is of physical interest to consider the ratio of the delta density to the density of pions in the hadron gas. In Fig.3 we present this ratio versus temperature for three values of the baryon density. It is seen that the form of the ratio as a function of temperature essentially changes if one takes into account finiteness of the delta decay width. The ratio monotonically decreases when temperature increases what is not the case for $\Gamma = 0$.

IV. CONCLUSIONS

Establishing the relation between our transport theory approach and the S-matrix formulation of statistical mechanics, we have made the idea of the profile function more convincing. On the other hand limitations of this concept are also better seen. The S-matrix expansion of grand potential ^{1/2} is rapidly convergent for classical systems. In the region, where quantum effects are important the expansion practically fails since one should take into account numerous multiparticle diagrams for realistic description of the system. Analogous situation occurs,

with our approach. To make the idea of the profile function adequate for quantum systems one should include the dependence of this function on the medium, i.e., the unstable particle lifetime should be density and temperature dependent.

We have applied our model to the description of the hadron gas of nucleons, pions and deltas. It has been shown that if one treats the isobars as stable particles their number in the gas is significantly underestimated particularly at the temperature smaller than about 60 MeV.

One of us (St.M.) is grateful to G.Baym for calling his attention to Ref.2.

REFERENCES

1. Mrówczyński St. JINR, E2-85-85, Dubna, 1985; Ann.Phys. in print.
2. Dashen R., Ma S.-K., Bernstein H.J. Phys.Rev., 1969, 187, p.345.
3. Das Gupta S., Mekijan A.Z. Phys.Rep., 1981, 72, p.131.
4. Gaździcki M., Mrówczyński St. Phys.Rev., 1984, C30, p.388.
5. Shimitzu F. et al. Nucl.Phys., 1982, A389, p.445.

Received by Publishing Department
on April 9, 1986.

SUBJECT CATEGORIES OF THE JINR PUBLICATIONS

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches
19.	Biophysics

Денисенко К.Г., Мрувчински С.
Дельта-изобары в адронном газе

E2-86-218

Найдена связь между описанием систем с нестабильными частицами в теории транспорта, предложенной одним из авторов, и описанием таких систем в рамках S-матричной формулировки статистической механики. Изучаются равновесные характеристики адронного газа, состоящего из нуклонов, дельта-изобар и пионов. Принята во внимание конечность ширины распада дельта-изобар. Показано, что если дельта-частицы рассматривать как стабильные, то их количество существенно занижается при температуре ниже 60 МэВ.

Работа выполнена в Лаборатории высоких энергий ОИЯИ.

Сообщение Объединенного института ядерных исследований. Дубна 1986

Denisenko K.G., Mrówczyński St.
Deltas in Hadron Gas

E2-86-218

The relation between the author's transport theory approach to the systems with unstable particles and the S-matrix formulation of statistical mechanics is established. Then, the equilibrium characteristics of classical gas of nucleons, deltas and pions are studied. The finiteness of the delta decay width is taken into account. It is shown that if one treats the delta isobars as stable particles, their number is significantly underestimated at the temperatures smaller than about 60 MeV.

The investigation has been performed at the Laboratory of High Energies, JINR.

Communication of the Joint Institute for Nuclear Research. Dubna 1986