

ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

C-57

E4-87-297

A.A.Chumbalov*, S.S.Kamalov

**OFF-SHELL EFFECTS
IN THE COHERENT π^0 -PHOTOPRODUCTION
OFF NUCLEI**

Submitted to "Physics Letters B"

* Kazakh State University, Alma-Ata

I. INTRODUCTION

In the present paper we discuss the off-shell effects in the coherent π^0 -photoproduction analysed in the framework of the distorted-wave impulse approximation (DWIA). In order to formulate the problem let us write T-matrix of the process in terms of DWIA in the momentum space [1]

$$\langle \underline{q}_c | T(E) | \underline{k} \lambda \rangle = \langle \underline{q}_c | V_{\pi N} | \underline{k} \lambda \rangle + \frac{i}{(2\pi)^3} \int \frac{\langle \underline{q}_c | T'_{\pi N}(E) | \underline{q} \rangle \langle \underline{q} | V_{\pi N} | \underline{k} \lambda \rangle}{E - E(\underline{q}) + i\epsilon} d\underline{q}, \quad (1)$$


where

$$\langle \underline{q} | V_{\pi N} | \underline{k} \lambda \rangle = \int \rho(\underline{p}', \underline{p}) \langle \underline{q}, \underline{p}' | t_{\pi N}(\omega) | \underline{p}, \underline{k} \lambda \rangle d\underline{p} d\underline{p}', \quad (2)$$

\underline{q} and $\underline{p}(\underline{p}')$ are the pion and initial (final) nucleon momenta, respectively, $\underline{k} = E_{\gamma} \hat{k}$ and $\lambda = \pm 1$ are the photon momentum and polarization,

$$\rho(\underline{p}', \underline{p}) = \langle 0 | \sum_{j=1}^A \delta(\underline{p}' - \underline{p}_j) \delta(\underline{p} - \underline{p}_j) | 0 \rangle \quad (3)$$

is the nuclear density, $t_{\pi N}(\omega)$ is the pion photoproduction t -matrix on the free nucleon. Here, the quantity ω has the meaning of the full pion-nucleon (πN) energy in the πN -centre of-mass system when pion is on-shell (i.e. when $E_{\pi} = [m_{\pi} + q_0^2]^{1/2}$). $T'_{\pi N}(E) = T_{\pi N}(E)(A-1)/A$ is the matrix of elastic pion-nuclear scattering that depends on the full energy E for a pion-nuclear (πA) system (i.e. $E = E_{\pi}(q_0) + E_A(q_0)$). This matrix can be determined by solving the πA -scattering problem [2, 3] in the KMT version of the Multiple Scattering Theory [4].



It follows from eqs.(1,2) that to take into account the strong πA -interaction in the final state it is necessary to perform the integration over q from 0 to ∞ . As a consequence, the problem of determination of the $t_{\pi Y}(\omega)$ -matrix in the off-shell region (i.e. when $q \neq q_0$) arises. This problem is closely connected with the choice of the reaction energy ω since different off-shell extrapolation of the $t_{\pi Y}(\omega)$ -matrix can be realized by different determination of the parameter ω in the off-shell region. The main task of the present work is to investigate the sensitivity of the π^0 -photoproduction process to the different choice of ω .

Note that an analogous problem arises in the πA -scattering. In this case it has been shown[5,6] that various assumptions about ω and its relation with the energy E lead to different results in the Δ_{33} -region. We think that coherent π^0 -photoproduction must be more sensitive to different determination of the reaction energy ω since in this process the nonresonant pion S -wave contribution is absent and therefore the resonance (3,3)-multipole dominates. As a consequence, we have a very sharp energy dependence for the $t_{\pi Y}(\omega)$ matrix. In such a situation it is very important to know the correct determination of the reaction energy ω .

II. ON-SHELL PHOTOPRODUCTION AMPLITUDE

For the determination of the $t_{\pi Y}$ -matrix in eq.(2) the expression for the corresponding free nucleon amplitude $\tilde{f}_{\pi Y}$ in the πN c.m. frame is used. The on-shell relation for $t_{\pi Y}$ and $\tilde{f}_{\pi Y}$ is

$$\langle \underline{q}, \underline{p}' | t_{\pi Y}(\omega) | \underline{p}, \underline{k}, \lambda \rangle = -2\pi \delta(\underline{P}_f - \underline{P}_i) \sqrt{\frac{W_i W_f}{E_Y E_N E_N'}} \langle \underline{\tilde{q}} | \tilde{f}_{\pi Y}^{(\lambda)}(\omega) | \underline{\tilde{k}} \rangle. \quad (4)$$

where W_i and W_f are the invariants for YN and πN systems

$$W_i^2 = (E_Y + E_N(p))^2 - \underline{P}_i^2; \quad W_f^2 = (E_\pi(q) + E_N(p'))^2 - \underline{P}_f^2 \quad (5)$$

with $\underline{P}_i = \underline{k} + \underline{p}$, $\underline{P}_f = \underline{q} + \underline{p}'$ and $E_N(p) = (M_N^2 + p^2)^{1/2}$. The pion ($\underline{\tilde{q}}$) and photon ($\underline{\tilde{k}}$) momenta in the c.m. frame are connected with the corresponding momenta in an arbitrary frame by the Lorents transformation

$$\underline{\tilde{q}} = \underline{q} + \left[\frac{\underline{P}_f \cdot \underline{q}}{E_\pi(q) + E_N(p') + W_f} - E_\pi(q) \right] \frac{\underline{P}_f}{W_f}; \quad \underline{\tilde{k}} = \underline{k} + \left[\frac{\underline{P}_i \cdot \underline{k}}{E_Y + E_N(p) + W_i} - E_Y \right] \frac{\underline{P}_i}{W_i}. \quad (6)$$

The general expression for $\tilde{f}_{\pi Y}$ and its partial decomposition is well-known[7]. Here, we use the spin-independent part of $\tilde{f}_{\pi Y}$ since only it contributes in the coherent π^0 -photoproduction on zero-spin nuclei. Keeping only p - and d -pion partial waves, one obtains

$$\langle \underline{\tilde{q}} | \tilde{f}_{\pi Y}^{(\lambda)}(\omega) | \underline{\tilde{k}} \rangle = [(2M_{1+} + M_{1-}) + 3\tilde{\alpha}(3M_{2+} + 2M_{2-})] [\underline{\tilde{q}} \times \underline{\tilde{k}}] \cdot \underline{\epsilon}_\lambda, \quad (7)$$

where $\underline{\tilde{q}}$ and $\underline{\tilde{k}}$ are the unit pion and photon momenta in the πN c.m. frame, respectively, $\tilde{\alpha} = \underline{\tilde{q}} \cdot \underline{\tilde{k}}$. The multipoles $M_{\ell\pm}(\omega)$ are taken from refs.[7,8].

For averaging over nucleon momentum distribution in the nucleus for the calculation of $V_{\pi Y}$ (see eq.(2)), it is convenient to perform the following substitution

$$\underline{p} = -\frac{\underline{k}}{A} - \frac{A-1}{2A}(\underline{k} - \underline{q}) + \underline{v} \equiv \underline{p}_i^{\text{eff}} + \underline{v}, \quad (8a)$$

$$\underline{p}' = -\frac{\underline{q}}{A} + \frac{A-1}{2A}(\underline{k} - \underline{q}) + \underline{v} \equiv \underline{p}_f^{\text{eff}} + \underline{v}. \quad (8b)$$

According to ref.[6], for zero-spin nuclei linear in v terms in $t_{\pi Y}$ give zero contributions and quadratic ones in v are of the order $(m_\pi/M)^2 \sim 0.02$. Consequently, for our purpose it is sufficient to neglect v in (7) and take $p \approx p_i^{\text{eff}}$, $p' \approx p_f^{\text{eff}}$

(the factorization approximation). Then, the plane wave part $V_{\pi Y}$ can be expressed as

$$\langle \underline{q} | V_{\pi Y} | \underline{k} \lambda \rangle = -2\pi A \left[\frac{W_i W_f}{E_Y E_\pi(q) E_N(p) E_N(p')} \right]^{1/2} \langle \tilde{q} | \tilde{f}_{\pi Y}^{(\lambda)}(\omega) | \tilde{k} \rangle F_A^{ch}(\underline{Q}) / f_p^{ch}(\underline{Q}), \quad (9)$$

where $F_A^{ch}(\underline{Q})$ and $f_p^{ch}(\underline{Q})$ are the nuclear and proton charge form factors, respectively, $\underline{Q} = \underline{k} - \underline{q}$ is the transferred momentum.

Note that the factorization approximation guarantees the momentum and energy conservation in the πA , πN and YN systems simultaneously when $q = q_0$. However, in the off-shell region (where $q \neq q_0$) the total energy for the πN -system disagrees with the total energy for the YN -system (i.e. $W_i \neq W_f$ when $q \neq q_0$). In this region the physical meaning of the parameter ω is uncertain.

III. OFF-SHELL EFFECTS. RESULTS AND DISCUSSION

As it has been noted above, the connection (4) between the matrices $t_{\pi Y}(\omega)$ and $\tilde{f}_{\pi Y}(\omega)$ is correct only in the on-shell sense. While computing the principal value of integral in (1), one postulates that eq.(4) is relevant in the off-shell region as well. However, the uncertainty in the definition of the energy ω in the off-shell region remains. In what follows, to study the effects of different definition of ω , we shall use the following alternative expressions:

$$\omega_0 = W_i \approx [(E_\pi(q_0) + E_N(p'))^2 - \underline{p}'^2]^{1/2} \quad (10a)$$

$$\omega_1 = [m_\pi^2 + M_N^2 + 2E_\pi(q_0)E_N(p') - 2\underline{q} \cdot \underline{p}']^{1/2} \quad (10b)$$

$$\omega_2 = W_f = [(E_\pi(q) + E_N(p'))^2 - \underline{p}'^2]^{1/2}. \quad (10c)$$

According to ref.[6], the reaction energy ω may be determined also as $\omega_3 = (W_i + W_f)/2$ and $\omega_4 = \sqrt{W_i W_f}$.

All definitions for ω_i ($i = 0-4$) are on-shell equivalent but they provide us with rather different values in off-shell-region. For the illustration of this fact the dependence of ω_0 , ω_1 and ω_2 on pion momentum q is plotted in Fig.1. in the case of π^0 -photoproduction off ^{12}C at $E_Y^{\text{LAB}} = 290$ MeV and $\theta_\pi = 25^\circ$ (in the πA c.m. frame $E_Y = 283$ MeV, $q_0 = 1.26 \text{ fm}^{-1}$ and $\omega(q_0) = 1200$ MeV).

The off-shell behaviour of the partial amplitudes $M_{\ell\pm}(\omega)$ at the fixed reaction energy ω_i ($i = 0-4$) may be determined, e.g., from the separable model of πN -interaction[9] like it was done in the πA -scattering problem[2,3]

$$M_{\ell\pm}(\omega) = M_{\ell\pm}(\omega_i) g_{\pi N}^{(\ell)}(\tilde{q}) / g_{\pi N}^{(\ell)}(\tilde{q}_i). \quad (11)$$

where

$$g_{\pi N}^{(\ell)}(\tilde{q}) = q^\ell / (1 + \alpha q^2)^2, \quad \alpha = 0.224 \text{ fm}^2. \quad (12)$$

In eqs. (11,12) \tilde{q}_i is the pion momentum in πN c.m. frame that corresponds to the energy ω_i for the πN -system. The appropriate threshold behaviour of the amplitude $M_{\ell\pm}(\omega)$ as $\tilde{q} \rightarrow 0$ is guaranteed by the term \tilde{q}^ℓ in eq.(12).

Notice that for $\omega_2 = W_f(q)$, one may not introduce the factor $g_{\pi N}^{(\ell)}(q)$ because in this case $\tilde{q} = q_2$. The right threshold behaviour of $\tilde{f}_{\pi Y}$ and its cutting off as $\tilde{q} \rightarrow \infty$ are provided by the energy dependence of $M_{\ell\pm}(\omega)$ multipoles for which one has[7]

$$M_{\ell\pm} \xrightarrow{\tilde{q} \rightarrow 0} \tilde{q}^\ell; \quad M_{\ell\pm} \xrightarrow{\tilde{q} \rightarrow \infty} \frac{1}{\omega^{\ell+1}}; \quad M_{\ell\pm} \xrightarrow{\tilde{q} \rightarrow \infty} \frac{1}{\omega^\ell}. \quad (13)$$




Fig. 1. q -dependence of different definition of the reaction energy ω in the off-shell region corresponding to the on-shell value $q_0 = 1.26 \text{ fm}^{-1}$ ($k_L \equiv E_Y^{\text{LAB}} = 290 \text{ MeV}$).

The energy dependence of the multipole $M_{1+}(\omega)$ dominant in (7) is plotted in Fig. 2. One can see that the region most sensitive to the choice of energy ω may be $E_Y^{\text{LAB}} \sim 290\text{--}340 \text{ MeV}$ where the real and imaginary parts of this multipole have sharp energy dependence. Our DWIA results* corresponding to the different definition for the off-shell behaviour of the parameter ω (see Fig.2) are depicted in Figs.3 and 4. As it was expected, Δ_{33} -region is the most sensitive to the choice of ω . In this region results may differ as much as 1.5-2 times. This sensitivity decreases with E_Y^{LAB} .

*The results of this paper differ from our previous results (see ref./14/). This is due to several reasons: 1) earlier, multipoles $M_{l+}(\omega)$ were calculated at fixed energy $\omega(q_0)$, now ω is the variable of integration; 2) in the Δ_{33} -resonance region we use the new BD-amplitude[8] instead of BDW [7]; 3) we use the new πA -optical potential[3] describing not only the differential cross sections[15] but also σ_{TOT} and σ_{el} for the elastic πA -scattering.




Fig. 2. Energy dependence of the $M_{1+}(\omega) = [M_{1+}^{(1)} + 2M_{1+}^{(3)}] / 3$ multipole [8].




Fig. 3. Angular distribution for $^{12}\text{C}(\gamma, \pi^+)^{12}\text{C}$ at the photon energy $k_L = 290 \text{ MeV}$ obtained by using the reaction energy $\omega = \omega_2$ (dash-dot), $\omega = \omega_1$ (dashed) and $\omega = \omega_2$ (solid). The data are from ref. [10].




Fig. 4. The same as in Fig. 3 at $k_L = 230$ Mev. The result corresponding to the choice $\omega = \omega_0$ is not shown because it only slightly differs from that corresponding to $\omega = \omega_2$. Dotted line is the result of the plane wave impulse approximation.

In our opinion $\omega = \omega_2$ is the most consistent choice. This is not only because this choice provides us with the best agreement with the experimental data from ref. [10]. Actually, such a conclusion may be considered as a consequence of the Relativistic Potential theory [11] with the help of which one can determine the off-shell relation between the $t_{\pi N}$ -matrix in an arbitrary frame and the corresponding amplitude $\tilde{f}_{\pi N}$ in πN c.m. frame. Such an expression was obtained in [12] for the πN -scattering matrix. Generalizing this method for the two-potentials problem [13] one can see that in expression (4) it is necessary to replace the amplitude $\tilde{f}_{\pi N}$ by the auxiliary matrix $\tilde{f}_{\pi N}$ connected with each other as follows:

$$\langle \tilde{q}' | \tilde{f}_{\pi N}^{(\lambda)}(\omega) | \tilde{k} \rangle = \langle \tilde{q}' | \tilde{f}_{\pi N}(W_f) | \tilde{k} \rangle - \frac{1}{(2\pi)^2} \int \frac{dq'}{\mu(q')} \langle \tilde{q}' | f_{\pi N}(W_f(q)) | q' \rangle \quad (14)$$

$$\times \langle q' | \tilde{f}_{\pi N}^{(\lambda)}(W_f(q)) | \tilde{k} \rangle \left[\frac{1}{W_f - W_f(q) + i\epsilon} - \frac{1}{\omega - W_f(q) + i\epsilon} \right],$$

where $\mu(q)$ is the reduced mass of πN -system, $W_f = E_\pi(\tilde{q}) + E_N(\tilde{q}) = \sqrt{[E_\pi(q) + E_N(p')]^2 - p^2}$ has the meaning of the eigenvalue of the relativistic free Hamiltonian of the πN -system ($\hat{h}_{\pi N}^0 = \hat{h}_\pi + \hat{h}_N$) in the c.m. frame, $f_{\pi N}$ is the πN -scattering amplitude. Note that we neglect in (14) the terms of $\mathcal{P}_i^2/W_i W_f$ order and use the first Born approximation for the electromagnetic interaction.

It can easily be seen from eq. (14) that if we set up $\omega = W_f$ the contribution of the second term in eq. (14) will be zero. As a result, we obtain the simplest off-shell connection (4) between the $t_{\pi N}$ -matrix in an arbitrary frame and the corresponding amplitude $\tilde{f}_{\pi N}$ in the πN -c.m. frame (half off-shell connection [12]).

IV. SUMMARY

We have demonstrated the strong sensitivity of the coherent π^0 -photoproduction off nuclei to the choice of the reaction energy ω for the elementary $t_{\pi N}(\omega)$ -matrix in the off-shell region. The main reason for such a sensitivity is the resonant energy dependence of the $t_{\pi N}$ -matrix. The best agreement with the experimental data was obtained when ω was chosen as the eigenvalue of the free relativistic Hamiltonian for the πN -system (i.e. $\omega^2 = (E_\pi(q) + E_N(p'))^2 - (q+p')^2$). This conclusion is consistent with the results of the Relativistic Potential theory [11, 22].

Note that in our calculations performed for the charged pion photoproduction off ^{16}O , ^{12}C and ^{10}B we have observed only 10-20% difference between the results corresponding to various choices of the energy ω . This is mainly due to the

fact that in the case of charged pion photoproduction of the nonresonant S -wave E_{0+} multipole dominates and consequently $t_{\pi\gamma}$ has smooth energy dependence.

The authors are grateful to R.A.Eramzhyan, M.Gmitro and R.Mach for interest in the work and valuable remarks.

REFERENCES

1. R.A.Eramzhyan, M.Gmitro, S.S.Kamalov and R.Mach. J.Phys. G: Nucl.Phys, 9 (1987) 605.
2. M.Gmitro, J.Kvasil and R.Mach. Phys.Rev. C31 (1984) 1349.
3. M.Gmitro S.S.Kamalov, R.Mach. Preprint INPCSAV, No.2 (1986).
(to be published in Phys.Rev. C).
4. A.Kerman, H.McManus and R.M.Thaler. Ann. of Phys. 8 (1959) 551.
5. T.I.Kopaleishvili, V.S.Shirdladse. Sov. J.Nucl.Phys. 32(1980) 1267.
6. R.Mach. Czech.J.Phys. B33 (1983) 773.
7. F.A.Berends, A.Donnachie, and D.L.Weaver. Nucl.Phys. B4(1967) 1.
8. F.A.Berends and A.Donnachie, Nucl.Phys. B84 (1975) 342.
9. J.T.Londergan, K.M.McVoy, and E.J.Moniz, Ann. of Phys. 78 (1973) 299.
10. J.Arends et al. Z.für Phys. A311 (1983) 567.
11. R.Fong and J.Sucher. J.Math.Phys. 5 (1964) 456.
12. L.Heller, G.E.Bohanon, and F.Tabakin. Phys.Rev. C13 (1976) 742.
13. M.L.Goldberger and K.M.Watson (N.Y.: Wiley) Collision Theory, 1964.

14. S.S.Kamalov, T.D.Kaipov, Phys.Lett. B162, (1985) 260.
15. R.A.Eramzhyan et al. Nucl.Phys. A429 (1984) 403.

Received by Publishing Department
on April 28, 1987.

НЕТ ЛИ ПРОБЕЛОВ В ВАШЕЙ БИБЛИОТЕКЕ?

Вы можете получить по почте перечисленные ниже книги, если они не были заказаны ранее.

Д9-82-664	Труды совещания по коллективным методам ускорения. Дубна, 1982.	3 р. 30 к.
Д3,4-82-704	Труды IV Международной школы по нейтронной физике. Дубна, 1982.	5 р. 00 к.
Д11-83-511	Труды совещания по системам и методам аналитических вычислений на ЭВМ и их применению в теоретической физике. Дубна, 1982.	2 р. 50 к.
Д7-83-644	Труды Международной школы-семинара по физике тяжелых ионов. Алушта, 1983.	6 р. 55 к.
Д2,13-83-689	Труды рабочего совещания по проблемам излучения и детектирования гравитационных волн. Дубна, 1983.	2 р. 00 к.
Д13-84-63	Труды XI Международного симпозиума по ядерной электронике. Братислава, Чехословакия, 1983.	4 р. 50 к.
Д2-84-366	Труды 7 Международного совещания по проблемам квантовой теории поля. Алушта, 1984.	4 р. 30 к.
Д1,2-84-599	Труды VII Международного семинара по проблемам физики высоких энергий. Дубна, 1984.	5 р. 50 к.
Д17-84-850	Труды III Международного симпозиума по избранным проблемам статистической механики. Дубна, 1984. /2 тома/	7 р. 75 к.
Д10,11-84-818	Труды V Международного совещания по проблемам математического моделирования, программирования и математическим методам решения физических задач. Дубна, 1983	3 р. 50 к.
	Труды IX Всесоюзного совещания по ускорителям заряженных частиц. Дубна, 1984 /2 тома/	13 р. 50 к.
Д4-85-851	Труды Международной школы по структуре ядра, Алушта, 1985.	3 р. 75 к.
Д11-85-791	Труды Международного совещания по аналитическим вычислениям на ЭВМ и их применению в теоретической физике. Дубна, 1985.	4 р.
Д13-85-793	Труды XII Международного симпозиума по ядерной электронике. Дубна, 1985.	4 р. 80 к.
Д3,4,17-86-747	Труды V Международной школы по нейтронной физике. Алушта, 1986.	4 р. 50 к.

Заказы на упомянутые книги могут быть направлены по адресу:
101000 Москва, Главпочтамт, п/я 79
Издательский отдел Объединенного института ядерных исследований

Чумбалов А.А., Камалов С.С.

E4-87-297

Эффекты схода с энергетической поверхности в процессах когерентного фоторождения π^0 -мезонов на ядрах

Продемонстрирована сильная чувствительность процесса когерентного фоторождения π^0 -мезонов на ядрах к разному выбору энергии реакции ω в элементарной амплитуде $t_{\pi\gamma}(\omega)$. Разные предположения о поведении ω во внеэнергетической области могут изменять дифференциальные сечения в 1.5-2 раза. Наилучшее согласие DWIA-результатов с экспериментальными данными получено при ω равном собственному значению свободного релятивистского гамильтониана пион-нуклонной системы. Такой результат находится в согласии со следствиями релятивистской потенциальной теории.

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1987

Chumbalov A.A., Kamalov S.S.

E4-87-297

Off-Shell Effects in the Coherent π^0 -Photoproduction off Nuclei-

The strong sensitivity of the coherent π^0 -photoproduction to the choice of the reaction energy ω in the elementary $t_{\pi\gamma}(\omega)$ -matrix is demonstrated. The best agreement of the DWIA-results with the experimental data is achieved when ω is chosen as an eigenvalue of the free relativistic Hamiltonian of the πN -system. This is in agreement with the consequences of the relativistic potential theory.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1987