

S.B.Drenska, A.I.Georgieva*, M.I.Ivanov*, S.Sht.Mavrodiev*, P.P.Raychev*, R.P.Roussev*

A PHENOMENOLOGICAL UNIFIED DESCRIPTION OF THE GROUND BAND ENERGY LEVELS OF ROTATIONAL NUCLEI

*Institute for Nuclear Research and Nuclear Energy, 1784 Sofia, Bulgaria

1988

Almost all of the existing nuclei show some degree of collectivity. As a result a lot of collective models and interpretations have been developed. However, they differ in language and formalism. The purpose of this paper is to unify the description of the rotational low lying levels in the nuclei, by using only the universal nuclear characteristics specific for a given nucleus.

The description should include a large region of nuclei. The latter can be done within the classification scheme proposed in '1' and 2'. According to this scheme all even-even nuclei with $Z \ge 20$, $A - Z \ge 20$ are systematized in terms of boson representations of the Sp(24,R) – algebra^{'3'}. These nuclei, whose valence nucleons occupy the same major nuclear shell are united in two The symplectic multiplets symplectic multiplets. are defined bv $(N_p, N_n \mid N_p, N_n)_{\pm}$, where N_p, N_n and N_p, N_n' are the numbers of protons and neutrons of the double magic nuclei at the beginning of two consecutive shells. The relevant multiplet is noted by + (-), when the total number of valence bosons is even (odd). Each nucleus corresponds to a definite cell in the symplectic multiplets, given by the irreducible unitary representations (IUR) of the group $U_{\pi}(6) \oplus U_{\mu}(6)$ according to the reduction

$$\operatorname{Sp}(24, \mathbb{R}) \xrightarrow{F_0} U(6, 6) \xrightarrow{N} U_{\pi}(6) \oplus U_{\nu}(6).$$
(1)

It is well known that the last group in (1) is the group of dynamical symmetry of IBM-2⁴⁴. Its UIT's are given by the proton and neutron boson numbers N_{π} and N_{ν} , which can be found by counting the valence proton and neutron pairs. By means of these quantum numbers the quantities N – total boson number and F_{0} – third projection of the F – spin are:

$$N = N_{\pi} + N_{\nu}$$
, $F_{o} = (N_{\pi} - N_{\nu})/2$. (2)

Obviously, we have used some group theoretical apparatus in terms of quantities introduced in one of the most popular models IBM-2. But we would like to underline that the physical meaning of these quantities is easily kept in mind, so their direct relation to the nuclear characteristics as number of nucleons of both types N_p , N_n is as simple as that:

$$N = (A - A^{mag})/2, \qquad F = (M_T - M_{T}^{mag})/2, \qquad (3)$$
1
1

where $A = N_p + N_n$ is the mass number and $M_T = (N_p - N_n)/2$ is the third projection of the isospin, A^{mag} , M_T^{mag} are the mass number and third projection of the isospin of the double magic nucleus at the beginning of the shell.

An empirical investigation of the energies of the first excited 2^+ levels of the nuclei from a given multiplet was completed in^{/1,2/}. The curves obtained show a stable periodical structure in the F₀-multiplets of a given shell. Follow up, all the excited states of the ground band present similar behaviour up to the occurrence of the back-banding in the spectrum. This periodical structure is observed in all shells investigated and is especially stable in the case of heavy and superheavy nuclei.

In the most simplified picture the well deformed nuclei, which demonstrate rotational spectra are to be found far from the spherical magic nuclei. In the case of our classification they correspond for each F_o to the lowest almost constant part of the N-dependence curves of the 2^+ energies. In order to specify this we tried to describe all the energies in the ground bands by means of the classical formulae of the type

 $E_{rot} = L(L+1)/2J,$ (4)

where we assume, that the inertion parameter 1/2J is a polynomial up to a second degree in the independent combinations of N_{π} , N_{ν} , N_{p} , N_{n} , N'_{p} , N'_{n} .

On the other hand the same assumptions are to be reached using the idea of the chain of groups, named general dynamical groups (GDGS), along which one can classify the nuclei. As it was already mentioned we classify the nuclei according to the chain (1), so the coefficient in the Hamiltonian will depend on the invariants of the groups in this chain; in this particular case -N, F_o and the numbers defining the relevent Sp(24, R) – multiplet, namely N_p , N_n , N_p' , N_n' .

The classification chain will lead to the group of dynamical symmetry (DG) whose reduction will generate the spectrum of the already specified nucleus. The terms of the Hamiltonian itself are to be the first and second order Casimir operators of a subchain of DG. As we are trying to describe the well deformed nuclei, an obvious candidate is the chain

 $U(6) \supset SU(3) \supset SO(3),$

which describes their spectrum in most algebraic models. On the states of the ground band, defined as: $|[\lambda]_6$, $[\lambda, 0]_3$, L > this type of Hamiltonian has eigenvalues

$$E_{rot} = F(N, F_{o}, N_{p}, N_{n}, N_{p}', N_{n}') L(L+1).$$
(5)

In order to describe the rotational energies using (5) it was necessary to introduce two new quantum numbers. These are N_{π} and N_{ν} , defining the numbers of valence proton and neutron boson holes for the relevant nucleus. For convenience the following system of the independent quantum numbers was chosen:

$$N = (N_{\pi} + N_{\nu}), \quad F_{o} = (N_{\pi} - N_{\nu})/2, \quad A_{p} = N_{p} + N_{p}'$$

$$N' = (N_{\pi}' + N_{\nu}'), \quad F_{o} = (N_{\pi}' - N_{\nu}')/2, \quad A_{N} = N_{n} + N_{n}'.$$
(6)

Thus the mathematical expression of the energy is

$$E_{rot} = L(L+1) \{C_{o} + C_{1}N + C_{2}N' + C_{3}F_{o} + C_{4}F_{o}' + C_{5}A_{P} + C_{6}A_{N} + C_{7}N^{2} + C_{8}N'^{2} + C_{9}F_{o}^{2} + C_{1o}F_{o}'^{2} + C_{11}NF_{o} + (7) + C_{12}N'F_{o}' + C_{13}A_{P}^{2} + C_{14}A_{N}^{2} + C_{15}A_{P}A_{N}\}.$$

All the rotational nuclei belonging to the multiplets $(50, 50 | 82, 82)_{\pm}$; $(50, 82 | 82, 126)_{\pm}$; $(82, 126 | 126, ...)_{\pm}$ have been considered. It is obvious that we are looking for the solution of the overdetermined system of equations:

$$E_{rot}(L,N,N',F_{o},F',A_{P},A_{N},(C_{i})) = E_{rot}^{exp}(L).$$
(8)

 $E_{rot}^{exp}(L)$ are the experimental values of the energies $^{/5/}$ of the ground bands for $L = 2^+, 4^+, \ldots$ up to the occurrence of back-banding.

Using an autoregularized iterational method of Gauss – Newton type^{'6'} the coefficients C_i are evaluated for 328 excited states of 70 nuclei. As a result for the rotational energies E_{rot} (L) the following simple formulae were obtained:

$$E_{rot}(L) = L(L+1) \{ D_1(N+N') + D_2(F_0 + F_0') + D_3(N^2 + N'^2/2) + D_4(F_0^2 - F_0'^2/2) + D_5A_N \}.$$
(9)

The values of the parameters D_i $(i=1,\ldots,5)$ are given in Table 1 together with their uncertainties $\pm \Delta D_i$.

		$\frac{Table \ 1}{\pm \Delta D_{i}}$		
i	Di			
1	16.44	0.37		
2	-36.82	3.06		
3	1.92	0.06		
4	-9.67	0.57		
5	-3.38	0.04		

Obviously, the values of (N + N'), $(F_0 + F'_0)$ and A_N are constants for the nuclei belonging to a given Sp(24, R)-multiplet. The necessity of introducing the quantum numbers N', F'_0 reveals the assymmetry of the transition from the spherical nuclei to the well deformed ones and vice versa. Hence, for the manifestation of the nuclear collectivity it is important not only how many valence nucleons the relevant nucleus has but as well how

more you need to fill up the shell. The values of the coefficients in front of the nuclear characteristics show the different contribution of each one to the parameter of inertia J.

In order to investigate this, it is more convenient to transform (9), reduced for the nuclei only from a given shell, in the following way:

 $E_{tot}^{shell} = \{ (D_3 + D_4/4) (N_{\pi}^2 + N_{\nu}^2) + (D_3 - D_4/4) (N_{\pi}^2 + N_{\nu}^2) + (D_3 - D_4/4) (N_{\pi}^2 + N_{\nu}^2) \}$ + $2(D_{2} - D_{4}/4) N_{\pi}N_{\nu} + 2N_{\pi}N_{\nu}(D_{2} + D_{4}/4) L(L + 1);$

or using the coefficients from Table 1 for the rotational parameter in a given shell one has

 $1/2J = 8.675N_{\pi}N_{\nu} - 0.995N_{\pi}N_{\nu} - 0.4985(N_{\pi}^2 + N_{\nu}^2) + 4.3375(N_{\pi}^2 + N_{\nu}^2)$

Obviously, the most important part in the rotational motion plays the interaction between the proton and neutron systems, depending on the product $N_{\pi}N_{\mu}^{\prime \prime \prime \prime \prime}$. And as a new result one can consider the rather big coefficient in front of the term $(N_{\pi}^{2} + N_{\nu}^{2})$ depending on the number of boson holes, or which is the same, on the number of valence nucleons which are needed to fill up the shell.

In Table 2 the experimental $5^{1/2}$ and theoretical values of the energies for each L are compared, for the nucleus from each multiplet with the longest rotational band. It must be mentioned that for the unknown values of N_n and N'_n for the third multiplet we use those values that the shell model predicts, i.e. $N_{p} = 126$ and $N_{n} = 184$.

The empirical formulae (9) obtained on the basis of the large investigation of the energies of the nuclei classified in a simple symplectic scheme can be used as a tool for evaluating the degree of the deformation or the validity of the rotational description of the nuclear spectra.

(5	50,50 82	2,82)		(50,50 82,82)						
6.0	Nd ¹²⁸ 68	1		60 Nd ¹³⁰						
1.	Е "хр	E _{th}		Ŀ	L _{yxp}	E t. h				
2	0.134	0.120		2	0.159	0.132				
4	0.426	0.401		4	0.485	0.441				
U.	0.850	0.842		Ċ	0.940	0.927				
				(50,82 82,126]						
1562	Sni 94			$\frac{162}{60}$ Nd $\frac{2}{2}$						
L	E	Eth		L	Е	E th				
2	0.076	0.061	-	2	0.076	0.069				
4	0.250	0.225		4	0.241	0.228				
6	0.518	0.471		Ċ	0.488	0.480				
8	0.878	0.808		8	0.810	0.822				
^{1 b o} Gd _{9 2}				164 68 4 6						
L	E	E thr		L	E	E				
2	0.089	0.076	- -	2	0.091	0.078				
4	0.288	0.253		4	0.299	0.260				
6	0.585	0.532		6	0.614	0.545				
[:] 8	0.965	0.912		8	1.025	0.935				
10	1.416	1.394		10	1.518	1.428				
12	1.924	1.977		12	2.083	2.025				
180	W	L	- '	¹⁸⁴ 0s						
	E	E		Ĺ	E	E				
2	0.104	0.08	1	2	0.120	0.098				
4	0.338	0.29		4	0.384	0.327				
6	0.688	0.62		6	0.774	0.686				
8	1.138	1.07		8	1.275	1.176				
10	1.660	1.64		10	1.870	1.797				

4

Table 2

172	Dy				¹⁷⁶ Yb				Table	2 (contin.)	
	E x	Р	E _{th}	L.	E exp		E th	L	E exp	E Lh	
2	0.08	1	0.072	2	0.082	0.	.078	2	0.093	0.083	
4	0.26	6	0.240	4	0.272	0.	261	4	0.307	0.277	
6	0.54	9	0.504	6	0.565	0.	548	6	0.632	0.582	
8	0.92	1	0.864	8	0.954	0.	940	8	1.059	0.998	
10	1.37	5	1.320	10	1.431	1.	436	10	1.571	1.524	
12	1.90	3	1.871	12	1.985	2.	036	12	2.151	2.162	
14	2.49	4	2.519	14	2.602	2.	741	14	2.778	2.910	
(82,126)126,184)											
232	Th 142				230 92140	,			¹ ²⁴⁴ Pu 94 ¹⁵⁰		
L	E _{& x}	P.	E _{th}	L	E	t	Ë th	L	Eexp	E _{th}	
2	0.04	9	0.043	2	0.045	0.	.040	2	0.046	0.041	
4	0.16	2	0.143	4	0.148	0.	133	4	0.156	0.137	
6	0.33	3	0.300	6	0.307	0.	279	6	0.318	0.288	
8	0.55	7	0.515	8	0.518	0.	479	8	0.535	0.493	
10	0.82	27	0.787	10	0.776	Ο.	. 731	10	0.801	0.754	
12	1.13	7	1.116	12	1.077	1.	.037	12	1.114	1.069	
14	1.48	3	1.502	14	1.415	1.	. 396	14	1.467	1.439	
16	1.85	9	1.946	16	1.788	1.	808	16	1.858	1.864	
$\frac{250}{98}Cf = \frac{250}{152} I = \frac{250}{100} I = \frac{1}{150}$											
		L	E •xp	E	L h	L	t.	хр	E		
		2	0.043	0.0	042	2	0.0	946	0.049		
		4 0.142 0.140 4		0.1	60	0,165					
		6 0.296 0.294 6 0		0.3	32	0.346					
		8	0.509	0.9	503	8	0.5	064	0.593		

ACNOWLEDGEMENTS

The authors are very grateful to Dr.D. Karadjov for his permanent interest and useful discussions and to the Laboratory of Theoretical Physics of JINR, Dubna, for the opportunity to realize the calculations.

This paper is supported in part by Contract No.330 with the Ministry of Science, Culture and Education, Sofia, Bulgaria.

REFERENCES

۰ ۱۰

Ù.

ł

....

۰,۶

1

- I. Georgieva A.I., Ivanov M.I., Raychev P.P., Roussev R. Communication of JINR E4-87-927, 1987, Dubna.
- 2. Georgieva A.I., Ivanov M.I., Raychev P.P., Roussev R. Comp. Rend. Bulg. Acad. Sci., 1988, 41, No.2, 55, p.
- 3. Georgieva A.I., Ivanov M.I., Raychev P.P., Roussev R. Int. J. Theor. Phys., 1985, 25, p.205.
- 4. Arima A., Otsuka T., Iachello F., Talmi I. Phys. Lett., 1977, 66B, p.205.
- 5. Sakai M. Atomic Data: Nucl. Data Tables, 1984, 31, p.399.
- 6. Aleksandrov L. Math. Phys. & Comp. Math., 1971, 11, No.1, p.36.
- 7. Casten R.F., Von Brentano P., Gellberg A., Harter H. J.Phys. G.: Nucl.Phys., 1986, 12, p.711.

Received by Publishing Department on December 28, 1988.

6

WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices — in US \$, including the packing and registered postage.

D2-84-366	Proceedings of the VII International Conference on the Problems of Quantum Field Theory. Alushta, 1984.	11.00
D1,2-84-599	Proceedings of the VII International Seminar on High Energy Physics Problems. Dubna, 1984.	12.00
D17-84-850	Proceedings of the III International Symposium on Selected Topics in Statistical Mechanics. Dubna, 1984 (2 volumes).	22.00
	Proceedings of the IX All-Union Conference on Charged Particle Accelerators. Dubna, 1984. (2 volumes)	25.00
D11-85-791	Proceedings of the International Conference on Computer Algebra and Its Applications in Theoretical Physics. Dubna, 1985.	12.00
D13-85-793	Proceedings of the XII International Symposium on Nuclear Electronics, Dubna, 1985.	14.00
D4-85-851	Proceedings of the International School on Nuclear Structure Alushta, 1985.	11.00
D1,2-86-668	Proceedings of the VIII International Seminar on High Energy Physics Problems, Dubna, 1986 (2 volumes)	23.00
D3,4,17-86-747	Proceedings of the V International School on Neutron Physics. Alushta, 1986.	25.00
Ð9.87.105	Proceedings of the X-All-Union Conference on Charged Particle Accelerators. Dubna, 1986 (2 volumes)	25.00
D7-87-68	Proceedings of the International School-Seminar on Heavy Ion Physics. Dubna, 1986.	25.00
D2-87-123	Proceedings of the Conference "Renormalization'Group-86". Dubna, 1986.	12.00
D4-87-692	Proceedings of the International Conference on the Theory of Few Body and Quark-Hadronic Systems. Dubna, 1987.	12.00
D2-87-798	Proceedings of the VIII International Conference on the Problems of Quantum Field Theory, Alushta, 1987.	10.00
D14-87-799	Proceedings of the International Symposium on Muon and Pion Interactions with Matter. Dubna, 1987.	13.00
D17-88-95	Proceedings of the IV International Symposium on Selected Topics in Statistical Mechanics, Dubna, 1987.	14 00
E1,2-88-426	Proceedings of the 1987 JINR-CERN School of Physics. Varna, Bulgaria, 1987.	14.00
Orders 1	or the above-mentioned books can be sent at the address: Publishing Department, JINR Head Post Office. P.O.Box 79 101000 Moscow. USSR	

0

Дренска С.Б. и др. Е4-88-906 Единое феноменологическое описание уровней основной полосы ротационных ядер На основании симплектической классификации ядер получена простая феноменологическая пятипараметрическая формула для единого описания возбужденных состояний основной полосы хорошо деформированных четно-четных ядер. Теоретические результаты сравниваются с экспериментальными энергиями 328 уровней 70 ядер.

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Сообщение Объединенного института ядерных исследований. Дубна 1988

Drenska S.B. et al. E4-88-906 A Phenomenological Unified Description of the Ground Band Energy Levels of Rotational Nuclei

On the basis of a symplectic classification of nuclei a phenomenological five-parameter formulae for a unified description of the lowlying excited states of the ground bands of the well deformed eveneven nuclei has been derived. The theoretical results are compared with the experimental data for 328 levels of 70 nuclei.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Communication of the Joint Institute for Nuclear Research. Dubna 1988