

Объединенный институт ядерных исследований

дубна

7 65

E5-88-622

V.I.Inozemtsev, N.A.Kostov

NEW INTEGRABLE SYSTEMS OF INTERACTING NONLINEAR WAVES

Submitted to "Physics Letters A".

1988

In the physical science the following Hamiltonian system

$$i\frac{d}{dt} u_{j} + \gamma |u_{j}|^{2} u_{j} + \sum_{k=1}^{n} M_{jk} u_{k} = 0 , j=1,..,n$$
 (1)

is now widely recognized as a generic model of self-trapping in condensed matter physics. In the case n=2 this system is completely integrable in the Liouville sense, due to the existence of the following conserved quantities H=- $\sum_{k=1}^{n} |u_{j}^{4}| - \sum_{k=1}^{n} M_{j,k} u_{j}u_{k}$

N=∑ {u_j²} j,k j,k j^k N=∑ {u_j²} which are in involution. In the case of n>2 there is j=1 conjecture that this system is not integrable.

The system (1) is a very special case of the following general vector equations

$$\mathbf{i} \frac{\mathbf{d}}{\mathbf{d}t} \mathbf{u}_{j} = \sum_{\mathbf{k}} \mathbf{r}_{j\mathbf{k}}^{(1)} \mathbf{u}_{\mathbf{k}} + \sum_{\mathbf{k},1} \mathbf{r}_{j\mathbf{k}1}^{(2)} \mathbf{u}_{\mathbf{k}} \mathbf{u}_{1} + \sum_{\mathbf{k},1,m} \mathbf{r}_{j\mathbf{k}1m}^{(3)} \mathbf{u}_{\mathbf{k}} \mathbf{u}_{1} \mathbf{u}_{m}^{+} \dots$$
(2)

which describe nonlinear interaction of n waves. These equations have a large number of applications in condensed matter physics, hydrodynamics, metereology, molecular dynamics, etc. The structure of constant tensors $\gamma^{(\alpha)}$, $\alpha=1,\ldots,\theta$ is determined by the particular physics of the processes under investigation.

After the pioneer work [2] many works of physical and mathematical character [3-7] have been devoted to the problems of classification of symmetries, reductions, and applications of the system (2). As special cases of the integrable equations of type (2) we may identify almost all finite dimensional integrable dynamical systems known at the present time, for example, one dimensional reductions of generalized matrix nonlinear Schrodinger equations [5], Toda lattice dynamical systems [6], equations of motion of n-dimensional rigid body in the external gravitational and electromagnetic fields [6] and many others.

The aim of the present article is the construction and analysis of reductions leading to two new integrable dynamical systems of the special type which belong to the class (2). One of these in the two-dimensional case is isomorphic to the nonperiodical Toda lattice; the other in the case n= 2 is the

1



so called dymer system equations [1] In the later case we obtain the general solutions using the finite-gap integration method. These solutions have been obtained in [1] by direct methods.

We start the analysis of the dymer equations (system (2) with n=2) with the Lax representation

$$\frac{d}{dt} L = [L,A], \qquad (3)$$

where the matrices L,A have the following form

$$\mathbf{L} = \begin{bmatrix} 0 & \boldsymbol{\varepsilon} & \mathbf{u}_1 & \mathbf{0} \\ \boldsymbol{\varepsilon} \lambda & 0 & \mathbf{0} & \mathbf{u}_2 \\ \mathbf{i} \mathbf{u}_1^* & 0 & 0 & -\boldsymbol{\varepsilon} / \lambda \\ \mathbf{0} & \mathbf{i} \mathbf{u}_2^* & -\boldsymbol{\varepsilon} & \mathbf{0} \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} -1 |u_1|^2 - \epsilon^2 \lambda & 0 & 0 & -\epsilon u_1 / \lambda - \epsilon u_2 \\ 0 & -1 |u_2|^2 - \epsilon^2 \lambda & -\epsilon u_1 \lambda - \epsilon u_2 & 0 \\ 0 & -1 \epsilon u_1^* - 1 \epsilon u_2^* / \lambda & 1 |u_1|^2 + \epsilon^2 / \lambda & 0 \\ -1 \epsilon \lambda u_2^* - 1 \epsilon u_1^* & 0 & 0 & 1 |u_2|^2 + \epsilon^2 / \lambda \end{bmatrix}$$

The central idea of the method of finite-gap integration [8] is the construction of the Baker-Akhiezer function (BA-function). By definition the BA-function is the solution of the following matrix linear equations:

$$L(\lambda)\Psi = \mu\Psi, \frac{d}{dt}\Psi = A(\lambda)\Psi.$$
 (4)

Generally, BA-function is explicitly written in terms of Riemann's theta functions which are associated with the affine part of some algebraic curve. In our case this curve is hyperelliptic of the genus 3:

$$= \mu^2 \epsilon (\lambda + \lambda^{-1}) + P_2(\mu^2) = 0.$$
 (5)

It is easy to see that K is equivalent to

ĸ

$$\mathbf{K}_{1}: \ \mathbf{w}^{2} = (\xi^{4} - (\mathbf{u}^{2} + \mathbf{u}^{-2})\xi^{2} + 1)(\xi^{4} - (\mathbf{v}^{2} + \mathbf{v}^{-2})\xi^{2} + 1), \tag{6}$$

where u,v are the rational functions of the first integrals of (2) when n=2, these integrals are the coefficients of the polynomial in (5). We point out that K is exactly the spectral curve of the two-particle Toda system associated with the Kac-Moody algebra $D_2^{(1)}$, see [11]. The spectral curve K₁ possesses the dyhedral group of automorphysms and due to this fact the 3-dimensional theta functions, which naturally exist in finite-gap method [10] reduce to one dimensional ones i.e. reduce to elliptic functions. The simplest way to demonstrate this reduction is the following: all Abelian integrals of the first type associated with the curve K₁ are easily related to $\int dx (x + \sqrt{\eta v})$ $[x(x-1)(x-\eta)(x-\nu)(x-\eta\nu)]^{-1/2}$ which after the rank 2 Jacobi reduction goes into elliptic integrals. These results agree with the results of paper [1].

Owing to the conjecture of nonintegrability of (2) the following problem naturally arises: Are there integrable cases of dynamical systems of the type (1) ? The structure of (1) gives us a possibility to find the Lax representation in the following form:

$$\mathbf{L} = \begin{bmatrix} \mathbf{l}_1 & \mathbf{q} \\ \mathbf{r} & -\mathbf{l}_2 \end{bmatrix}, \quad \mathbf{A} = \begin{bmatrix} \mu_1 & \mathbf{x}_1 \\ -\mathbf{x}_2 & \mu_2 \end{bmatrix}$$
(7)

where $l_{1,2}, \mu_{1,2}, q, r, \chi_{1,2}$ are matrices of the type n x n. The Lax representation is equivalent to the following matrix equations:

$$\begin{aligned} \frac{d}{dt} \mathbf{1}_{1} &= [\mathbf{1}_{1}, \mu_{1}] - (q_{2} + x_{1}\mathbf{r}), \quad \frac{d}{dt} \mathbf{1}_{2} &= [\mathbf{1}_{2}, \mu_{2}] - (\mathbf{r}x_{1} + x_{2}\mathbf{q}), \end{aligned} \tag{8} \\ \frac{d}{dt} \mathbf{q} &= \mathbf{q} \ \mu_{2} - \mu_{1}\mathbf{q} + (\mathbf{1}_{1}x_{1} + x_{1}\mathbf{1}_{2}), \\ \frac{d}{dt}\mathbf{r} &= \mathbf{r} \ \mu_{1} - \mu_{2}\mathbf{r} + (\mathbf{1}_{2}x_{2} + x_{2}\mathbf{1}_{1}). \end{aligned} \tag{9}$$

Is it possible to specify the structure of $(1_{\alpha}, \mu_{\alpha}, \chi_{\dot{\alpha}})$ in such a way that (9) reduces to an equation of first order with cubic nonlinearities and (8) reduces to identity? The answer to this question is positive. The simplest construction arises after the following choice of the matrices entering in (8),(9):

$$x_1 = -l_1q - ql_2, \qquad x_2 = l_2r + r_1l_1,$$

$$\mu_1 = -qr - m_1, \qquad \mu_2 = rq - m_2.$$

where $[m_1, l_1] = [m_2, l_2] = 0$, l_1 and l_2 are constant matrices. The
system of equations (9) for q and r in this case has the

form

$$\frac{d}{dt}q=2qrq - [(1_1^2+m_1)q + 21_1q1_2 + q(1_2^2+m_2)],$$
(10)

$$\frac{d}{dt}r=-2rqr + [(1_2^2+m_2)r + 21_2r1_1 + r(1_1^2+m_1)].$$

Using phase transformation it is possible to cancel in (10) terms of the following type $r(l_1^2+m_1)$, $q(l_2^2+m_2)$ put in (10) $m_1=-l_1^2$, $m_2=-l_2^2$. We get: and so we shall

$$\frac{d}{dt}q = 2qrq - 2l_1q l_2, \qquad (11)$$

BVB

 $\frac{d}{dt}r = -2rqr - 2l_2rl_1,$ where the matrices l_1 and l_2 are arbitrary. By similarity transformation it is possible to reduce one of them to the Jordan normal form. It is easy to see that it is impossible to put (10) in the form of the self-trapping equation . Indeed, the sum with cubic terms is absent only when the matrices q and r are diagonal, i.e.

$$\mathbf{q}_{\mathbf{i}\mathbf{j}} = \mathbf{Q}_{\mathbf{i}} \delta_{\mathbf{i}\mathbf{j}}$$
, $\mathbf{r}_{\mathbf{i}\mathbf{j}} = \mathbf{R}_{\mathbf{i}} \delta_{\mathbf{i}\mathbf{j}}$, $\mathbf{i}, \mathbf{j}=1,\ldots,n$ (12)

From (10) we obtain, that the matrices 1,q1, , 1,r1, must be diagonal too. Up to the phase and gauge ambiguities this condition is satisfied only for the two sets of matrices (1_1 , 1_2). Each choice corresponds to the irreducible system of equations, i.e. this system does't split to independent subsystems:

$$(\mathbf{A}) \qquad (\mathbf{1}_1)_{\mathbf{j}\mathbf{k}} = \sqrt{e} \quad \delta_{\mathbf{j},\mathbf{k}+1}, \qquad (\mathbf{1}_2)_{\mathbf{j}\mathbf{k}} = \sqrt{e} \quad \delta_{\mathbf{j},\mathbf{k}+1} \qquad (13)$$

 $(1_{1})_{jk} = \sqrt{\varepsilon} (\delta_{j+1,k} + \lambda \delta_{j,n} \delta_{k,1}),$ (B)

$$(1_2)_{jk} = \sqrt{\epsilon} (\delta_{j,k+1} + \lambda^{-1} \delta_{j+1,k} \delta_{k,n}).$$

The corresponding systems (10) may be written in the following form

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbf{Q}_{j} = 2(\mathbf{Q}_{j}^{2}\mathbf{R}_{j} - \mathbf{e}\mathbf{Q}_{j+1}) , \quad \frac{\mathrm{d}}{\mathrm{d}t}\mathbf{R}_{j} = -2(\mathbf{Q}_{j}\mathbf{R}_{j}^{2} - \mathbf{e}\mathbf{R}_{j-1}).$$
(15)

In the case (A) we have $Q_{n+1}=0, R_0=0$ ("nonperiodic conditions"), and in the case (B) we impose the cyclic conditions $Q_{n+1}=Q_1$, $R_0=R_n$. The Lax matrices in the case (B) contain an explicit dependence on a spectral parameter.

The simplest reduction of the system (15) is obtained by imposing the following cyclic conditions $R_i = i Q_{n+1-i}^*$, $\varepsilon = i \delta$, where δ is a real number.

$$\frac{d}{dt}Q_{j} = i (Q_{j}^{2}Q_{n+1-j}^{*} - \delta Q_{j+1})$$
(16)

The system (16) has a different structure of nonlinear term in comparison with nonintegrable self-trapping equations. Its physical applications (if any) are unknown.

In the case n=2 another reduction of the system (15) is possible $R_1 = i Q_1^*$, $R_2 = i Q_2^*$. This reduction immediately gives the completely integrable system (16) in the so called "periodic" case. When $-\infty < j < \infty$ we may consider the system (15) as a differential-difference equations of the first order. We shall present the cnoidal-type solutions of this system elsewhere.

Now we want to investigate only the finite dimensional case. It is easy to see that the systems (15),(16) are Hamiltonian and the Hamiltonians have the form H= 1/4 tr (L⁴). The later integrals are the part of more general integrals of motion I_{L} = $(1/2k)tr(L^{2k})$.

The leading terms of these integrals have the form $k - \stackrel{T}{\underline{\Gamma}} = Q_{j}^{k} R_{j}^{k}$ i.e. they are polynomials of k-th order of the variables $\{\bar{Q}_j, R_j\}$ which proves that the integrals in the set $\{I_k^-\}$ are functionally independent. Using the standard method (see for example [9]) it is easy to prove that these integrals are in involution with each

(14)

other. This completes the proof of the integrability of the system (15) in the Liouville sense.

For the dynamical systems of type (B) we may write the spectral curve det($L(\lambda) - \mu E$) = 0 in the more suitable form $|W_{ij}|= 0$, where we introduce the notation:

$$W_{ij} = (Q_i R_i + \epsilon Q_i / Q_{i-1} - \mu^2) \delta_{i,j} - \delta_{i,j+1} Q_{i-1} \mu \sqrt{\epsilon} + \mu \delta_{i+1,j} \sqrt{\epsilon} - \sqrt{\epsilon} \mu / \lambda \delta_{i,1} \delta_{j,n} Q_1 / Q_n + \sqrt{\epsilon} \mu \lambda \delta_{i,n} \delta_{j,1}$$

It is easy to express the explicit dependence of the spectral curve on the spectral parameter λ

$$\mathbb{K} : \mu^n \, \varepsilon^{n/2} (\lambda + \lambda^{-1}) + \mathbb{P}_n(\mu^2) = 0$$

where P_n is the polynomial of degree n in μ^2 . Its coefficients constitute the full set of involutive integrals of motion. The genus of the curve equals 2n-1, the number of involutions on K are different for odd and even n

$$n = 2 k , T_1: \lambda \rightarrow -\lambda , T_2: \mu \rightarrow -\mu$$

$$n = 2k-1, T: \lambda \rightarrow -\lambda , \mu \rightarrow -\mu$$

When n=2 besides T_1 , T_2 , an additional involution exists (see the previous text). The system (15) is similar (in the sence of linearization of these systems on the hyperelliptic algebraic curves) to the Adler-van Moerbeke generalized Toda system [11] associated with the Kac-Moody algebra $D_n^{(1)}$. The linearization of these dynamical systems occurs on the Prym varieties of the Jacobeans ($Prym(K_0) \subset Jac(K)$). The structure of the Prym(K_0) will presumably allow us to express the explicit solutions of (15) in terms of the Prym theta functions.

An open problem is the relation of the systems (15), (16) with the periodic and nonperiodic Toda systems [11,12]. In the simplest case n=2 we are able to find the following isomorphism between the system of the type (A) and the usual nonperiodic Toda lattice with the canonical variables (p_1, x_1, p_2, x_2) :

$$Q_1 = p_1 e^{x_1/2}, Q_2 = e^{x_2}, R_1 = e^{-x_1}, R_2 = p_2 e^{-x_2/2}.$$

In our opinion, it is of great interest to find the Lie algebra interpretation of the Lax matrix representation in the case (A) and the Kac-Moody algebra interpretation in the case (B). The investigation of the quasiperiodic dynamics of the system (B), including the construction of the BA-function on the Prym varieties of the spectral curve will be given in a separate paper. We also expect that these dynamical systems are a subclass of a more complicated family of integrable equations of the nonlinear wave interaction type. Their classification is far from being complete.

Acknowledgements.

We are very much indebted to A.V. Mikhailov for many discussions and suggestions.

References

- Eilbeck J.C., Lomdahl P.S., Scott A.C. Physica, 1985, D16, p.319; Kenkre V.M., Tsironis G.P. - Phys.Rev, 1987, B35, p.1473.
- Benney D.J., Newell A.C. Studies in Appl.Math., 1967, 46 p.133.
- 3. Akhmanov S.A., Khokhlov R.V. The problems in nonlinear optics, VINITI.
- 4. Maslov V.P., Uspekhi.Matem.Nauk, 1986, 41, p.19.
- 5. Fordy A., Kulish P., Commun. Math. Phys. 1983, 89, p.427.
- Olshanetsky H.A., Perelomov A.M., Reyman A.G., Semenov-Tian-Shansky M.A., Integrable systems II, (VINITI, Moscow, 1987), p.86
- Fordy A., Wojciechovski S., Marshall I., Phys.Lett. 1986, A13, p.395.
- 8. Dubrovin B.A., Russ.Math.Surveys, 1981, 36, p.11.
- 9. Perelomov A.M., Lett.Math.Phys., 1977, 1, p.531.
- Belokolos E.D., Enol´skii V.Z., Bobenko A.I., Matveev V.B., Uspekhi.Matem.Nauk, 1986, 41, p.3.
- 11. Adler M., van Moerbeke P., Adv. in Math., 1980, 38, p.318.
- 12. Inozemtsev V.I., JINR preprint, 1988, R5-88-235.

Received by Publishing Department

on August 11, 1988.

WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices - in US \$, including the packing and registered postage.

D13-84-63

D13-84-63	Proceedings of the XI International Symposium on Nuclear Electronics. Bratislava, Czechoslovakia, 1983.	12.00
E1,2-84-160	Proceedings of the 1983 JINR-CERN School of Physics. Tabor, Czechoslovakia, 1983.	6.50
D2-84-366	Proceedings of the VII International Conference on the Problems of Quantum Field Theory. Alushta, 1984.	11.00
D1,2-84-599	Proceedings of the VII International Seminar on High Finergy Physics Problems. Dubna, 1984.	12.00
D17-84-850	Proceedings of the III International Symposium on Selected Topics in Statistical Mechanics. Dubna, 1984 (2 volumes).	22.00
	Proceedings of the IX All-Union Conference on Charged Particle Accelerators. Dubna, 1984. (2 volumes)	25.00
D11-85-791	Proceedings of the International Conference on Computer Algebra and Its Applications in Theoretical Physics. Dubna, 1985.	12.00
D13-85-793	Proceedings of the XII International Symposium on Nuclear Electronics, Dubna, 1985.	14.00
D4-85-851	Proceedings of the International School on Nuclear Structure	11.00
D1,2-86-668	Proceedings of the VIII International Seminar on High Energy Physics Problems, Dubna, 1986 (2 volumes)	23.00
D3,4,17-86-747	Proceedings of the V International School on Neutron Physics. Alushta, 1986.	25.00
D9-87-105	Proceedings of the X All-Union Conference on Charged Particle Accelerators. Dubna, 1986 (2 volumes)	25.00
D7-87-68	Proceedings of the International School-Seminar on Heavy Ion Physics. Dubna, 1986.	25.00
D 2-8 7-123	Proceedings of the Conference "Renormalization Group-86". Dubna, 1986.	12.00
D4-87-692	Proceedings of the International Conference on the Theory of Few Body and Quark-Hadronic Systems. Dubna, 1987.	12.00
D2-87-798	Proceedings of the VIII International Conference on the Problems of Quantum Field Theory. Alushta, 1987.	10.00
D14-87-799	Proceedings of the International Symposium on Muon and Pion Interactions with Matter. Dubna, 1987.	13.00
D17-88-95	Proceedings of the IV International Symposium on Selected Topics in Statistical Mechanics. Dubna, 1987.	14.00
01	rders for the above-mentioned books can be sent at the address: Publishing Department, JINR Head Post Office, P.O.Box 79 101000 Moscow, USSR	

Иноземцев В.И., Костов Н.А.

0 новых интегрируемых системах уравнений взаимодействующих нелинейных волн

Найдены две новые гамильтоновы системы уравнения типа $i \frac{d}{dt} \vec{u} = \vec{F}(\vec{u})$

с кубической нелинейностью, обладающие Лаксовым представлением и достаточным набором независимых инволютивных интегралов. Одна из них в случае двух степеней свободы содержит в качестве редукции известное уравнение нелинейных колебаний двухатомных молекул и допускает рассмотрение в рамках общей теории конечнозонного интегрирования.

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1988

Inozemtsev V.I., Kostov N.A.

E5-88-622

New Integrable Systems of Interacting Nonlinear Waves

We obtain two new Hamiltonian systems of the following type $i \frac{d}{dt} \vec{u} = \vec{F}(\vec{u})$ with cubic nonlinearity, which have the Lax representation and appropriate set of first integrals. One of these systems in the two dimensional case after appropriate reduction contains the so called dymer equation and the last equation can be investigated using the finite-gap integration method.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1988