

A-81

16/xi-70

ОБЪЕДИНЕННЫЙ  
ИНСТИТУТ  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ

Дубна



E7 - 5319

A.G. Artukh, V.V. Avdeichikov\*, J. Erö,  
G.F. Gridnev, V.L. Mikheev, V.V. Volkov,  
J. Wilczynski

NEUTRON-RICH ISOTOPES OF LIGHT  
ELEMENTS PRODUCED IN TRANSFER  
REACTIONS WITH HEAVY IONS


1970

8566/2 pr  
A.G. Artukh, V.V. Avdeichikov\*, J. Erö,\*\*  
G.F. Gridnev, V.L. Mikheev, V.V. Volkov,  
J. Wilczynski\*\*\*

**NEUTRON-RICH ISOTOPES OF LIGHT  
ELEMENTS PRODUCED IN TRANSFER  
REACTIONS WITH HEAVY IONS**

Submitted to the Conference on the Properties  
of Nuclei Far from the Region of Beta-Stability  
Leysin, Switzerland, 1970

- 
- \* On leave from the V.G. Khlopin Radium Institute,  
Leningrad, USSR
  - \*\* On leave from the Central Research Institute for Physics,  
Budapest, Hungary
  - \*\*\* On leave from the Institute of Nuclear Physics, Cracow,  
Poland



## 1. Introduction

In the last few years the study of the neutron-rich isotopes of light elements has grown in importance. The discovery of such nuclei as  ${}^8\text{He}$  and  ${}^{11}\text{Li}$  has shown that neutron excess in light nuclei may reach a considerable value. This in turn arose the question of the boundary for the particle stability of nuclides with large neutron excess.

The boundary for the proton-rich nuclei can be estimated quite accurately, since the Coulomb repulsion between protons gives strong limitation for the proton excess. In the case of neutron-rich nuclei the problem is not so clear. Different theoretical approaches give quite different results.

The limitations for the neutron excess follow mainly from the Pauli's exclusion principle. For a nucleus with large neutron excess more single particle levels of high energy must be occupied than for the nucleus with the same  $A$  but with  $N-Z = 0$ . Some of these high lying levels may turn out to be unbound.

However, the application of the shell model to light nuclei with large neutron excess arises serious doubts. Recently, the multidimensional harmonic (K-harmonic) method<sup>/1,2/</sup> was used for binding energy calculations in a region of light nuclei. In the framework of this method the ground state solution of the Schrodinger equation for the system of nucleons can be obtained with a good accuracy. To do that, the nucleon-nucleon force must, of course, be known sufficiently well. For the neutron-rich nuclides the knowledge of the  $V^{(88)}$  potential (it is the potential for nucleon-nucleon interaction in the  $S = 1, T = 1$  state) is especially important. However, the magnitude of  $V^{(88)}$  cannot be reliably determined from the experiments on nucleon-nucleon scattering.

As has been pointed out in ref.<sup>/2/</sup>, a small contribution of attractive  $V^{(88)}$  potential leads to the possibility of the existence of  $^{10}\text{He}$  and even  $^{22}\text{He}$ . This statement may, of course, be inverted: studying the properties of light nuclides with large neutron excess one can obtain information on nuclear forces acting between the neutrons. Thus, experiments in this field have a fundamental character.








## 2. Production and Identification of Neutron-Rich Isotopes of Light Elements

Several methods have been used until now for production of light neutron-rich nuclides: spontaneous fission of  $^{252}\text{Cf}$ , fission of  $^{235}\text{U}$  and  $^{239}\text{Pu}$  induced by thermal neutrons, reactions with high-energy protons, and finally, reactions of negative pion capture and recharging. In our experiments the neutron-rich





isotopes of a number of light elements were produced in transfer reactions with heavy ions. These reactions usually occur in two-nuclei collision close to tangential ones. Coulomb and centrifugal barriers due to a large  $Z$  and the angular momentum of relative motion prevent the fusion of two nuclei into the compound nucleus. However the intensive interaction due to overlapping of nuclear surfaces ensures the transfer of a considerable number of nucleons between nuclei. Large variety of the multi-nucleon transfer reactions leads to the production of these isotopes: the proton stripping (projectile - x protons), the neutron pick-up (projectile + y neutrons) and the nucleon exchange reactions (projectile - x protons + y neutrons). Several types of reactions resulting in the production of the neutron-rich nuclides observed in our experiments are listed in Table 1.

Table 1  
Types of reactions resulting in the production of neutron-rich isotopes of light elements









### 1) neutron pick-up



### 2) proton stripping



3) nucleon exchange



The main experimental difficulty in the study of the direct reactions with heavy ions is the problem of identification of the reaction products. The radiochemical method is not suitable for the detection of nuclides with large neutron excess since they decay in a fraction of a second. Also, the  $dE/dx, E$  technique alone is not good enough because of an insufficient isotope resolution.

For an unambiguous identification of the projectile transformation products in transfer reactions we use a heavy-ion identification method which combines a magnetic spectrometer with the  $dE/dx, E$  technique<sup>/3/</sup> (see Fig. 1). The reaction products pass the magnetic spectrometer and are detected in a counter telescope placed in the focal plane of the spectrometer. The telescope consists of two semiconductor detectors: a thin  $\Delta E$  detector and a thick  $E - \Delta E$  detector.

The pulses from both detectors, after amplification, are sent to a two-dimensional, 64x64 channel pulse-height analyser. Since only the reaction products with discrete energy values  $E = \text{const } Z_{\text{ion}}^2 / M$  can pass the magnetic spectrometer, the values of  $\Delta E$  and  $E - \Delta E$  pulses are also discrete, and therefore, the unambiguous identifi-

cation of each isotope in two-dimensional spectrum is possible (an example of such a two-dimensional spectrum is shown in Fig. 2).

The method allows an unambiguous identification of all the isotopes of light elements emitted from the target with energies higher than 2-3 MeV per nucleon. In comparison with the standard  $dE/dx, E$  technique our method gives much better separation of neighbouring isotopes. Moreover, the method allows to remove the elastically scattered particles from the reaction product beam before it gets into the telescope. This gives the possibility to detect the reaction products formed with very small cross sections.

### 3. New Isotopes of Carbon, Nitrogen, Oxygen, Fluorine and Neon Produced in Transfer Reactions with Heavy Ions

The results obtained earlier<sup>/4/</sup> as well as the data from some additional experiments show that the cross sections for the reactions: (projectile - x protons), (projectile + y neutrons) and (projectile - x protons + y neutrons), leading to the production of neutron-rich isotopes of light elements are large when heavy elements are used as targets. Therefore, the target of  $^{282}\text{Th}$  was used in our experiments. It was bombarded with  $^{18}\text{O}$ ,  $^{22}\text{Ne}$ ,  $^{15}\text{N}$  and  $^{11}\text{B}$  ions.

Experiments were performed with the 310-cm heavy ion cyclotron of Nuclear Reaction Laboratory, JINR, at Dubna.

The reaction products were detected at an angle corresponding to the surface collision trajectory. The angular distributions in these reactions should have the wide maxima near this angle (see, for example, refs.<sup>/5,6/</sup>).

At a given magnetic rigidity BR all the reaction products could be detected only within narrow energy intervals. Measuring the yields of nuclides at different values of BR the energy spectra of the reaction products could be obtained.

As an example, the dependence of the yields of oxygen isotopes on the magnetic rigidity BR is shown in Fig. 3 for the case in which a thorium target was bombarded with 122 MeV  $^{18}\text{O}$  ions. The curves of the isotope yields have been obtained by summing along  $\Delta E$  axis the number of pulses belonging to the isotopes of the given element in the fixed charged state.

In our first experiment in bombardment of  $^{232}\text{Th}$  with  $^{18}\text{O}$  ions we expected to produce the  $^{22}\text{O}$  isotope in the pick-up reaction of 4 neutrons. The experiment was performed for magnetic rigidity at which the heaviest isotopes of oxygen were observed with maximum yields. The results of the 6-hour irradiation are shown in Fig. 4. Apart from a number of already known nuclides, three new nuclides:  $^{22}\text{O}$ ,  $^{20}\text{N}$  and  $^{18}\text{C}$  have been obtained. It should be noted that  $^{20}\text{N}$  and  $^{18}\text{C}$  nuclides were produced in the nucleon exchange reactions: (projectile - 1 proton + 3 neutrons) and (projectile - 2 protons + 2 neutrons), respectively. Thus, the nucleon exchange reactions turned out to be a very effective way of producing the nuclides with large neutron excess.

In the next series of experiments the thorium target was bombarded with 174 MeV  $^{22}\text{Ne}$  ions. In these experiments four new nuclides:  $^{28}\text{F}$ ,  $^{24}\text{F}$ ,  $^{25}\text{Ne}$  and  $^{26}\text{Ne}$  were obtained (Fig. 5). Finally, after increasing the efficiency of the detecting system and increasing the time of irradiation we succeeded to obtain subsequent four new nuclides:  $^{31}\text{N}$ ,  $^{28}\text{O}$ ,  $^{24}\text{O}$  and  $^{25}\text{F}$  (Fig. 6). In order

to reduce fluorine light isotope background the oxygen nuclei were detected in the  $^{+7}\text{O}$  charge state. The  $^{+7}\text{O}$  yield was somewhat reduced as compared to that of  $^{+8}\text{O}$  but the yield of the fluorine light isotopes was depressed strongly. Fig. 7 shows magnetic rigidity and the production cross sections of some heavy isotopes of oxygen in bombarding  $^{232}\text{Th}$  by  $^{22}\text{Ne}$  ions. These data were used to choose magnetic rigidity in detecting still heavier isotopes.

In bombarding the thorium target with 145 MeV  $^{16}\text{N}$  and 87 MeV  $^{11}\text{B}$  ions all the heaviest known isotopes of helium, lithium, beryllium and boron ( $^8\text{He}$ ,  $^{11}\text{Li}$ ,  $^{12}\text{Be}$ ,  $^{15}\text{B}$ ) were produced.

A list of eleven new nuclides produced in transfer reactions with heavy ions, together with the heaviest isotopes of light elements known earlier is presented in Table 2.

Table 2

Isotopes of light elements with large neutron excess produced in transfer reactions with heavy ions

a) known isotopes

$^8\text{He}$ ,  $^{11}\text{Li}$ ,  $^{12}\text{Be}$ ,  $^{15}\text{B}$ ,  $^{17}\text{C}$ ,  $^{19}\text{N}$ ,  $^{21}\text{O}$ ,  $^{22}\text{F}$ ,  $^{24}\text{Ne}$

b) new isotopes

$^{18}\text{C}$ ,  $^{20}\text{N}$ ,  $^{22}\text{O}$ ,  $^{28}\text{F}$ ,  $^{25}\text{Ne}$

$^{21}\text{N}$ ,  $^{28}\text{O}$ ,  $^{24}\text{F}$ ,  $^{26}\text{Ne}$

$^{24}\text{O}$ ,  $^{25}\text{F}$

It should be noted that the yields of neutron-rich isotopes in reactions with heavy ions are much higher than in reactions with high-energy protons. This gives possibility of spectroscopic study of nuclei with large neutron excess. In the Laboratory of Nuclear Reactions, JINR, the experiments in this field were initiated with the use of the magnetic mass-separator operating on-line with the heavy ion cyclotron.

#### 4. The Boundary of Particle-Stability for Light Nuclides with Large Neutron Excess, Evidence for Particle-Instability of $^{13}\text{Be}$ and $^{14}\text{Be}$

It is interesting to compare the present list of known nuclides with the theoretical predictions concerning the boundary of particle stability (Fig. 8). It can be seen from Fig. 8 that according to Garvey and Kelson<sup>/7/</sup> and Vinogradov and Nemirovsky<sup>/8/</sup> the isotopes of H, He, Li, Be and B which are heavier than already known nuclides ( $^3\text{H}$ ,  $^8\text{He}$ ,  $^{11}\text{Li}$ ,  $^{12}\text{Be}$  and  $^{15}\text{B}$ ), should be particle-unstable.

However, the question of the existence of  $^{10}\text{He}$  and  $^{14}\text{Be}$  has still been unsettled. The calculations of Baz, Demin and Zhukov<sup>/2/</sup>, performed in the framework of the K-harmonic method, did not exclude the possibility of particle stability of  $^{10}\text{He}$ . As for  $^{14}\text{Be}$ , only Garvey and Kelson<sup>/7/</sup> gave a definite answer. They predict that the neutron binding energy in  $^{18}\text{Be}$  is equal to -2.7 MeV and the binding energy of two neutrons in  $^{14}\text{Be}$  is equal to -2.4 MeV. According to Vorobiev et al.<sup>/9/</sup> these quantities are -0.1 MeV and +1.9 MeV, respectively. The analysis in the framework of the shell model<sup>/8/</sup> did not provide an unambiguous answer.

Some years ago Poskanzer et al.<sup>/10/</sup> reported their experimental results suggested that  $^{18}\text{Be}$  is probably particle-unstable.

We undertook experiments with the aim to give an answer to the question of the existence of  $^{10}\text{He}$  and  $^{14}\text{Be}$ . The experiments on  $^{10}\text{He}$  are under way. Here we report only the results of the experiment on  $^{14}\text{Be}$ .

The experiment can give a definite answer to the above question because one can estimate the expected yield of a given nuclide from the cross section systematics for multi-nucleon transfer reactions, as reported by us elsewhere<sup>/11/</sup>. The cross sections for the nucleon stripping - and nucleon exchange reactions (projectile - x protons  $\pm$  y neutrons) depend exponentially on the Q-values, calculated for the ground state masses of the reaction products. As an example of this dependence the data on cross sections for production of a large number of nuclides in the bombardment of a thorium with 137 MeV  $^{18}\text{O}$  ions are shown in Fig. 9.

The search for  $^{14}\text{Be}$  was performed in the  $^{16}\text{Ne} + ^{282}\text{Th}$  reaction at the energy of  $^{15}\text{N}$  ions equal to 145 MeV. Magnetic rigidity was chosen according to the energy spectra for some reactions (Fig. 10). The values of differential cross sections  $(d\sigma/d\Omega)_{40^\circ}$  for the production of carbon, boron, beryllium and lithium isotopes in this reaction are shown in Fig. 11. The extrapolated line for beryllium isotopes (shown in Fig. 11) should not contain any significant error, since the neighbouring lines are almost parallel and fit well the experimental points in a wide range of the cross section values.

In our experiment we detected about 20000 events of  $^{12}\text{Be}$ , whereas no effect due to the  $^{13}\text{Be}$  and  $^{14}\text{Be}$  ions was observed. The background, defining the upper limit of the  $^{13}\text{Be}$  and  $^{14}\text{Be}$  yields was about 3 events. Assuming that  $^{13}\text{Be}$  and  $^{14}\text{Be}$  are extremely weakly bound the expected yields of  $^{13}\text{Be}$  and  $^{14}\text{Be}$  should be about 550 events and 30 events, respectively. In the case of the binding energy of two neutrons in  $^{14}\text{Be}$  equal to 1.9 MeV, as suggested by Vorobiev et al.<sup>[9]</sup>, the expected yield of  $^{14}\text{Be}$  should be even larger (about 70 events).

Fig. 11 shows that even so weakly bound nuclei as  $^{14}\text{B}$  and  $^{11}\text{Li}$  are produced with yields consistent with our systematics. Therefore, we can conclude that in all probability the  $^{13}\text{Be}$  and  $^{14}\text{Be}$  nuclei are particle-unstable.

The problem of the particle-stability of  $^{10}\text{He}$  is now under study in a similar experiment.

The authors express their deep gratitude to Academician G.N. Flerov for his support and stimulating interest in our work. Thanks are also due to B.A. Zager and the cyclotron crew for their cooperation and to V.I. Vakatov and L.P. Chelnokov for help with the electronic apparatus.

#### R e f e r e n c e s

1. A.I. Baz and M.V. Zhukov. Kurchatov Institute, report IAE-1883, Moscow, 1969.
2. A.I. Baz, V.F. Demin and M.V. Zhukov. *Yadernaya Fiz.*, 9, 1184 (1969).
3. A.G. Artukh, V.V. Avdeichikov, J. Erö, G.F. Gridnev, V.L. Mikheev and V.V. Volkov. *Nucl.Instr. and Methods.*, 83, 72 (1970).

4. V.V. Volkov, L. Pomorski, J. Tys and G.N. Flerov. Proc. of the Conf. on Direct Interactions and Nuclear Reactions Mechanisms (ed. by E. Clemental and C. Villi) Gordon and Breach, New York - London, 1963, p. 994.
5. H. Kumpf and E.D. Donetz. *JETP (USSR)*, 44, 798 (1963).
6. V.V. Volkov, G.F. Gridnev, G.N. Zorin and L.P. Chelnokov. *Nucl. Phys.*, A126, 1 (1969).
7. G.T. Garvey and I. Kelson. *Phys.Rev.Lett.*, 16, 197 (1966).
8. B.N. Vinogradov, P.E. Nemirovsky. *Yadernaya Fiz.*, 10, 505 (1969).
9. A.A. Vorobiev, V.T. Gratchev, Yu.K. Zalite, I.A. Kondurov, A.M. Nikitin and D.M. Seleverstov. Report FTI-232, Leningrad, 1969.
10. A.M. Poskanzer, S.W. Cospers, E.K. Hyde and J. Cerny. *Phys. Rev.Lett.*, 17, 1271 (1966).
11. A.G. Artukh, V.V. Avdeichikov, J. Erö, G.F. Gridnev, V.L. Mikheev, V.V. Volkov and J. Wilczynski. Preprint JINR, E7-5241, Dubna, 1970. To be published in Nuclear Physics.

Received by Publishing Department  
on August 11, 1970.






Fig. 1. Comparison of the methods of particle identification: (A) the standard  $dE/dx, E$  method; (B) the magnetic analysis with the measurements of energy; (C) the combination of the magnetic analysis with the  $dE/dx, E$  method.




Fig. 2. Two-dimensional ( $\Delta E, E-\Delta E$ ) spectrum obtained in bombardment of the  $^{213}\text{Th}$  target with  $^{174}\text{MeV } ^{32}\text{Ne}$  ions. The area of each circle is proportional to the square root of the number of counts.




Fig. 3. Yields of oxygen isotopes from the  $^{282}\text{Th} + ^{18}\text{O}$  reaction ( $E(^{18}\text{O}) = 122$  MeV) at four different values of the magnetic rigidity BR. a) BR = 7.87 kGs.m, the ion flux through the target  $J = 2.3 \cdot 10^{14}$  particles; b) BR = 8.03 kGs.m,  $J = 1.9 \cdot 10^{15}$  particles; c) BR = 8.10 kGs.m,  $J = 1.9 \cdot 10^{15}$  particles; d) BR = 8.17 kGs.m,  $J = 1.4 \cdot 10^{16}$  particles.




Fig. 4. Yields of oxygen, nitrogen, carbon, boron and beryllium isotopes from the  $^{282}\text{Th} + ^{18}\text{O}$  reaction.  $E(^{18}\text{O}) = 122$  MeV; BR = 8.17 kGs.m; the  $^{18}\text{O}$  flux through the target  $J = 1.4 \cdot 10^{16}$  particles.





Fig. 5. Yields of neon, fluorine, oxygen, nitrogen and carbon isotopes from the  $^{22}\text{Ne} + ^{232}\text{Th}$  reaction.  $E(^{22}\text{Ne}) = 174$  MeV. (A) BR = 9.92 kGs.m, the  $^{22}\text{Ne}$  flux through the target  $J = 9.2 \cdot 10^{15}$  particles; (B) BR = 9.74 kGs.m,  $J = 2.3 \cdot 10^{15}$  particles.




Fig. 6. Yields of neon, fluorine, oxygen, nitrogen and carbon isotopes from the  $^{22}\text{Ne} + ^{232}\text{Th}$  reaction.  $E(^{22}\text{Ne}) = 174$  MeV; BR = 10.4 kGs.m; the  $^{22}\text{Ne}$  flux through the target  $J = 5.1 \cdot 10^{16}$  particles.




Fig. 7. Yields of <sup>20</sup>O, <sup>21</sup>O and <sup>22</sup>O isotopes from the <sup>22</sup>Ne + <sup>232</sup>Th reaction as a function of magnetic rigidity BR (on the left) and as a function of the outgoing particle energy (on the right). The arrows point the predicted outgoing particle energies for reactions leading to the ground states of the both reaction products.




Fig. 8. Theoretical predictions concerning the boundary of particle-stability for light nuclides with large neutron excess. The solid line shows the predictions of Garvey and Kelson<sup>7/</sup>, the dashed line is shown according to Vinogradov and Nemirovsky<sup>8/</sup>.




Fig. 9. Differential cross sections  $(\frac{d\sigma}{d\Omega})_{40^\circ}$  for production of Be, B, C and N isotopes in the  $^{232}\text{Th} + ^{16}\text{O}$  reaction as function of  $Q_{e.e.}$ .




Fig. 10. Energy spectra of  $^{14}\text{C}$ ,  $^{15}\text{C}$ ,  $^{16}\text{C}$ ,  $^{18}\text{B}$ ,  $^{14}\text{B}$ ,  $^{15}\text{B}$  and  $^{12}\text{Be}$  produced in the nucleon exchange reactions (projectile - x protons + y neutrons) in bombardment of  $^{232}\text{Th}$  target with  $^{15}\text{N}$   $^{145}\text{MeV}$   $^{15}\text{N}$  ions. The dashed curves show the predicted spectra of  $^{13}\text{Be}$  and  $^{14}\text{Be}$ . The arrows denote the energy values corresponding to the formation of both reaction products in the ground states.




Fig. 11. Differential cross sections  $(\frac{d\sigma}{d\Omega})_{40^\circ}$  for production of Li, Be, B and C isotopes in the  $^{232}\text{Th} + ^{15}\text{N}$  reaction as function of  $Q_{G.G.}$ . The  $Q_{G.G.}$ -value for  $^{13}\text{Be}$  corresponds to the particle-stability threshold ( $E_n=0$ ). For  $^{14}\text{Be}$  the left and right arrows indicate the  $Q$ -values for  $E_{n_1} = +1.9$  MeV and  $E_{n_1} = 0$ , respectively.