

ON SOLVING THE PROBLEM OF HEAVY ION COLLISIONS IN AN OPTICAL MODEL

S. Vinitzky

Joint Institute for Nuclear Research

E-mail: vinitzky@theor.jinr.ru

We present algorithm implements the solution of the inverse problem, i.e., calculates the unknown coupling constant $g(E)$ and scattering matrix $S(g(E), E)$ from condition $|S(g(E), E)|^2 = 1 - |T(E)|^2$ by means of the secant method. The required amplitudes of transmission $T(E)$ and reflection $R(E)$ subject also to the condition $|R(E)|^2 = 1 - |T(E)|^2$ of the model with incoming wave boundary conditions (IWBCs) are previously calculated by the standard MAPLE implemented KANTBP 4M program. The algorithm provides a one-to-one correspondence between the OM with a complex-valued potential and the model of IWBCs with a real-valued potential.

The efficiency of the proposed approach is shown by solving numerically the scattering problem and calculating the reference fusion cross section for a pair of heavy ions $^{16}\text{O} + ^{144}\text{Sm}$ in the single-channel approximation of the close-coupling method.

A.A. Gusev, O. Chuluunbaatar, V.L. Derbov, R.G. Nazmitdinov, S.I. Vinitzky, P.W. Wen, C.J. Lin, H. M. Jia, L. L. Hai, Symbolic-numerical algorithm for solving the problem of heavy ion collisions in an optical model with a complex potential, Lecture Notes in Computer Science 14139, pp. 128–140 (2023).